

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
15 January 2004 (15.01.2004)

PCT

(10) International Publication Number  
**WO 2004/005482 A2**

(51) International Patent Classification<sup>7</sup>: C12N  
(21) International Application Number: PCT/US2003/021339  
(22) International Filing Date: 8 July 2003 (08.07.2003)  
(25) Filing Language: English  
(26) Publication Language: English  
(30) Priority Data:  
60/393,756 8 July 2002 (08.07.2002) US  
60/396,627 19 July 2002 (19.07.2002) US  
60/417,172 10 October 2002 (10.10.2002) US

(71) Applicant: INVITROGEN CORPORATION [US/US];  
1600 Faraday Avenue, Carlsbad, CA 92008 (US).

(72) Inventor: BENNETT, Robert, P.; 1269 Rainbow Ridge  
Lane, Encinitas, CA 92024 (US).

(74) Agents: ESMOND, Robert, W. et al.; Sterne, Kessler,  
Goldstein & Fox, P.L.L.C., 1100 New York Avenue, Wash-  
ington, DC 20005 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND COMPOSITIONS FOR THE PRODUCTION, IDENTIFICATION AND PURIFICATION OF FU-  
SION PROTEINS

(57) Abstract: The present invention provides compositions and methods for producing fusion proteins that comprise an amino acid sequence tag. The amino acid sequence tag may be an amino acid sequence that is capable of being post-translationally modified; for example, the amino acid sequence may be an amino acid sequence that is capable of being biotinylated. The amino acid sequence tag may also be an amino acid sequence that is recognized by an antibody (or fragment thereof) or other specific interacting reagent. The invention includes isolated nucleic acid molecules comprising one or more nucleic acid sequences which encode an amino acid sequence tag. The nucleic acid molecules of the invention may also comprise one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases. The nucleic acid molecules of the invention can be used in recombinational cloning and/or topoisomerase-mediated cloning methods in order to produce polynucleotide constructs which encode fusion proteins that comprise an amino acid sequence tag. Also provided are host cells, kits and compositions comprising the nucleic acid molecules of the invention.



WO 2004/005482 A2

## METHODS AND COMPOSITIONS FOR THE PRODUCTION, IDENTIFICATION AND PURIFICATION OF FUSION PROTEINS

### BACKGROUND OF THE INVENTION

#### Field of the Invention

[0001] The present invention relates to compositions and methods for producing fusion proteins. More specifically, the invention relates to compositions and methods for producing fusion proteins that comprise an amino acid sequence tag. Exemplary amino acid sequence tags include amino acid sequences that are capable of being post-translationally modified, and amino acid sequences that are capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent.

[0002] The invention relates to nucleic acid molecules that can be used in recombinational cloning methods and/or topoisomerase-mediated cloning methods to produce polynucleotide constructs that encode fusion proteins, *e.g.*, fusion proteins that comprise one or more amino acid sequence tags. The invention also relates to methods for producing fusion proteins in a variety of prokaryotic and eukaryotic cell types. The invention also relates to methods for identifying and purifying fusion proteins by utilizing, *e.g.*, binding molecules and compositions that bind specifically to the fusion protein.

#### Related Art

[0003] Many areas of biotechnology and molecular biology rely on the production and purification of recombinant proteins. When recombinant proteins are produced *in vivo* they are generally produced in addition to a wide variety of endogenous proteins and other macromolecules in a host cell. Various strategies are employed to isolate and/or identify recombinant proteins from the cellular milieu. One strategy is to produce a fusion protein which comprises the protein of interest joined to an amino acid sequence tag.

[0004] When a fusion protein is produced that comprises a tag that is capable of being post-translationally modified, the post-translational modification can

-2-

be exploited to isolate or identify the fusion protein, especially when (a) very few or no endogenous proteins or molecules contain the same post-translational modification in the host cell, and (b) a molecule is available which is capable of physically interacting with the post-translationally modified protein.

[0005] One particular post-translational modification that has been used to isolate and/or identify recombinant fusion proteins is biotinylation. For instance, a fusion protein can be produced which comprises a protein of interest joined to an amino acid sequence to which a biotin moiety can be covalently bound. The biotinylation reaction will occur *in vivo*, i.e., in the host cell. The biotinylated fusion protein can then be isolated from the endogenous components of the host cell by providing a molecule that interacts specifically with the biotin moiety. Usually, the biotin-interacting molecule will be bound to a bead or other solid support which can be easily separated from the rest of the cellular components.

[0006] Amino acid sequences which are capable of being biotinylated include, for example, a domain the 1.3S subunit of *Propionibacterium shermanii* transcarboxylase (PSTCD) that is naturally biotinylated at lysine 89 of the domain. (Cronan, J.E., *J. Biol. Chem.* 265:10327-10333 (1990); Murtif, V.L., *et al.*, *Proc. Natl. Acad. Sci. USA* 82:5617-5621 (1985)). Another example is a 72 amino acid peptide derived from the C-terminus (amino acids 524-595) of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit. (Schwarz, E. *et al.*, *J. Biol. Chem.* 263:9640-9645 (1988)). Fusion proteins containing biotinylation domains have been shown to be biotinylated by endogenous biotinylation components in bacteria, yeast and mammalian cells. (Cronan, J.E., *J. Biol. Chem.* 265:10327-10333 (1990); Jank, M.M. *et al.*, *Protein Expr. Purif.* 17:123-127 (1999); Parrott, M.B. and Barry, M.A., *Biochem. Biophys. Res. Comm.* 281:993-1000 (2001); Parrott, M.B. and Barry, M.A., *Molecular Therapy* 1:96-104 (2000); U.S. Patent No. 5,252,466 and references cited therein).

[0007] Avidin has been shown to interact very strongly with biotin. The non-covalent interaction between avidin and biotin represents one of the strongest and most specific interactions commonly used in molecular biology. The interaction between avidin and biotin is estimated to have an affinity coefficient of  $10^{-14}$  to  $10^{-15}$ , which is several orders of magnitude greater than a typical antibody-antigen interaction. (Rosano, C. *et al.*, *Biomol. Eng.* 16:5-12 (1999); Green, N.M., *Methods Enzymol.* 184:51-67 (1990); Airenne, K.J. *et al.*, *Protein Expr. Purif.* 17:139-145 (1999); Wilchek, M. and Bayer, E.A., *Methods Enzymol.* 184:5-13 (1990)). Avidin analogs, including streptavidin are also available for specifically interacting with biotin.

[0008] As an alternative to producing a protein or polypeptide that is capable of being post-translationally modified, it is sometimes useful to produce a fusion protein that comprises an amino acid sequence that is identifiable by particular reagents, including, *e.g.*, antibodies (or fragments thereof) or other binding compounds that can recognize certain polypeptides or amino acid sequences.

[0009] In order to produce a recombinant fusion protein that comprises a particular amino acid sequence tag, a nucleic acid molecule must first be constructed which encodes the desired fusion protein. The construction of the recombinant nucleic acid molecule will generally involve the attachment of at least two individual nucleotide sequences: (1) a sequence encoding the protein of interest, and (2) a sequence encoding an amino acid sequence tag.

[0010] Multiple nucleic acid sequences can be joined using conventional *in vitro* cloning methods which employ restriction endonucleases and DNA ligation enzymes. More rapid and efficient methods are available, however, which involve site-specific recombination and/or topoisomerase-mediated joining of nucleic acid sequences. Recombinational and topoisomerase-mediated cloning methods have been described in detail elsewhere. (Hartley, J.L., *et al.*, *Genome Res.* 10:1788-1795 (2000); Shuman, S., *J. Biol. Chem.* 269:32678-32684 (1994); Shuman, S., *Proc. Natl. Acad. Sci. USA* 88:10104-10108 (1991); U.S. Patent Nos. 5,851,808, 5,888,732, 6,143,557, 6,171,861,

6,270,969, 6,277,608 and 6,410,317; and commonly owned, co-pending U.S. Patent Application No. 10/005,876 (filed 12/07/01)).

[0011] Briefly, recombinational cloning, specifically the Gateway™ Cloning System (available from Invitrogen Corporation), utilizes vectors that contain at least one and preferably at least two different site-specific recombination sites based on the bacteriophage lambda system (e. g., *att1* and *att2*) that are mutated from the wild type (*att0*) sites. Each mutated site has a unique specificity for its cognate partner *att* site of the same type (for example *attB1* with *attP1*, or *attL1* with *attR1*) and will not cross-react with recombination sites of the other mutant type or with the wild-type *att0* site. Nucleic acid fragments flanked by recombination sites are cloned and subcloned using the Gateway™ system by replacing a selectable marker (for example, *ccdB*) flanked by *att* sites on the recipient plasmid molecule, sometimes termed the Destination Vector. Desired clones are then selected by transformation of a *ccdB* sensitive host strain and positive selection for a marker on the recipient molecule. Similar strategies for negative selection (e.g., use of toxic genes) can be used in other organisms such as thymidine kinase (TK) in mammals and insects. Other recombinational cloning systems are available such as, e.g., Echo™ (Invitrogen Corporation) and Creator (Clontech).

[0012] Topoisomerase cloning can be used to generate a double-stranded recombinant nucleic acid molecule covalently linked in one strand. This method can be performed by contacting a first nucleic acid molecule which has a site-specific topoisomerase recognition site (e.g., a type IA or a type II topoisomerase recognition site), or a cleavage product thereof, at a 5' or 3' terminus, with a second (or other) nucleic acid molecule, and optionally, a topoisomerase (e.g., a type IA, type IB, and/or type II topoisomerase), such that the second nucleotide sequence can be covalently attached to the first nucleotide sequence. Topoisomerase cloning can also be used to generate a double-stranded recombinant nucleic acid molecule covalently linked in both strands. This method can be performed, for example, by contacting a first nucleic acid molecule having a first end and a second end, wherein, at the first

-5-

end or second end or both, the first nucleic acid molecule has a topoisomerase recognition site (or cleavage product thereof) at or near the 3' terminus; at least a second nucleic acid molecule having a first end and a second end, wherein, at the first end or second end or both, the at least second double stranded nucleotide sequence has a topoisomerase recognition site (or cleavage product thereof) at or near a 3' terminus; and at least one site specific topoisomerase (e.g., a type IA and/or a type IB topoisomerase), under conditions such that all components are in contact and the topoisomerase can effect its activity. A covalently linked double-stranded recombinant nucleic acid by this method is characterized, in part, in that it does not contain a nick in either strand at the position where the nucleic acid molecules are joined. The method may be performed by contacting a first nucleic acid molecule and a second (or other) nucleic acid molecule, each of which has a topoisomerase recognition site, or a cleavage product thereof, at the 3' termini or at the 5' termini of two ends to be covalently linked. Alternatively, the method can be performed by contacting a first nucleic acid molecule having a topoisomerase recognition site, or cleavage product thereof, at the 5' terminus and the 3' terminus of at least one end, and a second (or other) nucleic acid molecule having a 3' hydroxyl group and a 5' hydroxyl group at the end to be linked to the end of the first nucleic acid molecule containing the recognition sites. Topoisomerase cloning methods can be performed using any number of nucleic acid molecules having various combinations of termini and ends.

[0013] Cloning schemes are also available which use both recombinational cloning and topoisomerase cloning methods. Such methods may involve first joining two nucleic acid sequences using recombinational cloning to create a product nucleic acid molecule, followed by joining the product nucleic acid molecule to another nucleic acid molecule using topoisomerase cloning. Conversely, two nucleic acid molecules may be joined, first, by using topoisomerase cloning to create a product nucleic acid molecule, followed by joining the product nucleic acid molecule to another nucleic acid molecule using recombinational cloning.

[0014] Recombinational cloning methods, topoisomerase cloning methods, and combinations thereof, heretofore have not been described in the art for producing nucleic acid constructs that encode fusion proteins that comprise one or more amino acid sequence tags. Accordingly, a need exists in the art for rapid and efficient compositions and methods that enable the production of nucleic acid molecules which encode fusion proteins.

### BRIEF SUMMARY OF THE INVENTION

[0015] The present invention satisfies the aforementioned need in the art by providing compositions and methods for producing fusion proteins which comprise one or more amino acid sequences of interest and one or more amino acid sequence tags. An "amino acid sequence tag," as used herein, includes, *e.g.*, amino acid sequences that are capable of being post-translationally modified, and/or amino acid sequences that are capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent.

[0016] The invention includes isolated nucleic acid molecules comprising one or more nucleic acid sequences which encode an amino acid sequence tag. The isolated nucleic acid molecules of the invention may further comprise one or more recombination sites. Alternatively or additionally, the isolated nucleic acid molecules of the invention may further comprise one or more topoisomerase recognition sites and/or one or more topoisomerases. Thus, in certain embodiments, the invention includes isolated nucleic acid molecules comprising: (a) one or more recombination sites; (b) one or more topoisomerase recognition sites and/or one or more topoisomerases; and (c) one or more nucleic acid sequences which encode an amino acid sequence tag.

[0017] In addition to the aforementioned elements, the nucleic acid molecules of the invention may further comprise additional elements. Exemplary additional elements that may be included within the nucleic acid molecules of the invention include, *e.g.*, one or more promoters, one or more operators, one or more enhancers, one or more ribosome binding sites, one or more initiation codons, one or more nucleic acid sequences that encodes an amino acid



sequence that is capable of being cleaved by one or more proteases, one or more nucleic acid sequences of interest (*e.g.*, one or more nucleic acid sequences that encode one or more proteins or polypeptides of interest), one or more polyadenylation signals and/or one or more transcription termination regions. As understood by those skilled in the art, other elements may be included within the nucleic acid molecules of the invention depending on the circumstances under which the nucleic acids may be used.

[0018] In a preferred embodiment, the elements of the isolated nucleic acid molecules of the invention are arranged relative to one another such that a nucleic acid sequence of interest can be attached to the nucleic acid molecules of the invention, thereby producing a polynucleotide construct that encodes a fusion protein, the fusion protein comprising: (i) an amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest. The fusion protein may be, *e.g.*, an N-terminal fusion protein (*e.g.*, wherein an amino acid sequence tag is covalently attached at or near the N-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest). The fusion protein may also be, *e.g.*, a C-terminal fusion protein (*e.g.*, wherein an amino acid sequence tag is covalently attached at or near the C-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest). The fusion protein may also be, *e.g.*, an N-terminal and C-terminal fusion protein (*e.g.*, wherein an amino acid sequence tag is covalently attached at or near the N-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest and an amino acid sequence tag is covalently attached at or near the C-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest).

[0019] The invention also includes nucleic acid molecules that are created following the attachment of a nucleic acid sequence of interest to a nucleic acid molecule comprising: (a) a nucleic acid sequence that encodes an amino acid sequence tag; and/or (b) one or more recombination sites; and/or (c) one or more topoisomerase recognition sites and/or one or more topoisomerases.

[0020] In order to produce a polynucleotide sequence that encodes a fusion protein that comprises one or more amino acid sequence tags, a nucleic acid

sequence of interest may, for example, be inserted at or within 20 nucleotides of said one or more recombination sites. The nucleic acid sequence may also be inserted at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotides of the position of said one or more topoisomerases in order to produce a polynucleotide sequence that encodes a fusion protein that comprises an amino acid sequence tag.

[0021] The nucleic acid molecules of the invention may further comprise a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases. The position of such a nucleic acid sequence, relative to the other elements of the nucleic acid molecules of the invention, will be such that, a nucleic acid sequence of interest can be attached to the nucleic acid molecules of the invention, thereby producing a polynucleotide construct that encodes a fusion protein, the fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) the amino acid sequence tag, and on the other side by (iii) the amino acid sequence encoded by the amino acid sequence of interest.

[0022] In certain embodiments, the nucleic acid sequence that encodes an amino acid sequence tag may be, *e.g.*, a nucleic acid sequence that encodes an amino acid sequence that is capable of being post-translationally modified. For example, the nucleic acid sequence may be a nucleic acid sequence which encodes an amino acid sequence that is capable of being post-translationally modified by, *e.g.*, biotinylation, attachment of 4-phosphopanthetheine, attachment of lipoic acid, attachment of flavins, etc. In a preferred embodiment, the amino acid sequence is capable of being biotinylated. An exemplary nucleic acid sequence that encodes a protein or polypeptide having an amino acid sequence that is capable of being biotinylated is an amino acid sequence which encodes a portion of the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, *e.g.*, an amino acid sequence known as the Biotag™.

[0023] In certain other embodiments, the nucleic acid sequence that encodes an amino acid sequence tag may be, *e.g.*, a nucleic acid sequence which encodes an amino acid sequence that is capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent. Such amino acid sequences are known in the art and include, *e.g.*, a 6-Histidine tag, an epitope tag (*e.g.*, an amino acid sequence recognized by a specific antibody (or fragment thereof) such as, *e.g.*, the FLAG tag, the Myc tag, the HA tag, etc.) Thus, the nucleic acid molecules of the invention can, in some embodiments, be used to produce fusion proteins comprising: (i) an amino acid sequence which encodes an amino acid sequence that is capable of being recognized by a specific antibody (or fragment thereof) or other compound or reagent, and (ii) an amino acid sequence encoded by a nucleotide sequence of interest.

[0024] The invention also includes methods for producing polynucleotide constructs that encode fusion proteins that comprise one or more amino acid sequence tags. In certain embodiments, the invention generally includes methods of attaching a first nucleic acid molecule (*e.g.*, a nucleic acid molecule which has a nucleotide sequence which encodes a particular protein or polypeptide of interest) to a second nucleic acid molecule which comprises one or more nucleic acid sequence tags. The attachment of the first nucleic acid molecule to the second nucleic acid molecule may be accomplished by, *e.g.*, recombination (*e.g.*, recombinational cloning) and/or by topoisomerase-mediated cloning. The attachment of the first nucleic acid molecule to the second nucleic acid molecule will preferably result in a product polynucleotide construct which encodes a fusion protein, said fusion protein comprising: (i) the amino acid sequence tag; and (ii) the amino acid sequence encoded by the nucleotide sequence of the first nucleic acid molecule.

[0025] The invention also includes methods of producing fusion proteins that comprise one or more amino acid sequence tags. Also included are methods for producing fusion proteins that can be purified, concentrated or otherwise identified. The methods, according to this aspect of the invention, may comprise: (a) obtaining a host cell comprising a polynucleotide construct that encodes a fusion protein that comprises one or more amino acid sequence tags,

-10-

said polynucleotide construct produced according to a method of the invention; and (b) culturing said host cell under conditions wherein said fusion protein is produced by said host cell. The methods of the invention may further comprise culturing said host cell under conditions wherein said fusion protein is post-translationally modified in said host cell. In other embodiments of this aspect of the invention, the methods further comprise: (a) causing said fusion protein to be released from said host cell or treating said host cell such that said fusion protein is released from said host cell; and (b) contacting said fusion protein with a detecting composition comprising a molecule that is capable of interacting specifically with said fusion protein.

[0026] In certain exemplary embodiments, said fusion protein is a fusion protein that has been post-translationally modified, *e.g.*, a biotinylated fusion protein, and said detecting composition comprises avidin, streptavidin, or analogs and derivatives thereof.

[0027] The invention further comprises vectors comprising the nucleic acid molecules of the invention, host cells comprising the nucleic acid and/or vectors of the invention, and kits comprising the nucleic acid molecules, vectors, and/or host cells of the invention.

#### BRIEF DESCRIPTION OF THE DRAWINGS

[0028] Fig. 1 is a map which shows the general characteristics of pET104-DEST.

[0029] Figs. 2A-2C show the nucleotide sequence of pET104-DEST (SEQ ID NO:1).

[0030] Fig. 3 is a map which shows the general characteristics of pET104/GW/*lacZ*.

[0031] Fig. 4 is a map which shows the general characteristics of pET104/D-TOPO.

[0032] Figs. 5A-5B show the nucleotide sequence of pET104/D-TOPO (SEQ ID NO:2).

[0033] Fig. 6 is a map which shows the general characteristics of pET104/D/*lacZ*.

[0034] Fig. 7 is a map which shows the general characteristics of pcDNA6/Biotag<sup>TM</sup>-DEST.

[0035] Figs. 8A-8B show the nucleotide sequence of pcDNA6/Biotag<sup>TM</sup>-DEST (SEQ ID NO:3).

[0036] Fig. 9 is a map which shows the general characteristics of pcDNA6/Biotag<sup>TM</sup>-GW/*lacZ*.

[0037] Fig. 10 is a map which shows the general characteristics of pcDNA6/Biotag<sup>TM</sup>/D-TOPO.

[0038] Figs. 11A-11B show the nucleotide sequence of pcDNA6/Biotag<sup>TM</sup>/D-TOPO (SEQ ID NO:4).

[0039] Fig. 12 is a map which shows the general characteristics of pcDNA6/Biotag<sup>TM</sup>/*lacZ*.

[0040] Fig. 13 is a map which shows the general characteristics of pMT/Biotag<sup>TM</sup>-DEST.

[0041] Figs. 14A-14B show the nucleotide sequence of pMT/Biotag<sup>TM</sup>-DEST (SEQ ID NO:5).

[0042] Fig. 15 is a map which shows the general characteristics of pMT/Biotag<sup>TM</sup>/GW-*lacZ*.

[0043] Fig. 16 is a depiction of the recombination region of the expression clone resulting from pET104-DEST x entry clone, showing the nucleotide sequence of the recombination region (SEQ ID NO:25) and the amino acid sequence encoded therefrom (SEQ ID NO:26).

[0044] Fig. 17 is a schematic representation of the mechanism by which TOPO cloning is accomplished.

[0045] Fig. 18 is a flow-chart describing the general steps required for cloning and expressing a blunt-end PCR product using pET104/D-TOPO.

[0046] Fig. 19 is a depiction of a region of the pET104/D-TOPO vector surrounding the Biotag<sup>TM</sup>, showing the nucleotide sequence of the region (SEQ ID NO:27) and the amino acid sequence encoded therefrom (SEQ ID NO:28).

-12-

[0047] Fig. 20 is a depiction of the recombination region of the expression clone resulting from pcDNA6/Biotag™-DEST x entry clone, showing the nucleotide sequence of the recombination region (SEQ ID NO:29) and the amino acid sequence encoded therefrom (SEQ ID NO:30).

[0048] Fig. 21 is a flow-chart describing the general steps required for cloning and expressing a blunt-end PCR product using pcDNA6/Biotag™/D-TOPO.

[0049] Fig. 22 is a depiction of a region of the pcDNA6/Biotag™/D-TOPO vector surrounding the Biotag™, showing the nucleotide sequence of the region (SEQ ID NO:31) and the amino acid sequence encoded therefrom (SEQ ID NO:32).

[0050] Fig. 23 is a depiction of the recombination region of the expression clone resulting from pMT/Biotag™-DEST x entry clone, showing the nucleotide sequence of the recombination region (SEQ ID NO:33) and the amino acid sequence encoded therefrom (SEQ ID NO:34).

[0051] Fig. 24 is a map which shows the general characteristics of pCoHygro.

[0052] Fig. 25 is a map which shows the general characteristics of pCoBlast.

#### DETAILED DESCRIPTION OF THE INVENTION

[0053] The present invention relates generally to compositions and methods for producing nucleic acid molecules which encode fusion proteins, *e.g.*, fusion proteins that comprise one or more amino acid sequence tags. The invention also relates to methods for producing, purifying, concentrating and isolating fusion proteins using the compositions and methods described herein.

[0054] The invention relates to nucleic acid molecules comprising: (a) one or more recombination sites; and (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags.

[0055] The invention also relates to isolated nucleic acid molecules comprising: (a) one or more topoisomerase recognition sites and/or one or more topoisomerases; and (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags.

[0056] The invention also relates to isolated nucleic acid molecules comprising: (a) one or more recombination sites; (b) one or more topoisomerase recognition sites and/or one or more topoisomerases; and (c) one or more nucleic acid sequences which encode one or more amino acid sequence tags.

[0057] The nucleic acid molecules of the invention may be circular molecules, or they may be linear molecules.

[0058] As used herein, a nucleotide is a base-sugar-phosphate combination. Nucleotides are monomeric units of a nucleic acid molecule (DNA and RNA). The term nucleotide includes ribonucleoside triphosphates ATP, UTP, CTG, GTP and deoxyribonucleoside triphosphates such as dATP, dCTP, dITP, dUTP, dGTP, dTTP, or derivatives thereof. Such derivatives include, for example, [(S)dATP, 7-deaza-dGTP and 7-deaza-dATP. The term nucleotide as used herein also refers to dideoxyribonucleoside triphosphates (ddNTPs) and their derivatives. Illustrated examples of dideoxyribonucleoside triphosphates include, but are not limited to, ddATP, ddCTP, ddGTP, ddITP, and ddTTP. According to the present invention, a "nucleotide" may be unlabeled or detectably labeled by well known techniques. Detectable labels include, for example, radioactive isotopes, fluorescent labels, chemiluminescent labels, bioluminescent labels and enzyme labels.

[0059] As used herein, a nucleic acid molecule is a sequence of contiguous nucleotides (riboNTPs, dNTPs or ddNTPs, or combinations thereof) of any length which may encode a full-length polypeptide or a fragment of any length thereof, or which may be non-coding. As used herein, the terms "nucleic acid molecule" and "polynucleotide" and "polynucleotide construct" may be used interchangeably.

[0060] Polymerases for use in the invention include but are not limited to polymerases (DNA and RNA polymerases), and reverse transcriptases. DNA polymerases include, but are not limited to, *Thermus thermophilus* (Tth) DNA polymerase, *Thermus aquaticus* (Taq) DNA polymerase, *Thermotoga neopolitana* (Tne) DNA polymerase, *Thermotoga maritima* (Tma) DNA polymerase, *Thermococcus litoralis* (Tli or VENT™) DNA polymerase,

-14-

Pyrococcus furiosus (Pfu) DNA polymerase, DEEPVENT™ DNA polymerase, Pyrococcus woosii (Pwo) DNA polymerase, Pyrococcus sp KOD2 (KOD) DNA polymerase, Bacillus sterothermophilus (Bst) DNA polymerase, Bacillus caldophilus (Bca) DNA polymerase, Sulfolobus acidocaldarius (Sac) DNA polymerase, Thermoplasma acidophilum (Tac) DNA polymerase, Thermus flavus (Tfi/Tub) DNA polymerase, Thermus ruber (Tru) DNA polymerase, Thermus brockianus (DYNAZYME™) DNA polymerase, Methanobacterium thermoautotrophicum (Mth) DNA polymerase, mycobacterium DNA polymerase (Mtb, Mlep), E. coli pol I DNA polymerase, T5 DNA polymerase, T7 DNA polymerase, and generally pol I type DNA polymerases and mutants, variants and derivatives thereof. RNA polymerases such as T3, T5, T7 and SP6 and mutants, variants and derivatives thereof may also be used in accordance with the invention.

[0061] The nucleic acid polymerases used in the present invention may be mesophilic or thermophilic, and are preferably thermophilic. Preferred mesophilic DNA polymerases include Pol I family of DNA polymerases (and their respective Klenow fragments) any of which may be isolated from organism such as E. coli, H. influenzae, D. radiodurans, H. pylori, C. aurantiacus, R. prowazekii, T. pallidum, Synechocystis sp., B. subtilis, L. lactis, S. pneumoniae, M. tuberculosis, M. leprae, M. smegmatis, Bacteriophage L5, phi-C31, T7, T3, T5, SP01, SP02, mitochondrial from S. cerevisiae MIP-1, and eukaryotic C. elegans, and D. melanogaster (Astatke, M. et al., 1998, J. Mol. Biol. 278, 147-165), pol III type DNA polymerase isolated from any sources, and mutants, derivatives or variants thereof, and the like. Preferred thermostable DNA polymerases that may be used in the methods and compositions of the invention include Taq, Tne, Tma, Pfu, KOD, Tfi, Tth, Stoffel fragment, VENT™ and DEEPVENT™ DNA polymerases, and mutants, variants and derivatives thereof (U.S. Patent No. 5,436,149; U.S. Patent 4,889,818; U.S. Patent 4,965,188; U.S. Patent 5,079,352; U.S. Patent 5,614,365; U.S. Patent 5,374,553; U.S. Patent 5,270,179; U.S. Patent 5,047,342; U.S. Patent No. 5,512,462; WO 92/06188; WO 92/06200; WO 96/10640; WO 97/09451; Barnes, W.M., Gene 112:29-35 (1992); Lawyer,



F.C., et al., PCR Meth. Appl. 2:275-287 (1993); Flaman, J.-M, et al., Nucl. Acids Res. 22(15):3259-3260 (1994)).

[0062] Reverse transcriptases for use in this invention include any enzyme having reverse transcriptase activity. Such enzymes include, but are not limited to, retroviral reverse transcriptase, retrotransposon reverse transcriptase, hepatitis B reverse transcriptase, cauliflower mosaic virus reverse transcriptase, bacterial reverse transcriptase, Tth DNA polymerase, Taq DNA polymerase (Saiki, R.K., et al., Science 239:487-491 (1988); U.S. Patent Nos. 4,889,818 and 4,965,188), Tne DNA polymerase (WO 96/10640 and WO 97/09451), Tma DNA polymerase (U. S. Patent No. 5,374,553) and mutants, variants or derivatives thereof (see, e.g., WO 97/09451 and WO 98/47912). Preferred enzymes for use in the invention include those that have reduced, substantially reduced or eliminated RNase H activity. By an enzyme "substantially reduced in RNase H activity" is meant that the enzyme has less than about 20%, more preferably less than about 15%, 10% or 5%, and most preferably less than about 2%, of the RNase H activity of the corresponding wildtype or RNase H<sup>+</sup> enzyme such as wildtype Moloney Murine Leukemia Virus (M-MLV), Avian Myeloblastosis Virus (AMV) or Rous Sarcoma Virus (RSV) reverse transcriptases. The RNase H activity of any enzyme may be determined by a variety of assays, such as those described, for example, in U.S. Patent No. 5,244,797, in Kotewicz, M.L., et al., Nucl. Acids Res. 16:265 (1988) and in Gerard, G.F., et al., FOCUS 14(5):91 (1992), the disclosures of all of which are fully incorporated herein by reference. Particularly preferred polypeptides for use in the invention include, but are not limited to, M-MLV H<sup>-</sup> reverse transcriptase, RSV H<sup>-</sup> reverse transcriptase, AMV H<sup>-</sup> reverse transcriptase, RAV (rous-associated virus) H<sup>-</sup> reverse transcriptase, MAV (myeloblastosis-associated virus) H<sup>-</sup> reverse transcriptase and HIV H<sup>-</sup> reverse transcriptase. (See U.S. Patent No. 5,244,797 and WO 98/47912). It will be understood by one of ordinary skill, however, that any enzyme capable of producing a DNA molecule from a ribonucleic acid molecule (i.e., having reverse transcriptase activity) may be equivalently used in the compositions, methods and kits of the invention.

[0063] As used herein, a polypeptide is a sequence of contiguous amino acids, of any length. As used herein, the terms "peptide," "oligopeptide," or "protein" may be used interchangeably with the term "polypeptide."

[0064] As used herein, the term "amino acid sequence tag" is intended to mean any amino acid sequence that can be attached to, connected to, or linked to a heterologous amino acid sequence (*e.g.*, an amino acid sequence of interest) and that can be used to identify, purify, concentrate or isolate said heterologous amino acid sequence. The attachment of the amino acid sequence tag to the heterologous amino acid sequence may occur, *e.g.*, by constructing a nucleic acid molecule that comprises: (a) a nucleic acid sequence that encodes the amino acid sequence tag, and (b) a nucleic acid sequence that encodes a heterologous amino acid sequence. Exemplary amino acid sequence tags include, *e.g.*, amino acid sequences that are capable of being post-translationally modified. Other Exemplary amino acid sequence tags include, *e.g.*, amino acid sequences that are capable of being recognized and/or bound by an antibody (or fragment thereof) or other specific binding reagent.

[0065] As used herein, the expression "amino acid sequence that is capable of being post-translationally modified" is intended to mean any amino acid sequence, or portion thereof, that can be recognized, *in vivo* or *in vitro*, by an enzyme or other molecule that is capable of covalently attaching a chemical entity to one or more amino acids within the amino acid sequence.

[0066] As used herein, the term "post-translationally modified protein" is intended to mean at least one protein or polypeptide that has undergone or has been subjected to a post-translational modification. The term "post-translational modification" is intended to mean a modification that can take place *in vivo* (within a cell) or *in vitro* (outside a cell) whereby one or more chemical entities are covalently attached to at least one amino acid within the post-translational modification site by means of one or more enzymatic reactions. The site or sites include not only the amino acid that is modified, but any other amino acids, in the proper sequence, that are necessary to allow the post-translational modification to occur.

[0067] In the context of the present invention, the amino acid sequences that are capable of being post-translationally modified include amino acid sequences that are capable of being modified by any type of post-translational modification that provides a marker for a protein or polypeptide. The post-translational modifications that are included within the present invention include those that can be used, directly or indirectly, to identify a protein or polypeptide or to isolate it from a mixture of other materials, including other proteins, such as those found in a cell extract or in medium in which a host cell has been cultured and which contains the protein or polypeptide.

[0068] Amino acid sequences that are capable of being post-translationally modified include amino acid sequences that can be subjected to multiple (*e.g.*, 2, 3, 4, or 5 or more) post-translational modifications.

[0069] Preferred post-translational modifications are those that are utilized by a host cell to modify only a small number of proteins. Exemplary post-translational modifications that can be used with the present invention include biotinylation, attachment of 4-phosphopantetheine, attachment of lipoic acid and attachment of flavins and glycosylation. Further details regarding post-translational modifications of amino acid sequences can be found in U.S. Patent No. 5,252,466 and the references cited therein.

[0070] In a preferred embodiment of the invention, the amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being biotinylated (Parrott, M.B. and Barry, M.A., *Biochem. Biophys. Res. Comm.* 282:993-1000 (2001); Parrott, M.B. and Barry, M.A., *Mol. Ther.* 1:96-104 (2000)). Amino acid sequences that are capable of being biotinylated are known in the art. Exemplary amino acid sequences that are capable of being biotinylated include, *e.g.*, all or a portion of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, all or a portion of the *Propionibacterium shermanii* transcarboxylase 1.3S subunit, and all or a portion of the *Escherichia coli* biotin carboxyl carrier protein component of acetyl-CoA carboxylase.

[0071] According to certain embodiments of the invention, the amino acid sequence that is capable of being biotinylated is an amino acid sequence derived from the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit. In particular embodiments, the amino acid sequence that is capable of being biotinylated is a 72 amino acid peptide derived from the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit (Schwarz, E. *et al.*, *J. Biol. Chem.* 263:9640-9645 (1988)). This 72 amino acid sequence is also known as "the BIOTAG™." Biotin is covalently attached to the oxalacetate decarboxylase  $\alpha$  subunit and peptide sequencing has identified a single biotin binding site at lysine 561 of the protein. (Schwarz, E. *et al.*, *J. Biol. Chem.* 263:9640-9645 (1988)). When fused to a heterologous protein, the BIOTAG™ enables the *in vivo* biotinylation of the recombinant protein of interest. It is preferred that the entire 72 amino acid domain be used to ensure recognition by the cellular biotinylation enzymes. Additional details regarding cellular biotinylation enzymes and the mechanisms of biotinylation can be found in Chapman-Smith, A. and Cronan, J., *J. Nutr.* 129:477S-484S (1999).

[0072] Exemplary amino acid sequences that are capable of being biotinylated are listed in Table I. The nucleotide sequences encoding the exemplary amino acid sequence tags are listed in Table II.

TABLE I : Exemplary Amino Acid Sequences  
That are Capable of Being Biotinylated

Amino Acid Sequence Tag	Amino Acid Sequence
<i>K. pneumoniae</i> oxalacetate decarboxylase $\alpha$ subunit (Biotag™)	GAGTPVTAPLAGTIWKVLASEGQTVAAGE VLLILEAMKMETEIRAAQAGTVRGIKAG DAVAVGDTLMTLA (SEQ ID NO:6)
Mouse pyruvate decarboxylase domain	KALAVSDLNRAGQRQVFFELNGQLRSILVK DTQAMKEMHFHPKALKDVKGGQIGAPMPGK VIDIKVAAGDKVAKGQPLCVLSAMKMETV VTSPMEGTIRKVHVTKDMTLEGDDLIL (SEQ ID NO:7)
<i>P. shermanii</i> transcarboxylase domain	MKLKVTVNGTAYDVDVDVDKSHENPMGTI LFGGGTGGAPAPRAAGGAGAGKAGEGEIP APLAGTVSKILVKEGDTVKAGQTVLVLEA MKMETEINAPTDGKVEKVLVKERDAVQGG QGLIKIG (SEQ ID NO:8)
Human acetyl CoA Carboxylase domain	GSCVEVDVHRLSDGGLLSYDGSSYTTYM KEEVDRYRITIGNKTCVFEKENDPSVMRSPS AGKLIQYTVEDGGHVFAGQCYAEIEVMKM VMTLTAVESGCIHYVKRPGAALDPGCVLA KMQL (SEQ ID NO:9)
<i>E. coli</i> acetyl CoA carboxylase BCCP subunit	MDIRKIKKLIELVEESGISELEISEGEESVRIS RAAPAASFPVMQQAYAAPMMQQPAQSNA AAPATVPSMEAPAAAEISGHIVRSPMVGTF YRTPSPDAKAFIEVGQKVNVDGDTLCIVEAM KMMNQIEADKSGTVKAILVESGQPVEFDEP LVVIE (SEQ ID NO:10)

TABLE II : Nucleotide Sequences of Exemplary Amino Acid Sequence Tags

Amino Acid Sequence Tag	Nucleotide Sequence Encoding the Amino Acid Sequence Tag
<i>K. pneumoniae</i> oxalacetate decarboxylase $\alpha$ subunit (Biotag™)	ggcgccggcaccgggtgaccgccccgctggcgggcactatctgg aaggtgctggccagcgaagccagacgggtggccgagggcagggt gctgctgattctggaagccatgaagatggaaaccgaaatccgcgcc gcgcaggccgggaccgtgcgcggtatcgcggtgaaagccggcga cgcggtggcggtcggcgacaccctgatgacctggcg (SEQ ID NO:11)
Mouse pyruvate decarboxylase domain	aaagccctggctgtaagcgacctgaaccgtgctggccagaggcag gtgttcttgaactcaatgggcagcttcgatccattctggttaaagaca cccaggccatgaaggagatgcacttccatccaaggcttgaaggat gtgaagggccaaattggggccccgatgcctgggaaggtcatagac atcaaggtggcagcaggggacaaggtggctaaggccagccccctc tgtgtgctcagcgccatgaagatggagactgtggtgacttcgccat ggagggcactatccgaaagggtcatgttaccaggacatgactctgg aaggcgacgacctcatccta (SEQ ID NO:12)
<i>P. shermanii</i> transcarboxylase domain	atgaaactgaaggtaacagtcacgcgactgcgtatgacgttgacgt tgacgtcgacaagtcacacgaaaaccgatgggcaccatcctgttc ggcggcgccaccggcgccgcgcggcaccgcgcgcagcagggtg gcgcaggcgccggtgaaggccggagaggcgagattcccgtccg ctggccggcaccgtctcaagatcctcgtgaaggagggtgacacg gtcaaggctggtcagaccgtgctcgttcgaggccatgaagatgga gaccgagatcaacgctcccaccgacggcaaggctcgagaaggtcct tgtcaaggagcgtgacgccgtgcaggcggtcagggtctcatcaag atcggc (SEQ ID NO:13)
Human acetyl CoA Carboxylase domain	ggctcatgtgtagaagtagatgtacatcggtgagtacgggtggact gctcttgcctatgatggcagcagttacaccagtatatgaaggagga agtagacagatcgcatacaattggcaataaaacctgtgtgttga gaaggaaaatgaccatcggtgatgcgtcaccttctgctgggaagt taatccagtacattgtagaagatggaggtcatgtgttccggccagt gctatgcagagattgaggaatgaagatggtaatgactttgacagctg tggagtctggctgtatccattacgtcaagcgtcctggagcagctcttg accctggctgtgtactcgccaaaatgcaactg (SEQ ID NO:14)
<i>E. coli</i> acetyl CoA carboxylase BCCP subunit	atggatattcgtgaagattaaaaactgatcgagctggtgaagaatca ggcatctccgaactggaaattctgaaggcgaagagtcagtagcat tagccgtgcagctcctgccgaagttccctgtgatgcaacaagctta cgctgcaccaatgatgcagcagccagctcaatctaacgcagccgct ccggcgaccgttcttccatggaagcgccagcagcagcggaatc agtggtcacatcgtagtccccgatgggtggtactttctaccgcacc caagcccgacgcaaaagcgttcacgaagtgggtcagaaagtca acgtgggcgataccctgtgcatcgttgaagccatgaaatgatgaac cagatcgaagcgggacaaatccggtaccgtgaaagcaattctggtcg aaagtggacaaccggtagaatttgacgagccgctggtcgtcatcga g (SEQ ID NO:15)

[0073] An amino acid sequence tag, as used herein, may alternatively or additionally be an amino acid sequence that is capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent. The expression "amino acid sequence that is capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent" is intended to mean any amino acid sequence, or portion thereof, to which a particular compound or reagent can interact with or bind to, either covalently or non-covalently. Such amino acid sequences are known in the art. Preferred amino acid sequences that are capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent include, *e.g.*, those that are known in the art as "epitope tags." An epitope tag may be a natural or an artificial epitope tag. Natural and artificial epitope tags are known in the art, including, *e.g.*, artificial epitopes such as FLAG, Strep, or poly-histidine peptides. FLAG peptides include the sequence Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (SEQ ID NO:16) or Asp-Tyr-Lys-Asp-Glu-Asp-Asp-Lys (SEQ ID NO:17) (Einhauser, A. and Jungbauer, A., *J. Biochem. Biophys. Methods* 49:1-3:455-465 (2001)). The Strep epitope has the sequence Ala-Trp-Arg-His-Pro-Gln-Phe-Gly-Gly (SEQ ID NO:18). The VSV-G epitope can also be used and has the sequence Tyr-Thr-Asp-Ile-Glu-Met-Asn-Arg-Leu-Gly-Lys (SEQ ID NO:19). Another artificial epitope is a poly-His sequence having six histidine residues (His-His-His-His-His-His (SEQ ID NO:20). Naturally-occurring epitopes include the influenza virus hemagglutinin (HA) sequence Tyr-Pro-Tyr-Asp-Val-Pro-Asp-Tyr-Ala-Ile-Glu-Gly-Arg (SEQ ID NO:21) recognized by the monoclonal antibody 12CA5 (Murray *et al.*, *Anal. Biochem.* 229:170-179 (1995)) and the eleven amino acid sequence from human *c-myc* (Myc) recognized by the monoclonal antibody 9E10 (Glu-Gln-Lys-Leu-Leu-Ser-Glu-Glu-Asp-Leu-Asn (SEQ ID NO:22) (Manstein *et al.*, *Gene* 162:129-134 (1995)). Another useful epitope is the tripeptide Glu-Glu-Phe (SEQ ID NO:23) which is recognized by the monoclonal antibody YL 1/2. (Stammers *et al. FEBS Lett.* 283:298-302(1991)).

[0074] The nucleic acid molecules of the invention may include a variety of elements. The nucleic acid molecule of the invention preferably comprises one or more nucleic acid sequences which encode one or more amino acid sequence tags. The nucleic acid molecules may also comprise one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases.

[0075] The nucleic acid molecules of the invention may also comprise one or more selectable markers, one or more cloning sites, one or more restriction sites, one or more promoters, one or more operators (*e.g.*, a tet operator, a galactose operon operator, a lac operon operator, and the like), one or more operons, one or more origins of replication, one or more nucleotide sequences that encode a gene product which allows for negative selection, one or more nucleotide sequences which encode a repressor of at least one promoter, and one or more genes or gene products. Additional elements useful for molecular biology applications will be known to those skilled in the art and can be included within the nucleic acid molecules of the invention as well. The exact combination of elements, and their relative locations within the nucleic acid molecules of the invention, may vary depending on the intended uses of the nucleic acid molecules.

[0076] As used herein, a selectable marker is intended to include a nucleic acid segment that allows one to select for or against a molecule (*e.g.*, a replicon) or a cell that contains it, often under particular conditions. These markers can encode an activity, such as, but not limited to, production of RNA, peptide, or protein, or can provide a binding site for RNA, peptides, proteins, inorganic and organic compounds or compositions and the like. Examples of selectable markers include but are not limited to: (1) nucleic acid segments that encode products which provide resistance against otherwise toxic compounds (*e.g.*, antibiotics); (2) nucleic acid segments that encode products which are otherwise lacking in the recipient cell (*e.g.*, tRNA genes, auxotrophic markers); (3) nucleic acid segments that encode products which suppress the activity of a gene product; (4) nucleic acid segments that encode products which can be readily identified (*e.g.*, phenotypic markers such as



-23-

(-galactosidase, green fluorescent protein (GFP), enhanced green fluorescent protein (EGFP), and cell surface proteins); (5) nucleic acid segments that bind products which are otherwise detrimental to cell survival and/or function; (6) nucleic acid segments that otherwise inhibit the activity of any of the nucleic acid segments described in Nos. 1-5 above (e.g., antisense oligonucleotides); (7) nucleic acid segments that bind products that modify a substrate (e.g. restriction endonucleases); (8) nucleic acid segments that can be used to isolate or identify a desired molecule (e.g. specific protein binding sites); (9) nucleic acid segments that encode a specific nucleotide sequence which can be otherwise non-functional (e.g., for PCR amplification of subpopulations of molecules); (10) nucleic acid segments, which when absent, directly or indirectly confer resistance or sensitivity to particular compounds; and/or (11) nucleic acid segments that encode products which are toxic in recipient cells.

[0077] Exemplary selectable markers that can be included within the nucleic acid molecules of the invention include, *e.g.*, a gene encoding a product that confers resistance to chloramphenicol, *e.g.*, a chloramphenicol resistance gene (CmR), a gene encoding a product that confers resistance to ampicillin, *e.g.*, a gene which encodes  $\beta$ -lactamase, a gene encoding a product that confers resistance to other antibiotic compounds, a *ccdB* gene or other toxic genes (allowing for counterselection of the nucleic acid molecule), and a gene encoding a product that confers resistance to blasticidin, *e.g.*, a *bsd* resistance gene. Any other selectable marker gene known in the art can be include within the nucleic acid molecules of the invention.

[0078] A "cloning site," as used herein includes any nucleic acid regions which contain at least one restriction endonuclease cleavage sites. The nucleic acid molecules of the invention may also comprise "multiple cloning sites." A multiple cloning site is any nucleic acid region which contains two or more restriction endonuclease cleavage sites. "Restriction endonuclease cleavage sites are also referred to in the art as "restriction sites."

[0079] As used herein, a promoter is an example of a transcriptional regulatory sequence, and is specifically a nucleic acid sequence generally

described as the 5'-region of a gene located proximal to the start codon. The transcription of an adjacent nucleic acid segment is initiated at the promoter region. A repressible promoter's rate of transcription decreases in response to a repressing agent. An inducible promoter's rate of transcription increases in response to an inducing agent. A constitutive promoter's rate of transcription is not specifically regulated, though it can vary under the influence of general metabolic conditions.

[0080] Any promoter known to those skilled in the art can be included in the nucleic acid molecules of the invention. Exemplary promoters include, *e.g.*, the T7 promoter, the human cytomegalovirus (CMV) immediate early enhancer/promoter, the SV40 early promoter, a metallothionein (MT) promoter, including, *e.g.*, the *Drosophila* MT promoter. Other exemplary promoters include those that are inducible by, or can be repressed by, *e.g.*, certain carbon sources (*e.g.*, glucose, galactose, arabinose, etc.), salts, temperature changes (*e.g.*, temperatures greater than or less than the normal physiological growth temperature), and other molecules.

[0081] A number of operators are known in the art and can be included in the nucleic acid molecules of the invention. An example of an operator suitable for use with the invention is the tryptophan operator of the tryptophan operon of *E. coli*. The tryptophan repressor, when bound to two molecules of tryptophan, binds to the *E. coli* tryptophan operator and, when suitably positioned with respect to the promoter, blocks transcription. Another example of an operator suitable for use with the invention is operator of the *E. coli* tetracycline operon. Components of the tetracycline resistance system of *E. coli* have also been found to function in eukaryotic cells and have been used to regulate gene expression. For example, the tetracycline repressor, which binds to tetracycline operator in the absence of tetracycline and represses gene transcription, has been expressed in plant cells at sufficiently high concentrations to repress transcription from a promoter containing tetracycline operator sequences (Gatz *et al.*, *Plants* 2:397-404 (1992)). The tetracycline regulated expression systems are described, for example in U.S. Patent No. 5,789,156, the entire disclosure of which is incorporated herein by reference.

Additional examples of operators which can be used with the invention include the *Lac* operator and the operator of the molybdate transport operator/promoter system of *E. coli* (see, e.g., Cronin *et al.*, *Genes Dev.* 15:1461-1467 (2001) and Grunden *et al.*, *J. Biol. Chem.*, 274:24308-24315 (1999)).

[0082] Thus, in particular embodiments, the invention provides nucleic acid molecules that contain one or more operators which can be used to regulate expression in prokaryotic or eukaryotic cells. As one skilled in the art would recognize, when a nucleic acid molecule which contains an operator is placed under conditions in which transcriptional machinery is present, either *in vivo* or *in vitro*, regulation of expression will often be modulated by contacting the nucleic acid molecule with a repressor and one or more metabolites which facilitate binding of an appropriate repressor to the operator. Thus, the invention further provides nucleic acid molecules which encode repressors which modulate the function of operators.

[0083] The nucleic acid molecules of the invention may comprise one or more genes or partial genes. As used herein, a gene is a nucleic acid sequence that contains information necessary for expression of a polypeptide, protein or functional RNA (e.g., a ribozyme, tRNA, rRNA, mRNA, etc.). It includes the promoter and the structural gene open reading frame sequence (orf) as well as other sequences involved in expression of the protein. As used herein, a structural gene refers to a nucleic acid sequence that is transcribed into messenger RNA that is then translated into a sequence of amino acids characteristic of a specific polypeptide.

[0084] The range of positions of the various elements of the nucleic acid molecules of the invention, relative to one another, will be appreciated by persons having ordinary skill in the art. For example, a nucleic acid molecule within the scope of the invention may comprise (a) one or more recombination sites; and (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags. In a preferred embodiment, elements (a) and (b) will be positioned relative to one another such that a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more

-26-

recombination sites, thereby producing a polynucleotide construct that encodes a fusion protein. Such fusion protein may comprise: (i) the amino acid sequence tag, and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest.

[0085] Similarly, a nucleic acid molecule within the scope of the invention may comprise (a) one or more topoisomerase recognition sites and/or one or more topoisomerases; and (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags. In a preferred embodiment, elements (a) and (b) will be positioned relative to one another such that a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotides of the position of said one or more topoisomerases, thereby producing a polynucleotide construct that encodes a fusion protein. Such fusion protein may comprise: (i) the amino acid sequence tag, and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest.

[0086] Similarly, a nucleic acid molecule within the scope of the invention may comprise (a) one or more recombination sites; (b) one or more topoisomerase recognition sites and/or one or more topoisomerases; and (c) one or more nucleic acid sequences which encode one or more amino acid sequence tags. In a preferred embodiment, elements (a), (b) and (c) will be positioned relative to one another such that a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more recombination sites, thereby producing a polynucleotide construct that encodes a fusion protein. Such fusion protein may comprise: (i) the amino acid sequence tag, and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest. In another preferred embodiment, elements (a), (b) and (c) will be positioned relative to one another such that a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotides of the position of said one or more topoisomerases, thereby producing a polynucleotide construct that encodes a fusion protein. Such fusion protein may comprise: (i) the amino

-27-

acid sequence tag, and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest.

[0087] In certain embodiments, the nucleic acid molecules of the invention will comprise a nucleic acid sequence that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases. Amino acid sequences that can be recognized and/or cleaved by one or more proteases are known in the art. Exemplary amino acid sequences are those that are recognized by the following proteases: factor VIIa, factor IXa, factor Xa, APC, t-PA, u-PA, trypsin, chymotrypsin, enterokinase, pepsin, cathepsin B,H,L,S,D, cathepsin G, renin, angiotensin converting enzyme, matrix metalloproteases (collagenases, stromelysins, gelatinases), macrophage elastase, Cir, and Cis. The amino acid sequences that are recognized by the aforementioned proteases are known in the art. Exemplary sequences recognized by certain proteases can be found, *e.g.*, in U.S. Patent No. 5,811,252. A preferred amino acid sequence that is capable of being recognized and/or cleaved by a protease is the enterokinase (EK) recognition site (Asp-Asp-Asp-Asp-Lys (SEQ ID NO:24)).

[0088] The invention therefore also includes nucleic acid molecules comprising: (a) one or more recombination sites; (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags; and (c) one or more nucleic acid sequences that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases.

[0089] The invention also includes nucleic acid molecules comprising: (a) one or more topoisomerase recognition sites and/or one or more topoisomerases; (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags; and (c) one or more nucleic acid sequence that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases. In a preferred aspect, the nucleic acid sequence that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases is positioned such that, upon cleavage, the amino acid sequence tag is completely or partially removed from the amino acid sequence of interest. In another aspect, the nucleic acid sequence that

-28-

encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases is positioned such that, upon cleavage, other sequences (*e.g.*, topoisomerase recognition sequences and/or recombination sites) may be removed from the amino acid sequence of interest.

[0090] The invention also includes nucleic acid molecules comprising: (a) one or more recombination sites; (b) one or more topoisomerase recognition sites and/or one or more topoisomerases; (c) one or more nucleic acid sequences which encode one or more amino acid sequence tags; and (d) one or more nucleic acid sequence that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases. In a preferred aspect, the nucleic acid sequence that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases is positioned such that, upon cleavage, the amino acid sequence tag is completely or partially removed from the amino acid sequence of interest. In another aspect, the nucleic acid sequence that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases is positioned such that, upon cleavage, other sequences (*e.g.*, topoisomerase recognition sequences and/or recombination sites) may be removed from the amino acid sequence of interest.

[0091] The position of a nucleic acid sequence that encodes an amino acid sequence that is capable of being recognized and/or cleaved by one or more proteases, relative to the other elements of the nucleic acid molecules of the invention will be such that a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more recombination sites, or at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotides of the position of said one or more topoisomerases, thereby producing a polynucleotide construct that encodes a fusion protein. Such fusion protein may comprise: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid sequence tag, and on the other side by (iii) the amino acid sequence encoded by said nucleic acid sequence of interest.

[0092] This arrangement of elements will enable the production of a fusion protein of interest comprising an amino acid sequence tag, and will also enable the subsequent cleavage of the fusion protein by a protease, thereby separating the amino acid sequence tag from the amino acid sequence encoded by said nucleic acid sequence of interest. If the fusion protein is a fusion protein that is capable of being post-translationally modified, cleavage by the protease can be accomplished either before or after the post-translational modification of the fusion protein.

[0093] In addition to comprising one or more nucleic acid sequences which encode one or more amino acid sequence tags and/or one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases and/or one or more nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases, the nucleic acid molecules of the invention may further comprise additional elements. Exemplary additional elements that can be included within the nucleic acid molecules of the invention include, *e.g.*, one or more promoters, one or more selectable markers, one or more origins of replication, one or more operators, one or more enhancers, one or more ribosome binding sites, one or more initiation codons, one or more nucleic acid sequences of interest (*e.g.*, one or more nucleic acid sequences encoding one or more protein or polypeptides of interest), one or more polyadenylation signals, and/or one or more transcription termination regions. As understood by those skilled in the art, other elements may be included within the nucleic acid molecules of the invention depending on the circumstances under which the nucleic acids are intended to be used.

[0094] The possible arrangements of the various elements of the nucleic acid molecules of the invention, relative to one another, will be appreciated by persons having ordinary skill in the art. Non-limiting, exemplary arrangements are as follows:

[0095] Exemplary arrangement I: (a) one or more promoters – (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (c) one or more nucleic acid sequences that encodes an amino acid sequence

that is capable of being cleaved by one or more proteases – (d) one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (e) one or more polyadenylation signals and/or one or more transcription termination regions.

[0096] Exemplary arrangement II: (a) one or more promoters – (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (c) one or more nucleic acid sequences that encodes an amino acid sequence that is capable of being cleaved by one or more proteases – (d) one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (e) one or more nucleic acid sequences of interest – (f) one or more polyadenylation signals and/or one or more transcription termination regions.

[0097] Exemplary arrangement III: (a) one or more promoters – (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (c) one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (d) one or more polyadenylation signals and/or one or more transcription termination regions.

[0098] Exemplary arrangement IV: (a) one or more promoters – (b) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (c) one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (d) one or more nucleic acid sequences of interest – (e) one or more polyadenylation signals and/or one or more transcription termination regions.

[0099] Exemplary arrangement V: (a) one or more promoters – (b) one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (c) one or more nucleic acid sequences that encodes an amino acid sequence that is capable of being cleaved by one or more proteases – (d) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (e) one or more polyadenylation signals and/or one or more transcription termination regions.

[00100] Exemplary arrangement VI: (a) one or more promoters – (b) one or more nucleic acid sequences of interest – (c) one or more recombination sites



and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (d) one or more nucleic acid sequences that encodes an amino acid sequence that is capable of being cleaved by one or more proteases – (e) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (f) one or more polyadenylation signals and/or one or more transcription termination regions.

[00101] Exemplary arrangement VII: (a) one or more promoter – (b) one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (c) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (d) one or more polyadenylation signals and/or one or more transcription termination regions.

[00102] Exemplary arrangement VIII: (a) one or more promoters – (b) one or more nucleic acid sequences of interest – (c) one or more recombination sites and/or one or more topoisomerase recognition sites and/or one or more topoisomerases – (d) one or more nucleic acid sequences which encode one or more amino acid sequence tags – (e) one or more polyadenylation signals and/or one or more transcription termination regions.

[00103] In the foregoing exemplary arrangements, it will be understood by those skilled in the art that one or more additional elements may be included between any of the specifically listed elements, and/or that any of the specifically listed elements may be omitted. It will also be understood that many variations on these exemplary arrangements are possible (e.g., addition and/or omission of various elements) such that the nucleic acid molecules of the invention will allow the insertion of a nucleic acid sequence of interest and/or the production of a polynucleotide construct that encodes a desired fusion protein.

[00104] Persons of ordinary skill in the art will readily understand how close together, or how far apart, the elements of the nucleic acid molecules of the invention can be in order to permit the insertion of a nucleic acid sequence of interest and/or the production of a polynucleotide construct that encodes a desired fusion protein. For example, any two or more of the foregoing elements may be arranged within the nucleic acid molecules of the invention

such that they are within about 500 nucleotides of one another. In certain embodiments, any two or more elements of the nucleic acid molecules will be within about 400 nucleotides of one another, within about 300 nucleotides of one another, within about 200 nucleotides of one another, within about 100 nucleotides of one another, within about 50 nucleotides of one another, within about 40 nucleotides of one another, within about 30 nucleotides of one another, within about 20 nucleotides of one another, within about 10 nucleotides of one another, within about 5 nucleotides of one another, within about 4 nucleotides of one another, within about 3 nucleotides of one another, within about 2 nucleotides of one another, or within about 1 nucleotide of one another. The elements of the nucleic acid molecules of the invention may alternatively be directly adjacent to one another (*e.g.*, with no nucleotides separating them), as long as such an arrangement permits the insertion of a nucleic acid sequence of interest and/or the production of a polynucleotide construct that encodes a desired fusion protein.

[00105] It will also be appreciated that the nucleic acid sequence of interest will be preferably designed such that, when it is inserted at or within 20 nucleotides of said one or more recombination sites or at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotides of the position of said one or more topoisomerases, the nucleic acid sequence of interest is in frame with the nucleic acid sequence tag.

[00106] The nucleic acid molecules of the invention are useful, *e.g.*, in the production of fusion proteins that comprise one or more amino acid sequence tags. The fusion protein may be, *e.g.*, an N-terminal fusion protein (*e.g.*, wherein an amino acid sequence tag is covalently attached at or near the N-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest). The fusion protein may also be, *e.g.*, a C-terminal fusion protein (*e.g.*, wherein an amino acid sequence tag is covalently attached at or near the C-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest). The fusion protein may also be, *e.g.*, an N-terminal and C-terminal fusion protein (*e.g.*, wherein an amino acid sequence tag is covalently attached

at or near the N-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest and an amino acid sequence tag is covalently attached at or near the C-terminus of the amino acid sequence encoded by said nucleic acid sequence of interest).

[00107] The nucleic acid molecules of the invention may comprise one or more (e.g., 2, 3, 4, 5, 6, 7, 8, etc.) recombination sites. As used herein, a recombination site is a recognition sequence on a nucleic acid molecule participating in an integration/recombination reaction by recombination proteins. Recombination sites are discrete sections or segments of nucleic acid on the participating nucleic acid molecules that are recognized and bound by a site-specific recombination protein during the initial stages of integration or recombination. For example, the recombination site for Cre recombinase is loxP which is a 34 base pair sequence comprised of two 13 base pair inverted repeats (serving as the recombinase binding sites) flanking an 8 base pair core sequence. See Fig. 1 of Sauer, B., *Curr. Opin. Biotech.* 5:521-527 (1994). Other examples of recognition sequences include the attB, attP, attL, and attR sequences described herein, and mutants, fragments, variants and derivatives thereof, which are recognized by the recombination protein (Int and by the auxiliary proteins integration host factor (IHF), FIS and excisionase (Xis). See Landy, *Curr. Opin. Biotech.* 3:699-707 (1993).

[00108] Recombination sites for use in the invention may be any nucleic acid sequence that can serve as a substrate in a recombination reaction. Such recombination sites may be wild-type or naturally occurring recombination sites or modified or mutant recombination sites. Examples of recombination sites for use in the invention include, but are not limited to, phage-lambda recombination sites (such as attP, attB, attL, and attR and mutants or derivatives thereof) and recombination sites from other bacteriophage such as phi80, P22, P2, 186, P4 and P1 (including lox sites such as loxP and loxP511). Novel mutated att sites (e. g., attB 1-10, attP 1-10, attR 1-10 and attL 1-10) are described in International Patent Application PCT/US00/05432, which is specifically incorporated herein by reference. Other recombination sites having unique specificity (i.e., a first site will recombine with its

-34-

corresponding site and will not recombine with a second site having a different specificity) are known to those skilled in the art and may be used to practice the present invention.

[00109] Corresponding recombination proteins for these systems may be used in accordance with the invention with the indicated recombination sites. Other systems providing recombination sites and recombination proteins for use in the invention include the FLP/FRT system from *Saccharomyces cerevisiae*, the resolvase family (e.g.,  $\phi$ , Tn3 resolvase, Hin, Gin and Cin), and IS231 and other *Bacillus thuringiensis* transposable elements. Other suitable recombination systems for use in the present invention include the XerC and XerD recombinases and the psi, dif and cer recombination sites in *E. coli*. Other suitable recombination sites may be found in United States patent nos. 5,851,808 and 6,410,317 which are specifically incorporated herein by reference. Preferred recombination proteins and mutant or modified recombination sites for use in the invention include those described in U.S. Patent Nos. 5,888,732, 6,171,861, 6,143,557, 6,270,969 and 6,277,608, and commonly owned, co-pending U.S. Application Nos. 09/438,358 (filed 11/12/99), 09/517,466 (filed 03/02/00), 09/695,065 (filed 10/25/00), 09/732,914 (filed 12/11/00), and international application Nos. WO 01/11058 and WO 01/42509, the disclosures of all of which are incorporated herein by reference in their entireties, as well as those associated with the GATEWAY™ Cloning Technology and Echo™ Cloning Technology available from Invitrogen Corporation (Carlsbad, CA).

[00110] The nucleic acid molecules of the invention may comprise one or more (e.g., 2, 3, 4, 5, 6, 7, 8, etc.) topoisomerase recognition sites and/or one or more topoisomerases. As used herein, a topoisomerase recognition sequence (alternatively and equivalently referred to herein as a "topoisomerase recognition site") is a particular sequence to which a topoisomerase recognizes and binds. Examples of topoisomerase recognition sites include, but are not limited to, the sequence 5'-GCAACTT-3' that is recognized by *E. coli* topoisomerase III (a type I topoisomerase); the sequence 5'-(C/T)CCTT-3'

which is a topoisomerase recognition site that is bound specifically by most poxvirus topoisomerases, including vaccinia virus DNA topoisomerase I; and others that are known in the art as discussed elsewhere herein.

[00111] Topoisomerases are categorized as type I, including type IA and type IB topoisomerases, which cleave a single strand of a double stranded nucleic acid molecule, and type II topoisomerases (gyrases), which cleave both strands of a nucleic acid molecule. Type IA and IB topoisomerases cleave one strand of a nucleic acid molecule. Cleavage of a nucleic acid molecule by type IA topoisomerases generates a 5' phosphate and a 3' hydroxyl at the cleavage site, with the type IA topoisomerase covalently binding to the 5' terminus of a cleaved strand. In comparison, cleavage of a nucleic acid molecule by type IB topoisomerases generates a 3' phosphate and a 5' hydroxyl at the cleavage site, with the type IB topoisomerase covalently binding to the 3' terminus of a cleaved strand. As disclosed herein, type I and type II topoisomerases, as well as catalytic domains and mutant forms thereof, are useful for generating ds recombinant nucleic acid molecules covalently linked in both strands according to a method of the invention.

[00112] Type IA topoisomerases include *E. coli* topoisomerase I, *E. coli* topoisomerase III, eukaryotic topoisomerase II, archeal reverse gyrase, yeast topoisomerase III, *Drosophila* topoisomerase III, human topoisomerase III, *Streptococcus pneumoniae* topoisomerase III, and the like, including other type IA topoisomerases (see Berger, *Biochim. Biophys. Acta* 1400:3-18, 1998; DiGate and Marians, *J. Biol. Chem.* 264:17924-17930, 1989; Kim and Wang, *J. Biol. Chem.* 267:17178-17185, 1992; Wilson et al., *J. Biol. Chem.* 275:1533-1540, 2000; Hanai et al., *Proc. Natl. Acad. Sci., USA* 93:3653-3657, 1996, U.S. Pat. No. 6,277,620, each of which is incorporated herein by reference). *E. coli* topoisomerase III, which is a type IA topoisomerase that recognizes, binds to and cleaves the sequence 5'-GCAACTT-3', can be particularly useful in a method of the invention (Zhang et al., *J. Biol. Chem.* 270:23700-23705, 1995, which is incorporated herein by reference). A homolog, the traE protein of plasmid RP4, has been described by Li et al., *J. Biol. Chem.* 272:19582-19587 (1997) and can also be used in the practice of

-36-

the invention. A DNA-protein adduct is formed with the enzyme covalently binding to the 5'-thymidine residue, with cleavage occurring between the two thymidine residues.

[00113] Type IB topoisomerases include the nuclear type I topoisomerases present in all eukaryotic cells and those encoded by vaccinia and other cellular poxviruses (see Cheng et al., *Cell* 92:841-850, 1998, which is incorporated herein by reference). The eukaryotic type IB topoisomerases are exemplified by those expressed in yeast, *Drosophila* and mammalian cells, including human cells (see Caron and Wang, *Adv. Pharmacol.* 29B:271-297, 1994; Gupta et al., *Biochim. Biophys. Acta* 1262:1-14, 1995, each of which is incorporated herein by reference; see, also, Berger, *supra*, 1998). Viral type IB topoisomerases are exemplified by those produced by the vertebrate poxviruses (vaccinia, Shope fibroma virus, ORF virus, fowlpox virus, and molluscum contagiosum virus), and the insect poxvirus (*Amsacta moorei* entomopoxvirus) (see Shuman, *Biochim. Biophys. Acta* 1400:321-337, 1998; Petersen et al., *Virology* 230:197-206, 1997; Shuman and Prescott, *Proc. Natl. Acad. Sci., USA* 84:7478-7482, 1987; Shuman, *J. Biol. Chem.* 269:32678-32684, 1994; U.S. Pat. No. 5,766,891; PCT/US95/16099; PCT/US98/12372,, each of which is incorporated herein by reference; see, also, Cheng et al., *supra*, 1998).

[00114] Type II topoisomerases include, for example, bacterial gyrase, bacterial DNA topoisomerase IV, eukaryotic DNA topoisomerase II, and T-even phage encoded DNA topoisomerases (Roca and Wang, *Cell* 71:833-840, 1992; Wang, *J. Biol. Chem.* 266:6659-6662, 1991, each of which is incorporated herein by reference; Berger, *supra*, 1998). Like the type IB topoisomerases, the type II topoisomerases have both cleaving and ligating activities. In addition, like type IB topoisomerase, substrate nucleic acid molecules can be prepared such that the type II topoisomerase can form a covalent linkage to one strand at a cleavage site. For example, calf thymus type II topoisomerase can cleave a substrate nucleic acid molecule containing a 5' recessed topoisomerase recognition site positioned three nucleotides from the 5' end, resulting in dissociation of the three nucleotide sequence 5' to the cleavage site and

covalent binding the of the topoisomerase to the 5' terminus of the nucleic acid molecule (Andersen et al., *supra*, 1991). Furthermore, upon contacting such a type II topoisomerase charged nucleic acid molecule with a second nucleotide sequence containing a 3' hydroxyl group, the type II topoisomerase can ligate the sequences together, and then is released from the recombinant nucleic acid molecule. As such, type II topoisomerases also are useful in the nucleic acid molecules and methods of the invention.

[00115] Structural analysis of topoisomerases indicates that the members of each particular topoisomerase families, including type IA, type IB and type II topoisomerases, share common structural features with other members of the family (Berger, *supra*, 1998). In addition, sequence analysis of various type IB topoisomerases indicates that the structures are highly conserved, particularly in the catalytic domain (Shuman, *supra*, 1998; Cheng et al., *supra*, 1998; Petersen et al., *supra*, 1997). For example, a domain comprising amino acids 81 to 314 of the 314 amino acid vaccinia topoisomerase shares substantial homology with other type IB topoisomerases, and the isolated domain has essentially the same activity as the full length topoisomerase, although the isolated domain has a slower turnover rate and lower binding affinity to the recognition site (see Shuman, *supra*, 1998; Cheng et al., *supra*, 1998). In addition, a mutant vaccinia topoisomerase, which is mutated in the amino terminal domain (at amino acid residues 70 and 72) displays identical properties as the full length topoisomerase (Cheng et al., *supra*, 1998). In fact, mutation analysis of vaccinia type IB topoisomerase reveals a large number of amino acid residues that can be mutated without affecting the activity of the topoisomerase, and has identified several amino acids that are required for activity (Shuman, *supra*, 1998). In view of the high homology shared among the vaccinia topoisomerase catalytic domain and the other type IB topoisomerases, and the detailed mutation analysis of vaccinia topoisomerase, it will be recognized that isolated catalytic domains of the type IB topoisomerases and type IB topoisomerases having various amino acid mutations can be included with the nucleic acid molecules and methods of the invention.

[00116] The various topoisomerases exhibit a range of sequence specificity. For example, type II topoisomerases can bind to a variety of sequences, but cleave at a highly specific recognition site (*see Andersen et al., J. Biol. Chem.*, 266:9203-9210, 1991, which is incorporated herein by reference.). In comparison, the type IB topoisomerases include site specific topoisomerases, which bind to and cleave a specific nucleotide sequence ("topoisomerase recognition site"). Upon cleavage of a nucleic acid molecule by a topoisomerase, for example, a type IB topoisomerase, the energy of the phosphodiester bond is conserved via the formation of a phosphotyrosyl linkage between a specific tyrosine residue in the topoisomerase and the 3' nucleotide of the topoisomerase recognition site. Where the topoisomerase cleavage site is near the 3' terminus of the nucleic acid molecule, the downstream sequence (3' to the cleavage site) can dissociate, leaving a nucleic acid molecule having the topoisomerase covalently bound to the newly generated 3' end.

[00117] The nucleic acid molecules of the invention are useful, *e.g.*, for the production of fusion proteins. As used herein, the term "fusion protein" is intended to include any polypeptide which contains amino acids derived from at least two different polypeptides. The nucleic acid molecules of the invention are especially useful, *e.g.*, for producing fusion proteins comprising (i) one or more amino acid sequence tags, and (ii) one or more amino acid sequence encoded by one or more nucleic acid sequences of interest.

[00118] The invention also includes vectors comprising any of the nucleic acid molecules described herein. As used herein, a vector is a nucleic acid molecule (preferably DNA) that provides a useful biological or biochemical property to an insert. Examples include plasmids, phages, autonomously replicating sequences (ARS), centromeres, and other sequences which are able to replicate or be replicated in vitro or in a host cell, or to convey a desired nucleic acid segment to a desired location within a host cell. A Vector can have one or more restriction endonuclease recognition sites at which the sequences can be cut in a determinable fashion without loss of an essential biological function of the vector, and into which a nucleic acid fragment can



be spliced in order to bring about its replication and cloning. Vectors can further provide primer sites, e.g., for PCR, transcriptional and/or translational initiation and/or regulation sites, recombinational signals, replicons, selectable markers, etc. Clearly, methods of inserting a desired nucleic acid fragment which do not require the use of recombination, transpositions or restriction enzymes (such as, but not limited to, UDG cloning of PCR fragments (U.S. Patent No. 5,334,575, entirely incorporated herein by reference), TA Cloning® brand PCR cloning (Invitrogen Corporation, Carlsbad, CA) (also known as direct ligation cloning), and the like) can also be applied to clone a fragment into a cloning vector to be used according to the present invention. The cloning vector can further contain one or more selectable markers suitable for use in the identification of cells transformed with the cloning vector.

[00119] Exemplary vectors that are encompassed by the present invention include, e.g., pET104-DEST (SEQ ID NO:1) (Fig. 1), pET104/GW/*lacZ* (Fig. 2), pET104/D-TOPO (SEQ ID NO:2) (Fig. 3), pET104/D/*lacZ* (Fig. 4), pcDNA6/Biotag™-DEST (SEQ ID NO:3) (Fig. 5), pcDNA6/Biotag™-GW/*lacZ* (Fig. 6), pcDNA6/Biotag™/D-TOPO (SEQ ID NO:4) (Fig. 7), pcDNA6/Biotag™/*lacZ* (Fig. 8), pMT/Biotag™-DEST (SEQ ID NO:5) (Fig. 9), and pMT/Biotag™/GW-*lacZ* (Fig. 10).

[00120] The invention also encompasses nucleic acid molecules having nucleic acid sequences that are at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to at least 25, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000, 3000 or 4000 contiguous nucleotides of the exemplary vectors pET104-DEST (SEQ ID NO:1), pET104/D-TOPO (SEQ ID NO:2), pcDNA6/Biotag™-DEST (SEQ ID NO:3), pcDNA6/Biotag™/D-TOPO (SEQ ID NO:4) and pMT/Biotag™-DEST (SEQ ID NO:5). The invention also encompasses nucleic acid molecules comprising one or more nucleic acid sequences which encode an amino acid sequence tag, wherein said one or more nucleic acid sequences are at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to at least 25, 50, 75,

100, 125, 150, 175 or 200 contiguous nucleotides of any one of SEQ ID Nos:11-15.

[00121] By a nucleic acid molecule having a nucleotide sequence at least, for example, 80% "identical" to a reference nucleotide sequence it is intended that the nucleotide sequence of the nucleic acid molecule is identical to the reference sequence except that the nucleotide sequence may include up to 20 nucleotide alterations per each 100 nucleotides of the nucleotide sequence of the reference nucleic acid molecule. In other words, to obtain a nucleic acid molecule having a nucleotide sequence at least 80% identical to a reference nucleotide sequence, up to 20% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides, up to 20% of the total nucleotides in the reference sequence, may be inserted into the reference sequence. These alterations of the reference sequence may occur, *e.g.*, at the 5' or 3' ends of the reference nucleotide sequence and/or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence and/or in one or more contiguous groups within the reference sequence.

[00122] As a practical matter, whether any particular nucleic acid molecule is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to, for instance, a specified number of contiguous nucleotides of the nucleotide sequences shown in SEQ ID NOs:1-5 and 11-15 can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence

and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

[00123] A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag *et al.*, *Comp. Appl. Biosci.* 6:237-245 (1990). In a sequence alignment, the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

[00124] If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by the results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence are calculated for the purposes of manually adjusting the percent identity score.

-42-

[00125] For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and, therefore, the FASTDB alignment does not show a match/alignment of the first 10 bases at the 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence), so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal, so that there are no bases on the 5' or 3' ends of the subject sequence which are not matched/aligned with the query. In this case, the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

[00126] The invention also includes host cells comprising any of the nucleic acid molecules and/or vectors described herein. As used herein, a host cell is any prokaryotic or eukaryotic organism that is a recipient of a replicable expression vector, cloning vector or any nucleic acid molecule. As used herein, the terms "host," "host cell," "recombinant host" and "recombinant host cell" may be used interchangeably. Representative host cells that may be used with the invention include, but are not limited to, bacterial cells, yeast cells, plant cells and animal cells. Preferred bacterial host cells include *Escherichia* spp. cells (particularly *E. coli* cells and most particularly *E. coli* strains DH10B, Stbl2, DH5, DB3, DB3.1 (preferably *E. coli* LIBRARY EFFICIENCY® DB3.1™ Competent Cells; Invitrogen Corporation, Carlsbad, CA), DB4 and DB5 (see U.S. Application No. 09/518,188, filed March 2, 2000, the disclosure of which is incorporated by reference herein in its entirety), *Bacillus* spp. cells (particularly *B. subtilis* and *B. megaterium* cells), *Streptomyces* spp. cells, *Erwinia* spp. cells, *Klebsiella* spp. cells, *Serratia* spp. cells (particularly *S. marcescens* cells), *Pseudomonas* spp. cells (particularly *P.*

aeruginosa cells), and Salmonella spp. cells (particularly *S. typhimurium* and *S. typhi* cells). Preferred animal host cells include insect cells (most particularly *Drosophila melanogaster* cells, *Spodoptera frugiperda* Sf9 and Sf21 cells and *Trichoplusia* High-Five cells), nematode cells (particularly *C. elegans* cells), avian cells, amphibian cells (particularly *Xenopus laevis* cells), reptilian cells, and mammalian cells (most particularly NIH3T3, CHO, COS, VERO, BHK and human cells). Preferred yeast host cells include *Saccharomyces cerevisiae* cells and *Pichia pastoris* cells. These and other suitable host cells are available commercially, for example from Invitrogen Corporation (Carlsbad, California), American Type Culture Collection (Manassas, Virginia), and Agricultural Research Culture Collection (NRRL; Peoria, Illinois).

[00127] The nucleic acid molecules and/or vectors of the invention may be introduced into host cells using well known techniques of infection, transduction, electroporation, transfection, and transformation. The nucleic acid molecules and/or vectors of the invention may be introduced alone or in conjunction with other the nucleic acid molecules and/or vectors and/or proteins, peptides or RNAs. Alternatively, the nucleic acid molecules and/or vectors of the invention may be introduced into host cells as a precipitate, such as a calcium phosphate precipitate, or in a complex with a lipid. Electroporation also may be used to introduce the nucleic acid molecules and/or vectors of the invention into a host. Likewise, such molecules may be introduced into chemically competent cells such as *E. coli*. If the vector is a virus, it may be packaged in vitro or introduced into a packaging cell and the packaged virus may be transduced into cells. Hence, a wide variety of techniques suitable for introducing the nucleic acid molecules and/or vectors of the invention into host cells are well known and routine to those of skill in the art. Such techniques are reviewed at length, for example, in Sambrook, J., et al., *Molecular Cloning, a Laboratory Manual*, 2nd Ed., Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press, pp. 16.30-16.55 (1989), Watson, J.D., et al., *Recombinant DNA*, 2nd Ed., New York: W.H. Freeman and Co., pp. 213-234 (1992), and Winnacker, E.-L., *From Genes to Clones*, New York:

VCH Publishers (1987), which are illustrative of the many laboratory manuals that detail these techniques and which are incorporated by reference herein in their entireties for their relevant disclosures.

[00128] The present invention also includes methods of producing a polynucleotide construct that encodes a fusion protein that comprises one or more amino acid sequence tags. Such methods may be accomplished *in vivo* (e.g., within a cell) or *in vitro* (outside a cell).

[00129] According to one embodiment, the invention includes a method of producing a polynucleotide construct that encodes a fusion protein that comprises one or more amino acid sequence tags, said method comprising: (a) obtaining a first nucleic acid molecule comprising (i) a nucleotide sequence of interest and (ii) at least a first recombination site; (b) obtaining a second nucleic acid molecule comprising (i) one or more nucleic acid sequences which encode one or more amino acid sequence tags, and (ii) at least a second recombination site; and (c) combining said first nucleic acid molecule with said second nucleic acid molecule under conditions sufficient to cause recombination of at least said first and second recombination sites thereby producing a polynucleotide construct that encodes a fusion protein that comprises one or more amino acid sequence tags.

[00130] In certain embodiments, the methods of the invention comprise: (a) obtaining a first nucleic acid molecule comprising a nucleotide sequence of interest flanked by at least a first and at least a second recombination sites that do not recombine with each other; (b) obtaining a second nucleic acid molecule comprising: (i) at least a third and fourth recombination sites that do not recombine with each other; and (ii) one or more nucleic acid sequences which encode one or more amino acid sequence tags; and (c) contacting said first nucleic acid molecule with said second nucleic acid molecule under conditions favoring recombination between said first and third and between said second and fourth recombination sites, thereby producing a product polynucleotide construct; wherein said product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleotide acid sequence of interest.

[00131] In other embodiments, the methods of the invention comprise: (a) obtaining a first nucleic acid molecule comprising a nucleotide sequence of interest; (b) obtaining a second nucleic acid molecule comprising at least two topoisomerase recognition sites, at least one topoisomerase, and at least one nucleic acid sequence which encodes one or more amino acid sequence tags; (c) mixing said first nucleic acid molecule with said second nucleic acid molecule; and (d) incubating said mixture under conditions such that said first nucleic acid molecule is inserted into said second nucleic acid molecule between said at least two topoisomerase recognition sites, thereby producing a product polynucleotide construct; wherein said product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleotide sequence of interest.

[00132] In other embodiments, the methods of the invention comprise: (a) obtaining a first nucleic acid molecule comprising a nucleotide sequence of interest; (b) obtaining a second nucleic acid molecule comprising (i) at least a first topoisomerase recognition site, flanked by (ii) at least a first recombination site, and (iii) at least a second topoisomerase recognition site flanked by (iv) at least a second recombination site, wherein said first and second recombination sites do not recombine with each other, and (v) at least one topoisomerase; (c) obtaining a third nucleic acid molecule comprising: (i) at least a third and fourth recombination sites that do not recombine with each other; and (ii) one or more nucleic acid sequences which encode one or more amino acid sequence tags; (d) mixing said first nucleic acid molecule with said second nucleic acid molecule; (e) incubating said mixture under conditions such that said first nucleic acid molecule is inserted into said second nucleic acid molecule between said at least two topoisomerase recognition sites, thereby producing a first product polynucleotide construct; (f) contacting said first product polynucleotide construct with said third nucleic acid molecule under conditions favoring recombination between said first and third and between said second and fourth recombination sites, thereby producing a second product polynucleotide construct; wherein said second product

polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleotide sequence of interest.

[00133] In particular embodiments of the invention, one or more of the nucleic acid molecules that are used in the practice of the methods will further comprise a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases, and wherein the product polynucleotide constructs encode a fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) an amino acid sequence tag, and on the other side by (iii) the amino acid sequence encoded by a nucleotide sequence of interest. Any of the amino acid sequences that are capable of being cleaved by one or more proteases, as described elsewhere herein, can be used with the methods of the invention. In a preferred embodiment, the amino acid sequence that is capable of being cleaved by one or more proteases is an amino acid sequence that is capable of being cleaved by enterokinase.

[00134] The methods of the invention involve the use of nucleic acid molecules comprising one or more nucleic acid sequences which encode one or more amino acid sequence tags. Any of the nucleic acid sequences, described elsewhere herein, which encode an amino acid sequence tag, can be used in the context of the methods of the invention. In certain embodiments of the invention, the amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified. For example, the amino acid sequence tag may be an amino acid sequence that is capable of being biotinylated.

[00135] Any of the nucleic acid molecules, vectors, and host cells described herein, including any variations or modifications of such nucleic acid molecules vectors, and host cells, can be included in the practice of the methods of the invention. The nucleic acid molecules that are used in the practice of the methods of the invention may be linear, or circular. If a linear nucleic acid molecule is used, the ends of the molecule may be blunt ended or, alternatively, may have one or more overhang ends. The nucleic acid



molecules that are used in the practice of the methods of the invention may be PCR products.

[00136] The methods of the invention may further comprise inserting a product polynucleotide construct into a host cell.

[00137] In certain embodiments, the methods of the invention comprise contacting a first nucleic acid molecule comprising a first and a second recombination site with a second nucleic acid molecule comprising a third and a fourth recombination site under conditions favoring recombination between a first and third and between a second and fourth recombination sites.

[00138] Exemplary recombination sites included within the nucleic acid molecules that are used in the practice of the methods of the invention include, but are not limited to, (a) *attB* sites, (b) *attP* sites, (c) *attL* sites, (d) *attR* sites, (e) *lox* sites, (f) *psi* sites, (g) *dif* sites, (h) *cer* sites, (i) *frt* sites, and mutants, variants, and derivatives of the recombination sites of (a), (b), (c), (d), (e), (f), (g), (h), or (i) which retain the ability to undergo recombination.

[00139] In particular embodiments, said first and said second nucleic acid molecules are combined in the presence of at least one recombination protein. Exemplary recombination proteins that can be used in the methods of the invention include, *e.g.*, Cre, Int, IHF, Xis, Fis, Hin, Gin, Cin, Tn3 resolvase, TndX, XerC and XerD.

[00140] Methods for combining nucleic acid molecules by recombination at particular sites are known in the art. Such methods include, *e.g.*, recombinational cloning methods.

[00141] Cloning systems that utilize recombination at defined recombination sites have been previously described in U.S. Patent Nos. 5,888,732, 6,143,557, 6,171,861, 6,270,969, and 6,277,608, and in commonly owned, co-pending U.S. Application No. 10/005,876 (filed 12/07/01), which are specifically incorporated herein by reference. In brief, the Gateway™ Cloning System, described in this application and the applications referred to in the related applications section, utilizes vectors that contain at least one and preferably at least two different site-specific recombination sites based on the bacteriophage lambda system (*e. g.*, *att1* and *att2*) that are mutated from the wild type (*att0*)

-48-

sites. Each mutated site has a unique specificity for its cognate partner *att* site of the same type (for example *attB*1 with *attP*1, or *attL*1 with *attR*1) and will not cross-react with recombination sites of the other mutant type or with the wild-type *att0* site. Nucleic acid fragments flanked by recombination sites are cloned and subcloned using the Gateway™ system by replacing a selectable marker (for example, *ccdB*) flanked by *att* sites on the recipient plasmid molecule, sometimes termed the Destination Vector. Desired clones are then selected by transformation of a *ccdB* sensitive host strain and positive selection for a marker on the recipient molecule. Similar strategies for negative selection (e.g., use of toxic genes) can be used in other organisms such as thymidine kinase (TK) in mammals and insects.

[00142] Mutating specific residues in the core region of the *att* site can generate a large number of different *att* sites. As with the *att1* and *att2* sites utilized in Gateway™, each additional mutation potentially creates a novel *att* site with unique specificity that will recombine only with its cognate partner *att* site bearing the same mutation and will not cross-react with any other mutant or wild-type *att* site. Novel mutated *att* sites (e. g., *attB* 1-10, *attP* 1-10, *attR* 1-10 and *attL* 1-10) are described in International Patent Application PCT/US00/05432, which is specifically incorporated herein by reference. Other recombination sites having unique specificity (i.e., a first site will recombine with its corresponding site and will not recombine or not substantially recombine with a second site having a different specificity) may be used to practice the present invention. Examples of suitable recombination sites include, but are not limited to, *loxP* sites and derivatives such as *loxP*511 (see U.S. Patent No. 5,851,808), *frt* sites and derivatives, *dif* sites and derivatives, *psi* sites and derivatives and *cer* sites and derivatives. The present invention provides novel methods using such recombination sites to join or link multiple nucleic acid molecules or segments and more specifically to clone such multiple segments into one or more vectors containing one or more recombination sites (such as any Gateway™ Vector including Destination Vectors).

[00143] In certain embodiments, the methods of the invention comprise (a) mixing a first nucleic acid molecule with a second nucleic acid molecule, said second nucleic acid molecule comprising at least two topoisomerase recognition sites and at least one topoisomerase, and (b) incubating the mixture under conditions such that said first nucleic acid molecule is inserted into said second nucleic acid molecule between said at least two topoisomerase recognition sites.

[00144] Methods for inserting a first nucleic acid molecule into a second nucleic acid molecule between topoisomerase recognition sites thereby producing a product polynucleotide construct, are known in the art. Exemplary methods are known in the art as Topoisomerase cloning, TOPO® cloning, and Directional TOPO® cloning. As used herein, the term "topoisomerase-mediated cloning" is intended to mean any method of combining two or more nucleic acid molecules using at least one topoisomerase recognition site on one or more of the nucleic acid molecules and one or more topoisomerase. Exemplary methods are described in commonly owned, co-pending U.S. Application No. 10/005,876 (filed 12/07/01), the disclosure of which is incorporated herein by reference in its entirety.

[00145] A method for generating a product polynucleotide construct using topoisomerase cloning can be performed, for example, by contacting a first nucleic acid molecule having a first end and a second end, wherein, at the first end or second end or both, the first nucleic acid molecule has a topoisomerase recognition site (or cleavage product thereof) at or near the 3' terminus; at least a second nucleic acid molecule having a first end and a second end, wherein, at the first end or second end or both, the at least second double stranded nucleotide sequence has a topoisomerase recognition site (or cleavage product thereof) at or near a 3' terminus; and at least one site specific topoisomerase (e.g., a type IA and/or a type IB topoisomerase), under conditions such that all components are in contact and the topoisomerase can effect its activity.

[00146] In one embodiment, the method is performed by contacting a first nucleic acid molecule and a second (or other) nucleic acid molecule, each of which has a topoisomerase recognition site, or a cleavage product thereof, at the 3' termini or at the 5' termini of two ends to be covalently linked. In another embodiment, the method is performed by contacting a first nucleic acid molecule having a topoisomerase recognition site, or cleavage product thereof, at the 5' terminus and the 3' terminus of at least one end, and a second (or other) nucleic acid molecule having a 3' hydroxyl group and a 5' hydroxyl group at the end to be linked to the end of the first nucleic acid molecule containing the recognition sites. As disclosed herein, the methods can be performed using any number of nucleic acid molecules having various combinations of termini and ends.

[00147] Method of the invention may involve the use of nucleic acid molecule that comprises at least one topoisomerase. The topoisomerase may be, *e.g.*, a type I topoisomerase. More specifically, the type I topoisomerase may be a type IB topoisomerase. Where a type IB topoisomerase is used, the type IB topoisomerase may be a topoisomerase selected, *e.g.*, from the group consisting of eukaryotic nuclear type I topoisomerase and a poxvirus topoisomerase. Poxvirus topoisomerases may be produced by or isolated from a virus selected from the group consisting of vaccinia virus, Shope fibroma virus, ORF virus, fowlpox virus, molluscum contagiosum virus and *Amsacta moorei* entomopoxvirus.

[00148] The present invention includes methods for producing a polynucleotide construct that encodes a fusion protein that comprises one or more amino acid sequence tags, using, for example, recombinational cloning or topoisomerase-mediated cloning. The methods of the invention may also involve the use of a combination of recombinational cloning and topoisomerase-mediated cloning. For example, the invention includes methods comprising the successive use of one or more recombinational cloning steps followed by one or more topoisomerase-mediated cloning steps. Alternatively, the invention also includes methods comprising the successive use of one or more topoisomerase-mediated cloning steps followed by one or more

-51-

recombinational cloning steps. Alternatively, the invention includes methods comprising the use of recombinational cloning and topoisomerase-mediated cloning in the same cloning step.

[00149] One example of the use of topoisomerase-mediated cloning followed by recombinational cloning to produce a polynucleotide construct that encodes a fusion protein capable of being post-translationally modified or that is capable of being recognized by an antibody (or fragment thereof) or other specific binding reagent, is as follows. A first nucleic acid molecule comprising a nucleotide sequence of interest is mixed with a second nucleic acid molecule comprising: (i) at least a first topoisomerase recognition site flanked by (ii) at least a first recombination site, and (iii) at least a second topoisomerase recognition site flanked by (iv) at least a second recombination site, wherein said first and second recombination sites do not recombine with each other, and (v) at least one topoisomerase. The mixture is incubated under conditions such that said first nucleic acid molecule is inserted into said second nucleic acid molecule between said at least two topoisomerase recognition sites, thereby producing a first product polynucleotide construct. The first product polynucleotide construct is then brought into contact with a third nucleic acid molecule comprising: (i) at least a third and fourth recombination sites that do not recombine with each other and (ii) one or more nucleic acid sequences which encode one or more amino acid sequence tags. The first product polynucleotide construct is contacted with said third nucleic acid molecule under conditions favoring recombination between said first and third and between said second and fourth recombination sites, thereby producing a second product polynucleotide construct. According to this exemplary method, said second polynucleotide construct will encode a fusion protein comprising: (i) said amino acid sequence tag, and (ii) the amino acid sequence encoded by said nucleotide sequence of interest.

[00150] Another example of the use of topoisomerase-mediated cloning followed by recombinational cloning to produce a polynucleotide construct that encodes a fusion protein that comprises an amino acid sequence tag, is as follows: A first nucleic acid molecule comprising a nucleotide sequence of

-52-

interest is mixed with a second nucleic acid molecule comprising: (i) at least a first topoisomerase recognition site flanked by (ii) at least a first recombination site, and (iii) at least a second topoisomerase recognition site flanked by (iv) at least a second recombination site, wherein said first and second recombination sites do not recombine with each other, (v) one or more nucleic acid sequences which encode one or more amino acid sequence tags, and (vi) at least one topoisomerase. The mixture is incubated under conditions such that said first nucleic acid molecule is inserted into said second nucleic acid molecule between said at least two topoisomerase recognition sites, thereby producing a first product polynucleotide construct. The first product polynucleotide construct is then brought into contact with a third nucleic acid molecule comprising: (i) at least a third and fourth recombination sites that do not recombine with each other. The first product polynucleotide construct is contacted with said third nucleic acid molecule under conditions favoring recombination between said first and third and between said second and fourth recombination sites, thereby producing a second product polynucleotide construct. According to this exemplary method, said second polynucleotide construct will encode a fusion protein comprising: (i) said amino acid sequence tag, and (ii) the amino acid sequence encoded by said nucleotide sequence of interest.

[00151] The invention also includes host cells comprising one or more polynucleotide construct that encodes a fusion protein, *e.g.*, a fusion protein that comprises one or more amino acid sequence tags, wherein said polynucleotide construct is produced according to a method of the invention.

[00152] The nucleic acid molecules and methods of the invention can be used, *e.g.*, to produce a fusion protein comprising one or more amino acid sequence tags, and an amino acid sequence encoded by a nucleic acid sequence of interest. Accordingly, the present invention includes methods for producing fusion proteins comprising one or more amino acid tags. The methods of the invention can be used to produce fusion proteins *in vitro* or *in vivo*. When *in vivo* methods are used, the fusion protein can be produced in either eukaryotic

or prokaryotic cells. Methods for producing proteins *in vivo* and *in vitro* are well known in the art.

[00153] According to certain embodiments, the invention provides methods for producing a fusion protein that comprises one or more amino acid sequence tags, said methods comprising: (a) obtaining a host cell comprising a polynucleotide construct that encodes a fusion protein that comprises one or more amino acid sequence tags, said polynucleotide construct produced according to a method of the invention; and (b) culturing said host cell under conditions wherein said fusion protein is produced by said host cell. The precise conditions for producing a fusion protein in a host cell will vary, depending on the host cell used and the nature of the fusion protein being produced, and will be appreciated by those of ordinary skill in the art. In certain embodiments, the methods of the invention further comprise culturing said host cell under conditions wherein said fusion protein is post-translationally modified in said host cell. For example, the fusion protein may be biotinylated in said host cell.

[00154] In yet other embodiments, the methods may further comprise causing said fusion protein to be released from said host cell or treating said host cell such that said fusion protein is released from said host cell; and (b) contacting said fusion protein with a detecting composition comprising a molecule that is capable of interacting with said fusion protein. In an exemplary embodiment, the fusion protein will be a post-translationally modified fusion protein, *e.g.*, a biotinylated fusion protein, and said detecting composition will comprise avidin or an avidin analogue (including *e.g.*, streptavidin).

[00155] Methods for treating a host cell such that a protein, produced therein, is released from said host cell, are well known in the art and include, *e.g.*, chemical disruption of the cell and physical disruption of the cell including, *e.g.*, boiling, freezing, grinding, and combinations of chemical and physical disruption of the cell. Such methods include producing a protein extract from said host cell.

[00156] Details regarding the production and detection of fusion proteins that comprise one or more amino acid sequence tags, in general, are known in the

art. (See, e.g., Parrott, M.B. and Barry, M.A., *Biochem. Biophys. Res. Comm.* 281:993-1000 (2001), Parrott, M.B. and Barry, M.A., *Mol. Ther.* 1:96-104 (2000), U.S. Patent No. 5,252,466, and references cited therein).

[00157] The invention also includes methods for purifying, isolating or concentrating fusion proteins that are produced using the compositions and methods of the invention. In one embodiment, the invention includes methods for purifying, isolating or concentrating fusion proteins that have been post-translationally modified by a post-translational modification reaction, either *in vivo* or *in vitro*. In another embodiment, the invention includes methods for purifying, isolating or concentrating fusion proteins that comprise an amino acid sequence that is capable of being recognized by one or more antibody (or fragment thereof) or other specific reagents.

[00158] In an exemplary embodiment, the fusion proteins of the invention are purified, isolated or concentrated by bringing the fusion proteins into contact with a composition that is capable of interacting with the amino acid sequence tag and/or with a molecular entity that is attached to the amino acid sequence tag. Such compositions that interact specifically with an amino acid sequence tag include, e.g., "detecting compositions." As used herein, the term "detecting composition" is intended to mean any composition comprising a molecule that is capable of interacting with an amino acid sequence tag or with a molecular entity that is attached to an amino acid sequence tag, e.g., a molecule that is capable of interacting with a molecular entity that was attached to the amino acid sequence tag in a post-translational modification reaction. Such molecules that interact with amino acid sequence tags include, e.g., proteins and polypeptides, including, e.g., antibodies (or fragments thereof including fab fragments, fc fragments, etc) specific for the amino acid sequence tag. Particular exemplary molecules that can be attached to a detecting composition include avidin, streptavidin, and derivatives and analogs of those two compounds, as well as metal compounds (e.g., arsenites and thallium) that bind to dithiols such as lipoic acid (U.S. Patent No. 5,252,466), and antibodies (or fragments thereof) specific for epitopes such as, e.g., the FLAG epitope, the Myc epitope, the HA epitope, etc.



[00159] Detecting compositions may further comprise a surface (including, e.g., a solid and semi-solid surface), a matrix or a substrate, to which the molecule that is capable of interacting with particular amino acid sequence tag (or molecular entity attached thereto) is attached. Exemplary surfaces, matrices and substrates include, e.g., agarose beads, plastic beads, microscope coverslips, microscope slides, magnetic beads, glass beads or planar surfaces. The attachment may be, e.g., covalent or non-covalent. The types of surfaces, matrices and substrates to which a molecule that is capable of interacting with an amino acid sequence tag (or molecular entity attached thereto) may be attached are known in the art (see, e.g., Zou, H. *et al.*, *J. Biochem. Biophys. Methods* 49:1-3:199-240 (2001), Zusman, R. and Zusman, I., *J. Biochem. Biophys. Methods* 49:1-3:175-187 (2001)). Exemplary detecting compositions include agarose beads to which avidin, streptavidin, or derivatives/analog thereof, are attached.

[00160] In certain embodiments, the detecting composition may be used to identify, concentrate or purify a fusion protein by, e.g., mixing the detecting composition with a solution or composition comprising the fusion protein of interest, wherein the mixing takes place in batch (e.g., in a vessel such as a beaker, flask, bottle, test tube, petri dish, or other suitable container) or through a column containing the detecting composition. The detecting composition may alternatively be applied to a solution, to a cell (e.g., a permeabilized cell), or to any other substance that is known to contain or suspected of containing the fusion protein of interest.

[00161] In certain embodiments, the fusion proteins of the invention will be post-translationally modified fusion proteins, e.g., fusion proteins that have been biotinylated at the amino acid sequence tag. The biotinylated fusion protein can be purified, isolated or concentrated from a mixture of other proteins and molecules by bringing the biotinylated fusion protein into contact with, e.g., a detecting composition comprising a molecule that specifically interacts with biotin. Such molecules include, e.g., avidin and avidin derivatives such as streptavidin. The detecting composition may further comprise a surface or support matrix that can be physically removed from a

mixture of proteins and other molecules, *e.g.*, agarose beads, or other equivalent beads.

[00162] In other embodiments, the fusion protein that is produced using the methods and compositions of the invention will comprise an amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by an amino acid sequence tag, and on the other side by an amino acid sequence encoded by a nucleic acid sequence of interest. After purifying, isolating or concentrating such a fusion protein, the fusion protein can be treated with a protease to separate the amino acid sequence tag from the amino acid sequence encoded by a nucleic acid sequence of interest.

[00163] The invention also includes compositions or reaction mixtures comprising one or more nucleic acid molecule of the invention. The compositions or reaction mixtures may additionally comprise, one or more additional components selected from the group consisting of one or more topoisomerases, one or more host cells (*e.g.*, host cells that may be competent for uptake of nucleic acid molecules) one or more recombination proteins, one or more vectors, one or more nucleotides, one or more primers, and one or more polypeptides having polymerase activity.

[00164] The invention also provides kits comprising the isolated nucleic acid molecules of the invention, which may optionally comprise one or more additional components selected from the group consisting of one or more topoisomerases, one or more recombination proteins, one or more vectors, one or more nucleotides, one or more primers, one or more polypeptides having polymerase activity, one or more host cells (*e.g.*, host cells that may be competent for uptake of nucleic acid molecules), one or more antibody (or fragment thereof), and one or more detecting compositions, including, *e.g.*, one or more support matrices complexed with avidin or an avidin analog.

[00165] It will be readily apparent to one of ordinary skill in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are obvious and may be made without departing from the scope of the invention or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly

-57-

understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention.

### EXAMPLE 1

#### A Gateway™-Adapted Destination Vector for Cloning and Expression of Biotinylated Fusion Proteins in *E. coli*

[00166] This example describes the pET104-DEST expression vector (Fig. 1). pET104-DEST is a 7.6 kb vector adapted for use with the Gateway™ Technology, and is designed to allow for high-level, inducible expression of biotinylated recombinant fusion proteins in *E. coli* using the pET system. Biotinylated recombinant protein may then be easily detected or immobilized to a solid support for other downstream applications.

[00167] The pET system was originally developed by Studier and colleagues and takes advantage of the high activity and specificity of the bacteriophage T7 RNA polymerase to allow regulated expression of heterologous genes in *E. coli* from the T7 promoter (Rosenberg, A.H. *et al.*, *Gene* 56:125-135 (1987); Studier, F.W. and Moffatt, B.A., *J. Mol. Biol.* 189:113-130 (1986); Studier, F.W. *et al.*, *Meth. Enzymol.* 185:60-89 (1990)).

[00168] The pET104-DEST vector comprises the following elements:

- (a) T7lac promoter for high-level, IPTG-inducible expression of the gene of interest in *E. coli* (Dubendorff, J.W., and Studier, F.W., *J. Mol. Biol.* 219:45-59 (1991); ); Studier, F.W. *et al.*, *Meth. Enzymol.* 185:60-89 (1990));
- (b) Biotag™ to allow biotinylation of the recombinant protein of interest for easy detection or use in other applications;
- (c) Enterokinase (EK) recognition site for cleavage of the Biotag™ from the recombinant protein;
- (d) Two recombination sites, attR1 and attR2, downstream of the CMV promoter for recombinational cloning of the gene of interest from an entry clone;

- (e) Chloramphenicol resistance gene (CmR) located between the two *attR* sites for counterselection;
- (f) The *ccdB* gene located between the *attR* sites for negative selection;
- (g) *lacI* gene encoding the lac repressor to reduce basal transcription from the T7lac promoter in the pET104-DEST vector and from the *lacUV5* promoter in the *E. coli* chromosome;
- (h) Ampicillin resistance gene for selection in *E. coli*; and
- (i) pBR322 origin for low-copy replication and maintenance of the plasmid in *E. coli*.

[00169] The control plasmid, pET104/GW/*lacZ* (Fig. 2), can be used as a positive control for expression in *E. coli*. pET104/GW/*lacZ* was generated using the Gateway LR recombination reaction between an entry clone containing the *lacZ* gene and pET104-DEST.

[00170] To recombine a gene of interest into pET104-DEST, an entry clone containing a gene of interest will be obtained. Details relating to choosing an entry vector and constructing an entry clone are available in the art (*See, e.g.*, U.S. Patent No. 6,270,969).

[00171] pET104-DEST is an N-terminal fusion vector and contains an ATG initiation codon. A Shine-Dalgarno ribosome binding site (RBS) is included upstream of the initiation. The gene of interest in the entry clone must: (a) be in frame with the N-terminal Biotag™ after recombination; and (b) contain a stop codon.

[00172] The entry clone will contain, *e.g.*, *attL* sites flanking the gene of interest. Genes in an entry clone are transferred to the destination vector backbone by mixing the DNAs with, *e.g.*, the Gateway LR Clonase Enzyme Mix. The resulting LR recombination reaction is then transformed into *E. coli* (*e.g.*, TOP10 or DH5 $\alpha$ -T1R) and the expression clone is selected using ampicillin. Recombination between the *attR* sites on the destination vector and the *attL* sites on the entry clone replaces the chloramphenicol (CmR) gene and the *ccdB* gene with the gene of interest and results in the formation of *attB*

sites in the expression clone. Details for setting up the recombination reaction, transforming *E. coli*, and selecting for the expression clone, are available in the art.

[00173] The recombination region of the expression clone resulting from pET104-DEST x entry clone is depicted in Fig. 11. Features of the recombination region are as follows:

- (a) shaded regions correspond to those DNA sequences transferred from the entry clone into the pET104-DEST vector by recombination. Non-shaded regions are derived from the pET104-DEST vector;
- (b) bases 568 and 2230 of the pET104-DEST sequence are marked.
- (c) The biotin binding site is labeled with an asterisk (\*).

[00174] The Expression clone can be confirmed following recombination. The *ccdB* gene mutates at a very low frequency, resulting in a very low number of false positives. True expression clones will be ampicillin-resistant and chloramphenicol-sensitive. Transformants containing a plasmid with a mutated *ccdB* gene will be both ampicillin- and chloramphenicol-resistant. To check a putative expression clone, transformants can be tested for growth on LB plates containing 30 µg/ml chloramphenicol. A true expression clone should not grow in the presence of chloramphenicol.

[00175] The expression construct may also be sequenced to confirm that the gene of interest is in frame with the Biotag™. The priming sites indicated in Fig. 11 can be used to sequence the insert.

[00176] Expression of the recombinant fusion protein can be induced by first transforming the expression clone into an appropriate *E. coli* strain for protein expression, *e.g.*, BL21 cells. The transformant is then grown to mid-log in LB containing 100 µg/ml ampicillin or 50 µg/ml carbenicillin, and IPTG is added to a final concentration of 0.5-1 mM.

[00177] Expression of the recombinant fusion protein can be detected, *e.g.*, by western blot analysis using, *e.g.*, streptavidin-HRP or streptavidin-AP

conjugates, or an antibody (or fragment thereof) specific for the protein of interest.

[00178] The recombinant fusion protein can then be purified. The presence of the N-terminal Biotag™ in pET104-DEST allows the recombinant fusion protein to be biotinylated. Once biotinylated, the recombinant fusion protein can be purified by taking advantage of the strong association between biotin and avidin (and its analogs including streptavidin). For example, streptavidin agarose-conjugated beads can be used to purify the recombinant fusion protein. Other streptavidin conjugates can also be used.

[00179] A streptavidin-agarose resin can be used for affinity purification of recombinant fusion proteins containing the Biotag™. The resin can be constructed by covalently linking streptavidin to cross-linked agarose beads via a 15-atom hydrophilic spacer arm specifically designed to reduce non-specific binding and to ensure optimal binding of biotinylated molecules. Streptavidin is bound to a final concentration of 2-3 mg streptavidin per ml of packed resin.

[00180] Recombinant fusion proteins may be purified with streptavidin-agarose under native or denaturing conditions. Methods for purifying biotinylated proteins are known in the art.

[00181] pET104-DEST contains an enterokinase (EK) recognition site to allow removal of the Biotag™ from the recombinant fusion protein, if desired. After digestion with enterokinase, 11 amino acids will remain at the N-terminus of the protein (*see* Fig. 11). Methods for digestion with enterokinase are known in the art.

## EXAMPLE 2

### Directional TOPO Cloning of Blunt-End PCR Products into a Vector for Biotinylated Expression in *E. coli*

[00182] This example describes directional TOPO cloning using the pET104/D-TOPO vector (Fig. 3).

[00183] pET104/D-TOPO is a 5.9 kb vector designed to facilitate rapid, directional TOPO cloning of blunt-end PCR products for regulated and biotinylated expression in *E. coli*. The pET104/D-TOPO vector comprises the following elements:

- (a) T7lac promoter for high-level, IPTG-inducible expression of the gene of interest in *E. coli* (Dubendorff, J.W., and Studier, F.W., *J. Mol. Biol.* 219:45-59 (1991); ); Studier, F.W. *et al.*, *Meth. Enzymol.* 185:60-89 (1990));
- (b) Directional TOPO cloning site for rapid and efficient directional cloning of blunt-end PCR products;
- (c) Biotag™ to allow biotinylation of the recombinant protein of interest for easy detection or use in other applications;
- (d) Enterokinase (EK) recognition site for cleavage of the Biotag™ from the recombinant protein;
- (e) *lacI* gene encoding the lac repressor to reduce basal transcription from the T7lac promoter in the pET104/D-TOPO vector and from the *lacUV5* promoter in the *E. coli* chromosome;
- (f) Ampicillin resistance gene for selection in *E. coli*; and
- (g) pBR322 origin for low-copy replication and maintenance of the plasmid in *E. coli*.

[00184] The control plasmid, pET104/D/*lacZ* (Fig. 4), can be used as a positive control for expression in *E. coli*. The gene encoding  $\beta$ -galactosidase was directionally TOPO cloned into the pET104/D-TOPO vector.

[00185] Topoisomerase I from *Vaccinia* virus binds to duplex DNA at specific sites and cleaves the phosphodiester backbone after 5'-CCCTT in one strand (Shuman, S., *Proc. Natl. Acad. Sci. USA* 88:10104-10108 (1991)). The energy from the broken phosphodiester backbone is conserved by formation of a covalent bond between the 3' phosphate of the cleaved strand and a tyrosyl residue (Tyr-274) of topoisomerase I. The phospho-tyrosyl bond between the DNA and enzyme can subsequently be attacked by the 5' hydroxyl of the original cleaved strand, reversing the reaction and releasing topoisomerase

(Shuman, S., *J. Biol. Chem.* 269:32678-32684 (1994)). TOPO cloning exploits this reaction to efficiently clone PCR products.

[00186] Directional joining of double-strand DNA using TOPO-charged oligonucleotides occurs by adding a 3' single-stranded end (overhang) to the incoming DNA (Cheng, C. and Shuman, S., *Mol. Cell. Biol.* 20:8059-8068 (2000)). This single-stranded overhang is identical to the 5' end of the TOPO-charged DNA fragment. A 4 nucleotide overhang sequence has been added to the TOPO-charged DNA and the TOPO system has been adapted to a "whole vector" format.

[00187] In this system, PCR products are directionally cloned by adding four bases to the forward primer (CACC). The overhang in the cloning vector (GTGG) invades the 5' end of the PCR product, anneals to the added bases, and stabilizes the PCR product in the correct orientation (*see* Fig. 12). Inserts can be cloned in the correct orientation with efficiencies equal to or greater than 90%.

[00188] The general steps required to clone and express a blunt-end PCR product are illustrated in Fig. 13.

[00189] The following factors should be considered when designing the forward PCR primer:

- (a) To enable directional cloning, the forward PCR primer must contain the sequence, CACC, at the 5' end of the primer. The 4 nucleotides, CACC, base pair with the overhang sequence, GTGG, in the pET104/D-TOPO vector.
- (b) To include the N-terminal Biotag™, it is important that the forward PCR primer be designed such that the gene of interest is in frame with the Biotag™. The initiation ATG codon is not needed. A Shine-Dalgarno ribosome binding site (RBS) is included upstream of the ATG in the N-terminal tag to ensure optimal spacing for proper translation initiation.
- (c) At least six non-native amino acids will be present between the EK cleavage site and the start of the gene of interest.



- (d) If it is desired to express the protein with a native N-terminus (*i.e.*, with out the Biotag™), the forward PCR primer should be designed to include: (i) a stop codon to terminate the Biotag™, and (ii) a second ribosome binding site (AGGAGG) 9-10 base pairs 5' of the initial ATG codon of the protein.

[00190] The following factors should be considered when designing the reverse PCR primer:

- (a) It is important to include a stop codon in the reverse primer or the reverse primer should be designed to hybridize downstream of the native stop codon.
- (b) To ensure that the PCR product clones directionally with high efficiency, the reverse PCR primer must not be complementary to the overhang sequence GTGG at the 5' end. A one base pair mismatch can reduce the directional cloning efficiency from 90% to 75%, and may increase the chances of the open reading frame cloning in the opposite orientation.

[00191] The diagram depicted in Fig. 14 is useful for designing suitable PCR primers to clone an express a PCR product using pET104/D-TOPO. The biotin binding site is designated with an asterisk (\*).

[00192] Once a desired PCR product has been produced, it can then be TOPO cloned into the pET104/D-TOPO vector. The recombinant vector can then be transformed into an appropriate *E. coli* strain.

[00193] It has been found that inclusion of salt (*e.g.*, 250 mM NaCl, 10 mM MgCl<sub>2</sub>) in the TOPO cloning reaction may result in an increase in the number of transformants. Therefore, it is recommended that salt be added to the TOPO cloning reaction.

[00194] Table III describes how to set up a TOPO cloning reaction (6 µl) for eventual transformation into either chemically competent *E. coli* or electrocompetent *E. coli*.

TABLE III

Setting up a TOPO Cloning Reaction

-64-

Reagents	Chemically competent <i>E. coli</i>	Electrocompetent <i>E. coli</i>
Fresh PCR product	0.5 to 4.0 $\mu$ l	0.5 to 4.0 $\mu$ l
Salt solution	1 $\mu$ l	--
Sterile water	Add to a final volume of 5 $\mu$ l	Add to a final volume of 5 $\mu$ l
TOPO vector	1 $\mu$ l	1 $\mu$ l

[00195] Mix reaction gently and incubate for 5 minutes at room temperature (22-23°C). For most applications, 5 minutes will yield sufficient colonies for analysis. Depending on the circumstances, the length of the TOPO cloning reaction can be varied from 30 seconds to 30 minutes. For routine subcloning of PCR products, 30 seconds may be sufficient. For large PCR products (>1 kb) or if a pool of PCR products is being cloned, increasing the reaction time may yield more colonies.

[00196] Place the reaction on ice or store the TOPO cloning reaction at -20°C overnight.

[00197] Once the TOPO cloning reaction has been performed, the pET104/D-TOPO construct will be transformed into competent *E. coli*. Methods for transforming *E. coli* with nucleic acids are known in the art.

[00198] Transformants can be analyzed by isolating plasmid DNA from transformant colonies. The isolated plasmid DNA can be checked by restriction analysis to confirm the presence and correct orientation of the insert. Additionally, the construct can be sequenced to confirm that the gene of interest is in frame with the N-terminal Biotag™. Forward and T7 reverse primers can be used to sequence the insert. Positive transformants can also be analyzed by PCR.

[00199] Expression of the recombinant fusion protein can be induced by first transforming the expression clone into an appropriate *E. coli* strain for protein expression, e.g., BL21 cells. The transformant is then grown to mid-log in LB containing 100  $\mu$ g/ml ampicillin or 50  $\mu$ g/ml carbenicillin, and IPTG is added to a final concentration of 0.5-1 mM.

- [00200] Expression of the recombinant fusion protein can be detected, *e.g.*, by western blot analysis using, *e.g.*, streptavidin-HRP or streptavidin-AP conjugates, or an antibody (or fragment thereof) specific for the protein of interest.
- [00201] The recombinant fusion protein can then be purified. The presence of the N-terminal Biotag™ in pET104/D-TOPO allows the recombinant fusion protein to be biotinylated. Once biotinylated, the recombinant fusion protein can be purified by taking advantage of the strong association between biotin and avidin (and its analogs including streptavidin). For example, streptavidin agarose-conjugated beads can be used to purify the recombinant fusion protein. Other streptavidin conjugates can also be used.
- [00202] A streptavidin-agarose resin can be used for affinity purification of recombinant fusion proteins containing the Biotag™. The resin can be constructed by covalently linking streptavidin to cross-linked agarose beads via a 15-atom hydrophilic spacer arm specifically designed to reduce non-specific binding and to ensure optimal binding of biotinylated molecules. Streptavidin is bound to a final concentration of 2-3 mg streptavidin per ml of packed resin.
- [00203] Recombinant fusion proteins may be purified with streptavidin-agarose under native or denaturing conditions. Methods for purifying biotinylated proteins are known in the art.
- [00204] pET104/D-TOPO contains an enterokinase (EK) recognition site to allow removal of the Biotag™ from the recombinant fusion protein, if desired. After digestion with enterokinase, 6 amino acids will remain at the N-terminus of the protein (*see* Fig. 14). Methods for digestion with enterokinase are known in the art.

### EXAMPLE 3

A Gateway-Adapted Destination Vector for Cloning and Expression of Biotinylated Fusion Proteins in Mammalian Cells

[00205] This example describes the pcDNA/Biotag™-DEST vector (Fig. 5). pcDNA6/Biotag™-DEST is a 7.0 kb vector adapted for use with the Gateway Technology, and is designed to allow high-level expression of biotinylated recombinant fusion proteins in mammalian cells. Biotinylated recombinant protein may then be easily detected or immobilized to a solid support for other downstream applications.

[00206] The pcDNA6/Biotag™-DEST vector contains the following elements:

- (a) The human cytomegalovirus (CMV) immediate early enhancer/promoter for high level constitutive expression of the gene of interest in a wide range of mammalian cells (Andersson, S. *et al.*, *J. Biol. Chem.* 264:8222-8229 (1989); Boshart, M. *et al.*, *Cell* 41:521-530 (1985); Nelson, J.A. *et al.*, *Molec. Cell Biol.* 7:4125-4129 (1987));
- (b) Biotag™ to allow biotinylation of the recombinant protein of interest for easy detection or use in other applications.
- (c) Enterokinase (EK) recognition site for cleavage of the Biotag™ from the recombinant protein;
- (d) Two recombination sites, *attR1* and *attR2*, downstream of the CMV promoter for recombinational cloning of the gene of interest from an entry clone;
- (e) Chloramphenicol resistance gene (CmR) located between the two *attR* sites for counterselection;
- (f) The *ccdB* gene located between the *attR* sites for negative selection;
- (g) Blasticidin (*bsd*) resistance gene for selection of stable cell lines using blasticidin;
- (h) Ampicillin resistance gene for selection in *E. coli*; and
- (i) pUC origin for high-copy replication and maintenance of the plasmid in *E. coli*.

[00207] The control plasmid, pcDNA6/Biotag™-GW/*lacZ* (Fig. 6), can be used as a positive control for transfection and expression in the mammalian cell line of choice. pcDNA6/Biotag™-GW/*lacZ* was generated using the Gateway LR recombination reaction between an entry clone containing the *lacZ* gene and pcDNA6/Biotag™-DEST.

[00208] To recombine a gene of interest into pcDNA6/Biotag™-DEST, an entry clone containing the gene of interest must first be obtained. Details relating to choosing an entry vector and constructing an entry clone are available in the art (*See, e.g.*, U.S. Patent No. 6,270,969).

[00209] pcDNA6/Biotag™-DEST is an N-terminal fusion vector and contains an ATG initiation codon in the context of a Kozak consensus sequence to ensure optimal translation initiation. The gene of interest in the entry clone must: (a) be in frame with the N-terminal Biotag™ after recombination; and (b) contain a stop codon.

[00210] The entry clone will contain, *e.g.*, *attL* sites flanking the gene of interest. Genes in an entry clone are transferred to the destination vector backbone by mixing the DNAs with, *e.g.*, the Gateway LR Clonase Enzyme Mix. The resulting LR recombination reaction is then transformed into *E. coli* (*e.g.*, TOP10 or DH5α-T1R) and the expression clone is selected using ampicillin. Recombination between the *attR* sites on the destination vector and the *attL* sites on the entry clone replaces the chloramphenicol (CmR) gene and the *ccdB* gene with the gene of interest and results in the formation of *attB* sites in the expression clone. Details for setting up the recombination reaction, transforming *E. coli*, and selecting for the expression clone, are available in the art.

[00211] The recombination region of the expression clone resulting from pcDNA6/Biotag™-DEST x entry clone is depicted in Fig. 15. Features of the recombination region are as follows:

- (a) shaded regions correspond to those DNA sequences transferred from the entry clone into the pcDNA6/Biotag™-DEST vector by

-68-

recombination. Non-shaded regions are derived from the pcDNA6/Biotag™-DEST vector;

(b) bases 1191 and 2853 of the pcDNA6/Biotag™-DEST sequence are marked.

(c) The biotin binding site is labeled with an asterisk (\*).

(d) Potential stop codons are underlined.

[00212] The Expression clone can be confirmed following recombination. The *ccdB* gene mutates at a very low frequency, resulting in a very low number of false positives. True expression clones will be ampicillin-resistant and chloramphenicol-sensitive. Transformants containing a plasmid with a mutated *ccdB* gene will be both ampicillin- and chloramphenicol-resistant. To check a putative expression clone, transformants can be tested for growth on LB plates containing 30 µg/ml chloramphenicol. A true expression clone should not grow in the presence of chloramphenicol.

[00213] The expression construct may also be sequenced to confirm that the gene of interest is in frame with the Biotag™. The priming sites indicated in Fig. 15 can be used to sequence the insert.

[00214] Before expression of the recombinant fusion protein can be induced, the expression clone must first be transfected into the mammalian cells of choice. Methods for transfecting mammalian cells are known in the art. Exemplary methods of transfection include calcium phosphate, lipid-mediated, and electroporation. Following transfection, a stable cell line can be generated.

[00215] Expression of the recombinant fusion protein can be assayed from either transiently transfected cells or stable cell lines. Expression of the recombinant fusion protein can be detected, e.g., by western blot analysis using, e.g., streptavidin-HRP or streptavidin-AP conjugates, or an antibody (or fragment thereof) specific for the protein of interest.

[00216] The recombinant fusion protein can then be purified. The presence of the N-terminal Biotag™ in pcDNA6/Biotag™-DEST allows the recombinant fusion protein to be biotinylated. Once biotinylated, the recombinant fusion

protein can be purified by taking advantage of the strong association between biotin and avidin (and its analogs including streptavidin). For example, streptavidin agarose-conjugated beads can be used to purify the recombinant fusion protein. Other streptavidin conjugates can also be used.

[00217] A streptavidin-agarose resin can be used for affinity purification of recombinant fusion proteins containing the Biotag™. The resin can be constructed by covalently linking streptavidin to cross-linked agarose beads via a 15-atom hydrophilic spacer arm specifically designed to reduce non-specific binding and to ensure optimal binding of biotinylated molecules. Streptavidin is bound to a final concentration of 2-3 mg streptavidin per ml of packed resin.

[00218] Recombinant fusion proteins may be purified with streptavidin-agarose under native or denaturing conditions. Methods for purifying biotinylated proteins are known in the art.

[00219] pcDNA6/Biotag™-DEST contains an enterokinase (EK) recognition site to allow removal of the Biotag™ from the recombinant fusion protein, if desired. After digestion with enterokinase, 12 amino acids will remain at the N-terminus of the protein (*see* Fig. 15). Methods for digestion with enterokinase are known in the art.

#### EXAMPLE 4

##### Directional TOPO Cloning of Blunt-End PCR Products into a Vector for Biotinylated Expression in Mammalian Cells

[00220] This example describes directional TOPO cloning using the pcDNA6/Biotag™/D-TOPO vector (Fig. 7).

[00221] pcDNA6/Biotag™/D-TOPO is a 5.3 kb expression vector designed to facilitate rapid directional cloning of blunt-end PCR products for high-level expression and biotinylation in mammalian cells. Biotinylated recombinant protein may then be easily detected or immobilized to a solid support for other

downstream applications. The pcDNA6/Biotag™/D-TOPO vector comprises the following elements:

- (a) The human cytomegalovirus (CMV) immediate early enhancer/promoter for high level constitutive expression of the gene of interest in a wide range of mammalian cells (Andersson, S. *et al.*, *J. Biol. Chem.* 264:8222-8229 (1989); Boshart, M. *et al.*, *Cell* 41:521-530 (1985); Nelson, J.A. *et al.*, *Molec. Cell Biol.* 7:4125-4129 (1987));
- (b) Biotag™ to allow biotinylation of the recombinant protein of interest for easy detection or use in other applications;
- (c) Enterokinase (EK) recognition site for cleavage of the Biotag™ from the recombinant protein;
- (d) TOPO cloning site for rapid and efficient directional cloning of blunt-end PCR products;
- (e) Blasticidin (*bsd*) resistance gene for selection of stable cell lines using blasticidin.

[00222] The control plasmid, pcDNA6/Biotag™/*lacZ* (Fig. 8), can be used as a positive control for expression in *E. coli*. The gene encoding  $\beta$ -galactosidase was directionally TOPO cloned into the pcDNA6/Biotag™/D-TOPO vector.

[00223] The theory behind topoisomerase cloning is described under Example 2, *supra*.

[00224] The general steps required to clone and express a blunt-end PCR product are illustrated in Fig. 16.

[00225] The following factors should be considered when designing the forward PCR primer:

- (e) To enable directional cloning, the forward PCR primer must contain the sequence, CACC, at the 5' end of the primer. The 4 nucleotides, CACC, base pair with the overhang sequence, GTGG, in the pcDNA6/Biotag™/D-TOPO vector.
- (f) To include the N-terminal Biotag™, it is important that the forward PCR primer be designed such that the gene of interest is



-71-

in frame with the Biotag™. The initiation ATG codon is not needed.

- (g) If it is desired to express the protein with a native N-terminus (*i.e.*, with out the Biotag™), the forward PCR primer should be designed to include: (i) a stop codon to terminate the Biotag™, and (ii) the ATG initiation codon within the context of a Kozak consensus sequence to ensure optimal translation initiation.

[00226] The following factors should be considered when designing the reverse PCR primer:

- (c) It is important to include a stop codon in the reverse primer or the reverse primer should be designed to hybridize downstream of the native stop codon.
- (d) To ensure that the PCR product clones directionally with high efficiency, the reverse PCR primer must not be complementary to the overhang sequence GTGG at the 5' end. A one base pair mismatch can reduce the directional cloning efficiency from 90% to 75%, and may increase the chances of the open reading frame cloning in the opposite orientation.

[00227] The diagram depicted in Fig. 17 is useful for designing suitable PCR primers to clone an express a PCR product using pcDNA6/Biotag™/D-TOPO. The biotin binding site is designated with an asterisk (\*).

[00228] Once a desired PCR product has been produced, it can then be TOPO cloned into the pcDNA6/Biotag™/D-TOPO vector. The recombinant vector can then be transformed into an appropriate *E. coli* strain.

[00229] It has been found that inclusion of salt (*e.g.*, 250 mM NaCl, 10 mM MgCl<sub>2</sub>) in the TOPO cloning reaction may result in an increase in the number of transformants. Therefore, it is recommended that salt be added to the TOPO cloning reaction.

[00230] Table IV describes how to set up a TOPO cloning reaction (6 µl) for eventual transformation into either chemically competent *E. coli* or electrocompetent *E. coli*.

TABLE IV  
Setting up a TOPO Cloning Reaction

Reagents	Chemically competent <i>E. coli</i>	Electrocompetent <i>E. coli</i>
Fresh PCR product	0.5 to 4.0 $\mu$ l	0.5 to 4.0 $\mu$ l
Salt solution	1 $\mu$ l	--
Sterile water	Add to a final volume of 5 $\mu$ l	Add to a final volume of 5 $\mu$ l
TOPO vector	1 $\mu$ l	1 $\mu$ l

[00231] Mix reaction gently and incubate for 5 minutes at room temperature (22-23°C). For most applications, 5 minutes will yield sufficient colonies for analysis. Depending on the circumstances, the length of the TOPO cloning reaction can be varied from 30 seconds to 30 minutes. For routine subcloning of PCR products, 30 seconds may be sufficient. For large PCR products (>1 kb) or if a pool of PCR products is being cloned, increasing the reaction time may yield more colonies.

[00232] Place the reaction on ice or store the TOPO cloning reaction at -20°C overnight.

[00233] Once the TOPO cloning reaction has been performed, pcDNA6/Biotag™/D-TOPO construct will be transformed into competent *E. coli*. Methods for transforming *E. coli* with nucleic acids are known in the art.

[00234] Transformants can be analyzed by isolating plasmid DNA from transformant colonies. The isolated plasmid DNA can be checked by restriction analysis to confirm the presence and correct orientation of the

-73-

insert. Additionally, the construct can be sequenced to confirm that the gene of interest is in frame with the N-terminal Biotag™. Forward and T7 reverse primers can be used to sequence the insert. Positive transformants can also be analyzed by PCR.

[00235] Before expression of the recombinant fusion protein can be induced, the expression clone must first be transfected into the mammalian cells of choice. Methods for transfecting mammalian cells are known in the art. Exemplary methods of transfection include calcium phosphate, lipid-mediated, and electroporation. Following transfection, a stable cell line can be generated.

[00236] Expression of the recombinant fusion protein can be assayed from either transiently transfected cells or stable cell lines. Expression of the recombinant fusion protein can be detected, *e.g.*, by western blot analysis using, *e.g.*, streptavidin-HRP or streptavidin-AP conjugates, or an antibody (or fragment thereof) specific for the protein of interest.

[00237] The recombinant fusion protein can then be purified. The presence of the N-terminal Biotag™ in pcDNA6/Biotag™/D-TOPO allows the recombinant fusion protein to be biotinylated. Once biotinylated, the recombinant fusion protein can be purified by taking advantage of the strong association between biotin and avidin (and its analogs including streptavidin). For example, streptavidin agarose-conjugated beads can be used to purify the recombinant fusion protein. Other streptavidin conjugates can also be used.

[00238] A streptavidin-agarose resin can be used for affinity purification of recombinant fusion proteins containing the Biotag™. The resin can be constructed by covalently linking streptavidin to cross-linked agarose beads via a 15-atom hydrophilic spacer arm specifically designed to reduce non-specific binding and to ensure optimal binding of biotinylated molecules. Streptavidin is bound to a final concentration of 2-3 mg streptavidin per ml of packed resin.

[00239] Recombinant fusion proteins may be purified with streptavidin-agarose under native or denaturing conditions. Methods for purifying biotinylated proteins are known in the art.

[00240] pcDNA6/Biotag™/D-TOPO contains an enterokinase (EK) recognition site to allow removal of the Biotag™ from the recombinant fusion protein, if desired. After digestion with enterokinase, 13 amino acids will remain at the N-terminus of the protein (see Fig. 17). Methods for digestion with enterokinase are known in the art.

### EXAMPLE 5

#### A Gateway™-Adapted Destination Vector for the Stable Expression of Biotinylated Fusion Proteins in *Drosophila* Schneider 2 Cells

[00241] This example describes the pMT/Biotag™-DEST vector (Fig. 9). pMT/Biotag™-DEST is a 5.4 kb vector adapted for use with the Gateway Technology, and is designed to allow high-level expression of biotinylated recombinant fusion proteins in *Drosophila* Schneider 2 (S2) cells. Biotinylated recombinant protein may then be easily detected or immobilized to a solid support for other downstream applications.

[00242] The pMT/Biotag™-DEST vector contains the following elements:

- (a) The *Drosophila* metallothionein (MT) promoter for high-level, metal-inducible expression of a gene of interest in S2 cells.
- (b) Biotag™ to allow biotinylation of the recombinant protein of interest for easy detection or use in other applications.
- (c) Two recombination sites, *attR1* and *attR2*, downstream of the MT promoter for recombinational cloning of the gene of interest form an entry clone.
- (d) Chloramphenicol resistance gene (CmR) located between the *attR* sites for counterselection.
- (e) The *ccdB* gene located between the *attR* sites for negative selection.

-75-

(f) pUC origin for high-copy replication and maintenance of the plasmid in *E. coli*.

(g) Ampicillin resistance gene for selection in *E. coli*.

[00243] The control plasmid, pMT/Biotag<sup>TM</sup>/GW-*lacZ* (Fig. 10), can be used as a positive control for transfection and expression in the mammalian cell line of choice. pMT/Biotag<sup>TM</sup>/GW-*lacZ* was generated using the Gateway LR recombination reaction between an entry clone containing the *lacZ* gene and pMT/Biotag<sup>TM</sup>-DEST.

[00244] To recombine a gene of interest into pMT/Biotag<sup>TM</sup>-DEST, an entry clone containing the gene of interest must first be obtained. Details relating to choosing an entry vector and constructing an entry clone are available in the art (*See, e.g.*, U.S. Patent No. 6,270,969).

[00245] pMT/Biotag<sup>TM</sup>-DEST is an N-terminal fusion vector and contains an ATG initiation codon. The gene of interest in the entry clone must: (a) be in frame with the N-terminal Biotag<sup>TM</sup> after recombination; and (b) contain a stop codon.

[00246] The entry clone will contain, *e.g.*, *attL* sites flanking the gene of interest. Genes in an entry clone are transferred to the destination vector backbone by mixing the DNAs with, *e.g.*, the Gateway LR Clonase Enzyme Mix. The resulting LR recombination reaction is then transformed into *E. coli* (*e.g.*, TOP10 or DH5 $\alpha$ -T1R) and the expression clone is selected using ampicillin. Recombination between the *attR* sites on the destination vector and the *attL* sites on the entry clone replaces the chloramphenicol (CmR) gene and the *ccdB* gene with the gene of interest and results in the formation of *attB* sites in the expression clone. Details for setting up the recombination reaction, transforming *E. coli*, and selecting for the expression clone, are available in the art.

[00247] The recombination region of the expression clone resulting from pMT/Biotag<sup>TM</sup>-DEST x entry clone is depicted in Fig. 18. Features of the recombination region are as follows:

-76-

- (e) shaded regions correspond to those DNA sequences transferred from the entry clone into the pMT/Biotag<sup>TM</sup>-DEST vector by recombination. Non-shaded regions are derived from the pMT/Biotag<sup>TM</sup>-DEST vector;
- (f) bases 1135 and 2797 of the pMT/Biotag<sup>TM</sup>-DEST sequence are marked.
- (g) The biotin binding site is labeled with an asterisk (\*).
- (h) Potential stop codons are underlined.

[00248] The basic steps needed to clone and express a protein using pMT/Biotag<sup>TM</sup>-DEST are as follows:

- (a) Establish a culture of S2 cells from supplied frozen stock.
- (b) Choose a Gateway entry vector and generate an entry clone containing the gene of interest.
- (c) Perform an LR recombination reaction between the entry clone containing the gene of interest and the pMT/Biotag<sup>TM</sup>-DEST vector. Transform *E. coli* and select for the expression clone.
- (d) Isolate plasmid DNA.
- (e) Transiently transfect S2 cells.
- (f) Induce, if necessary, and assay for expression of the protein.
- (g) Create stable cell lines expressing the protein of interest by cotransfecting the recombinant expression vector with a selection vector, pCoHygro (Fig. 19) or pCoBlast (Fig. 20), and select with the appropriate concentration of hygromycin-B or blasticidin, respectively.
- (h) Induce if necessary, and assay for expression of the protein.
- (i) Scale up expression, if desired.

[00249] Expression of the recombinant fusion protein can be detected, e.g., by western blot analysis using, e.g., streptavidin-HRP or streptavidin-AP conjugates, or an antibody (or fragment thereof) specific for the protein of interest.

- [00250] The recombinant fusion protein can then be purified. The presence of the N-terminal Biotag™ in pMT/Biotag™-DEST allows the recombinant fusion protein to be biotinylated. Once biotinylated, the recombinant fusion protein can be purified by taking advantage of the strong association between biotin and avidin (and its analogs including streptavidin). For example, streptavidin agarose-conjugated beads can be used to purify the recombinant fusion protein. Other streptavidin conjugates can also be used.
- [00251] A streptavidin-agarose resin can be used for affinity purification of recombinant fusion proteins containing the Biotag™. The resin can be constructed by covalently linking streptavidin to cross-linked agarose beads via a 15-atom hydrophilic spacer arm specifically designed to reduce non-specific binding and to ensure optimal binding of biotinylated molecules. Streptavidin is bound to a final concentration of 2-3 mg streptavidin per ml of packed resin.
- [00252] Recombinant fusion proteins may be purified with streptavidin-agarose under native or denaturing conditions. Methods for purifying biotinylated proteins are known in the art.
- [00253] pMT/Biotag™-DEST contains an enterokinase (EK) recognition site to allow removal of the Biotag™ from the recombinant fusion protein, if desired. After digestion with enterokinase, 11 amino acids will remain at the N-terminus of the protein (*see* Fig. 18). Methods for digestion with enterokinase are known in the art.
- [00254] Having now fully described the present invention in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious to one of ordinary skill in the art that the same can be performed by modifying or changing the invention within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any specific embodiment thereof, and that such modifications or changes are intended to be encompassed within the scope of the appended claims.

[00255] All publications, patents and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains, and are herein incorporated by reference to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference.



## WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising:
  - (a) one or more recombination sites; and
  - (b) one or more nucleic acid sequences which encode an amino acid sequence tag.
2. The isolated nucleic acid molecule of claim 1, further comprising at least one additional nucleic acid sequence selected from the group consisting of a selectable marker, a cloning site, a restriction site, a promoter, an operator, an operon, a nucleotide sequence encoding a gene product which allows for negative selection, an origin of replication, a nucleotide sequence which encodes a repressor of at least one promoter, and a gene or partial gene.
3. The isolated nucleic acid molecule of claim 1, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more recombination sites, thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest.
4. The isolated nucleic acid molecule of claim 1, further comprising a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases.
5. The isolated nucleic acid molecule of claim 4, wherein said amino acid sequence that is capable of being cleaved by one or more proteases is an amino acid sequence that is capable of being cleaved by enterokinase.
6. The isolated nucleic acid molecule of claim 4, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one

-80-

or more recombination sites thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid tag, and on the other side by (iii) the amino acid sequence encoded by said nucleic acid sequence of interest.

7. The nucleic acid molecule of claim 1, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.
8. The isolated nucleic acid molecule of claim 7, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being post-translationally modified by biotinylation, attachment of 4-phosphopanthetheine, attachment of lipoic acid or attachment of flavins.
9. The isolated nucleic acid molecule of claim 7, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being biotinylated.
10. The isolated nucleic acid molecule of claim 9, wherein said amino acid sequence that is capable of being biotinylated is all or a portion of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, all or a portion of the *Propionibacterium shermanii* transcarboxylase 1.3S subunit, or all or a portion of the *Escherichia coli* biotin carboxyl carrier protein component of acetyl-CoA carboxylase.
11. The isolated nucleic acid molecule of claim 9, wherein said amino acid sequence that is capable of being biotinylated is a portion of the C-

-81-

terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit.

12. The isolated nucleic acid molecule of claim 11, wherein said amino acid sequence that is capable of being biotinylated is the BIOTAG™.
13. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule is a circular molecule.
14. The isolated nucleic acid molecule of claim 1, wherein said nucleic acid molecule comprises two or more recombination sites.
15. The isolated nucleic acid molecule of claim 1, wherein said recombination sites are selected from the group consisting of: (a) *attB* sites, (b) *attP* sites, (c) *attL* sites, (d) *attR* sites, (e) *lox* sites, (f) *psi* sites, (g) *dif* sites, (h) *cer* sites, (i) *frt* sites, and mutants, variants, and derivatives of the recombination sites of (a), (b), (c), (d), (e), (f), (g), (h), or (i) which retain the ability to undergo recombination.
16. A vector comprising the isolated nucleic acid molecule of claim 1.
17. A host cell comprising the isolated nucleic acid molecule of claim 1.
18. A host cell comprising the vector of claim 16.
19. A method of producing a polynucleotide construct that encodes a fusion protein that comprises an amino acid sequence tag, said method comprising:
  - (a) obtaining a first nucleic acid molecule comprising a nucleotide sequence of interest flanked by at least a first and at least a second recombination sites that do not recombine with each other;

-82-

- (b) obtaining a second nucleic acid molecule comprising: (i) at least a third and fourth recombination sites that do not recombine with each other; and (ii) one or more nucleic acid sequences which encode an amino acid sequence tag; and
  - (c) contacting said first nucleic acid molecule with said second nucleic acid molecule under conditions favoring recombination between said first and third and between said second and fourth recombination sites, thereby producing a product polynucleotide construct;  
wherein said product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleotide acid sequence of interest.
20. The method of claim 19, wherein said second nucleic acid molecule further comprises a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases; and  
wherein said product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid sequence tag, and on the other side by (iii) the amino acid sequence encoded by said nucleotide sequence of interest.
21. The method of claim 20, wherein said amino acid sequence that is capable of being cleaved by one or more proteases is an amino acid sequence that is capable of being cleaved by enterokinase.
22. The method of claim 19, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.
23. The method of claim 22, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being post-translationally modified by biotinylation, attachment

of 4-phosphopanthetheine, attachment of lipoic acid or attachment of flavins.

24. The method of claim 22, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being biotinylated.
25. The method of claim of claim 24, wherein said amino acid sequence that is capable of being biotinylated is all or a portion of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, all or a portion of the *Propionibacterium shermanii* transcarboxylase 1.3S subunit, or all or a portion of the *Escherichia coli* biotin carboxyl carrier protein component of acetyl-CoA carboxylase.
26. The method of claim of claim 24, wherein said amino acid sequence that is capable of being biotinylated is a portion of the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit.
27. The method of claim 26, wherein said amino acid sequence that is capable of being biotinylated is the BIOTAG™.
28. The method of claim 19, wherein said second nucleic acid molecule is a vector.
29. The method of claim 19, wherein said first nucleic acid molecule is a circular nucleic acid molecule.
30. The method of claim 19, wherein said first nucleic acid molecule is a linear nucleic acid molecule.

31. The method of claim 30, wherein said first nucleic acid molecule is a PCR product.
32. The method of claim 19, further comprising inserting said product polynucleotide construct into a host cell.
33. The method of claim 20, further comprising inserting said product polynucleotide construct into a host cell.
34. The method of claim 19, wherein said second nucleic acid molecule comprises at least one additional nucleic acid sequence selected from the group consisting of a selectable marker, a cloning site, a restriction site, a promoter, an operator, an operon, a nucleotide sequence encoding a gene product which allows for negative selection, an origin of replication, a nucleotide sequence which encodes a repressor of at least one promoter, and a gene or partial gene.
35. The method of claim 19, wherein said first, second, third and fourth recombination sites are selected from the group consisting of: (a) *attB* sites, (b) *attP* sites, (c) *attL* sites, (d) *attR* sites, (e) *lox* sites, (f) *psi* sites, (g) *dif* sites, (h) *cer* sites, (i) *frt* sites, and mutants, variants, and derivatives of the recombination sites of (a), (b), (c), (d), (e), (f), (g), (h), or (i) which retain the ability to undergo recombination.
36. The method of claim 19, wherein said first and said second nucleic acid molecules are combined in the presence of at least one recombination protein.
37. The method of claim 36, wherein said recombination protein is selected from the group consisting of: (a) Cre, (b) Int, (c) IHF, (d) Xis, (e) Fis, (f) Hin, (g) Gin, (h) Cin, (i) Tn3 resolvase, (j) TndX, (k) XerC, and (l) XerD.

38. The method of claim 36, wherein said recombination protein is Cre.
39. An isolated nucleic acid molecule comprising:
- (a) one or more topoisomerase recognition sites and/or one or more topoisomerases; and
  - (b) one or more nucleic acid sequences which encode an amino acid sequence tag.
40. The isolated nucleic acid molecule of claim 39, further comprising at least one additional nucleic acid sequence selected from the group consisting of a selectable marker, a cloning site, a restriction site, a promoter, an operator, an operon, a nucleotide sequence encoding a gene product which allows for negative selection, an origin of replication, a nucleotide sequence which encodes a repressor of at least one promoter, and a gene or partial gene.
41. The isolated nucleic acid molecule of claim 39, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotide of the position of said one or more topoisomerases, thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest.
42. The isolated nucleic acid molecule of claim 39, further comprising a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases.
43. The isolated nucleic acid molecule of claim 42, wherein said amino acid sequence that is capable of being cleaved by one or more proteases is an amino acid sequence that is capable of being cleaved by enterokinase.

44. The isolated nucleic acid molecule of claim 42, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at the position of said one or more topoisomerases thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid tag, and on the other side by (iii) the amino acid sequence encoded by said nucleic acid sequence of interest.
45. The isolated nucleic acid molecule of claim 39, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.
46. The isolated nucleic acid molecule of claim 45, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being post-translationally modified by biotinylation, attachment of 4-phosphopanthetheine, attachment of lipoic acid or attachment of flavins.
47. The isolated nucleic acid molecule of claim 45, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being biotinylated.
48. The isolated nucleic acid molecule of claim 47, wherein said amino acid sequence that is capable of being biotinylated is all or a portion of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, all or a portion of the *Propionibacterium shermanii* transcarboxylase 1.3S subunit, or all or a portion of the *Escherichia coli* biotin carboxyl carrier protein component of acetyl-CoA carboxylase.



-87-

49. The isolated nucleic acid molecule of claim 47, wherein said amino acid sequence that is capable of being biotinylated is a portion of the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit.
50. The isolated nucleic acid molecule of claim 49, wherein said amino acid sequence that is capable of being biotinylated is the BIOTAG™.
51. The isolated nucleic acid molecule of claim 39, wherein said nucleic acid molecule is a circular molecule.
52. The isolated nucleic acid molecule of claim 39, wherein said nucleic acid molecule comprises two or more recombination sites.
53. The isolated nucleic acid molecule of claim 39, wherein said topoisomerase is a type I topoisomerase.
54. The isolated nucleic acid molecule of claim 53, wherein said type I topoisomerase is a type IB topoisomerase.
55. The isolated nucleic acid molecule of claim 54, wherein said type IB topoisomerase is selected from the group consisting of eukaryotic nuclear type I topoisomerase and a poxvirus topoisomerase.
56. The isolated nucleic acid molecule of claim 55, wherein said poxvirus topoisomerase is produced by or isolated from a virus selected from the group consisting of vaccinia virus, Shope fibroma virus, ORF virus, fowlpox virus, molluscum contagiosum virus and *Amsacta moorei* entomopoxvirus.
57. A vector comprising the isolated nucleic acid molecule of claim 39.

58. A host cell comprising the isolated nucleic acid molecule of claim 39.

59. A host cell comprising the vector of claim 57.

60. A method of producing a polynucleotide construct that encodes a fusion protein that comprises an amino acid sequence tag, said method comprising:

- (a) obtaining a first nucleic acid molecule comprising a nucleotide sequence of interest;
- (b) obtaining a second nucleic acid molecule comprising at least two topoisomerase recognition sites, at least one topoisomerase, and at least one nucleic acid sequence which encodes an amino acid sequence tag;
- (c) mixing said first nucleic acid molecule with said second nucleic acid molecule; and
- (d) incubating said mixture under conditions such that said first nucleic acid molecule is inserted into said second nucleic acid molecule between said at least two topoisomerase recognition sites, thereby producing a product polynucleotide construct;

wherein said product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleotide sequence of interest.

61. The method of claim 60, wherein said second nucleic acid molecule further comprises a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases; and

wherein said product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid sequence tag, and on the other side by (iii) the amino acid sequence encoded by said nucleotide sequence of interest.

62. The method of claim 61, wherein said amino acid sequence that is capable of being cleaved by one or more proteases is an amino acid sequence that is capable of being cleaved by enterokinase.
63. The method of claim 60, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.
64. The method of claim 63, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being post-translationally modified by biotinylation, attachment of 4-phosphopanthetheine, attachment of lipoic acid or attachment of flavins.
65. The method of claim 63, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being biotinylated.
66. The method of claim of claim 65, wherein said amino acid sequence that is capable of being biotinylated is all or a portion of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, all or a portion of the *Propionibacterium shermanii* transcarboxylase 1.3S subunit, or all or a portion of the *Escherichia coli* biotin carboxyl carrier protein component of acetyl-CoA carboxylase.
67. The method of claim of claim 65, wherein said amino acid sequence that is capable of being biotinylated is a portion of the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit.
68. The method of claim 67, wherein said amino acid sequence that is capable of being biotinylated is the BIOTAG™.

-90-

69. The method of claim 60, wherein said second nucleic acid molecule is a vector.
70. The method of claim 60, wherein said first nucleic acid molecule is a linear nucleic acid molecule.
71. The method of claim 70, wherein said first nucleic acid molecule is a blunt-end nucleic acid molecule.
72. The method of claim 60, wherein said first nucleic acid molecule is a PCR product.
73. The method of claim 60, further comprising inserting said product polynucleotide construct into a host cell.
74. The method of claim 61, further comprising inserting said product polynucleotide construct into a host cell.
75. The method of claim 60, wherein said second nucleic acid molecule comprises at least one additional nucleic acid sequence selected from the group consisting of a selectable marker, a cloning site, a restriction site, a promoter, an operator, an operon, a nucleotide sequence encoding a gene product which allows for negative selection, an origin of replication, a nucleotide sequence which encodes a repressor of at least one promoter, and a gene or partial gene.
76. The method of claim 60, wherein said topoisomerase is a type I topoisomerase.
77. The method of claim 76, wherein said type I topoisomerase is a type IB topoisomerase.

-91-

78. The method of claim 77, wherein said type IB topoisomerase is selected from the group consisting of eukaryotic nuclear type I topoisomerase and a poxvirus topoisomerase.
79. The method of claim 78, wherein said poxvirus topoisomerase is produced by or isolated from a virus selected from the group consisting of vaccinia virus, Shope fibroma virus, ORF virus, fowlpox virus, molluscum contagiosum virus and *Amsacta moorei* entomopoxvirus.
80. An isolated nucleic acid molecule comprising:
- (a) one or more recombination sites;
  - (b) one or more topoisomerase recognition sites and/or one or more topoisomerases; and
  - (c) one or more nucleic acid sequences which encode an amino acid sequence tag.
81. The isolated nucleic acid molecule of claim 80, further comprising at least one additional nucleic acid sequence selected from the group consisting of a selectable marker, a cloning site, a restriction site, a promoter, an operator, an operon, a nucleotide sequence encoding a gene product which allows for negative selection, an origin of replication, a nucleotide sequence which encodes a repressor of at least one promoter, and a gene or partial gene.
82. The isolated nucleic acid molecule of claim 80, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more recombination sites, thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest.

83. The isolated nucleic acid molecule of claim 80, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotides of the position of said one or more topoisomerases, thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid tag; and (ii) the amino acid sequence encoded by said nucleic acid sequence of interest.
84. The isolated nucleic acid molecule of claim 80, further comprising a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases.
85. The isolated nucleic acid molecule of claim 84, wherein said amino acid sequence that is capable of being cleaved by one or more proteases is an amino acid sequence that is capable of being cleaved by enterokinase.
86. The isolated nucleic acid molecule of claim 84, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more recombination sites, thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid sequence tag, and on the other side by (iii) the amino acid sequence encoded by said nucleic acid sequence of interest.
87. The isolated nucleic acid molecule of claim 84, wherein a nucleic acid sequence of interest can be inserted at or within 20 nucleotides of said one or more topoisomerase recognition sites and/or at or within 20 nucleotides of the position of said one or more topoisomerases, thereby producing a polynucleotide construct that encodes a fusion protein, said fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid

sequence tag, and on the other side by (iii) the amino acid sequence encoded by said nucleic acid sequence of interest.

88. The isolated nucleic acid molecule of claim 80, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.

89. The isolated nucleic acid molecule of claim 88, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being post-translationally modified by biotinylation, attachment of 4-phosphopanthetheine, attachment of lipoic acid or attachment of flavins.

90. The isolated nucleic acid molecule of claim 80, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being biotinylated.

91. The isolated nucleic acid molecule of claim 90, wherein said amino acid sequence that is capable of being biotinylated is all or a portion of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, all or a portion of the *Propionibacterium shermanii* transcarboxylase 1.3S subunit, or all or a portion of the *Escherichia coli* biotin carboxyl carrier protein component of acetyl-CoA carboxylase.

92. The isolated nucleic acid molecule of claim 90, wherein said amino acid sequence that is capable of being biotinylated is a portion of the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit.

93. The isolated nucleic acid molecule of claim 92, wherein said amino acid sequence that is capable of being biotinylated is the BIOTAG™.

-94-

94. The isolated nucleic acid molecule of claim 80, wherein said nucleic acid molecule is a circular molecule.
95. The isolated nucleic acid molecule of claim 80, wherein said nucleic acid molecule comprises two or more recombination sites.
96. The isolated nucleic acid molecule of claim 80, wherein said recombination sites are selected from the group consisting of: (a) *attB* sites, (b) *attP* sites, (c) *attL* sites, (d) *attR* sites, (e) *lox* sites, (f) *psi* sites, (g) *dif* sites, (h) *cer* sites, (i) *frt* sites, and mutants, variants, and derivatives of the recombination sites of (a), (b), (c), (d), (e), (f), (g), (h), or (i) which retain the ability to undergo recombination.
97. The isolated nucleic acid molecule of claim 80, wherein said topoisomerase is a type I topoisomerase.
98. The isolated nucleic acid molecule of claim 97, wherein said type I topoisomerase is a type IB topoisomerase.
99. The isolated nucleic acid molecule of claim 98, wherein said type IB topoisomerase is selected from the group consisting of eukaryotic nuclear type I topoisomerase and a poxvirus topoisomerase.
100. The isolated nucleic acid molecule of claim 99, wherein said poxvirus topoisomerase is produced by or isolated from a virus selected from the group consisting of vaccinia virus, Shope fibroma virus, ORF virus, fowlpox virus, molluscum contagiosum virus and *Amsacta moorei* entomopoxvirus.
101. A vector comprising the isolated nucleic acid molecule of claim 80.



102. A host cell comprising the isolated nucleic acid molecule of claim 80.
103. A host cell comprising the vector of claim 101.
104. A method of producing a polynucleotide construct that encodes a fusion protein that comprises an amino acid sequence tag, said method comprising:
- (a) obtaining a first nucleic acid molecule comprising a nucleotide sequence of interest;
  - (b) obtaining a second nucleic acid molecule comprising (i) at least a first topoisomerase recognition site flanked by (ii) at least a first recombination site, and (iii) at least a second topoisomerase recognition site flanked by (iv) at least a second recombination site, wherein said first and second recombination sites do not recombine with each other, and (v) at least one topoisomerase;
  - (c) obtaining a third nucleic acid molecule comprising: (i) at least a third and fourth recombination sites that do not recombine with each other; and (ii) one or more nucleic acid sequences which encode an amino acid sequence tag;
  - (d) mixing said first nucleic acid molecule with said second nucleic acid molecule;
  - (e) incubating said mixture under conditions such that said first nucleic acid molecule is inserted into said second nucleic acid molecule between said at least two topoisomerase recognition sites, thereby producing a first product polynucleotide construct;
  - (f) contacting said first product polynucleotide construct with said third nucleic acid molecule under conditions favoring recombination between said first and third and between said second and fourth recombination sites, thereby producing a second product polynucleotide construct;

wherein said second product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence tag; and (ii) the amino acid sequence encoded by said nucleotide sequence of interest.

105. The method of claim 104, wherein said third nucleic acid molecule further comprises a nucleic acid sequence that encodes an amino acid sequence that is capable of being cleaved by one or more proteases; and wherein said second product polynucleotide construct encodes a fusion protein comprising: (i) said amino acid sequence that is capable of being cleaved by one or more proteases, flanked on one side by (ii) said amino acid sequence tag, and on the other side by (iii) the amino acid sequence encoded by said nucleotide sequence of interest.

106. The method of claim 105, wherein said amino acid sequence that is capable of being cleaved by one or more proteases is an amino acid sequence that is capable of being cleaved by enterokinase.

107. The method of claim 104, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.

108. The method of claim 107, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being post-translationally modified by biotinylation, attachment of 4-phosphopantetheine, attachment of lipoic acid or attachment of flavins.

109. The method of claim 107, wherein said amino acid sequence that is capable of being post-translationally modified is an amino acid sequence that is capable of being biotinylated.

110. The method of claim of claim 109, wherein said amino acid sequence that is capable of being biotinylated is all or a portion of the *Klebsiella*

*pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit, all or a portion of the *Propionibacterium shermanii* transcarboxylase 1.3S subunit, or all or a portion of the *Escherichia coli* biotin carboxyl carrier protein component of acetyl-CoA carboxylase.

111. The method of claim 109, wherein said amino acid sequence that is capable of being biotinylated is a portion of the C-terminus of the *Klebsiella pneumoniae* oxalacetate decarboxylase  $\alpha$  subunit.

112. The method of claim 111, wherein said amino acid sequence that is capable of being biotinylated is the BIOTAG™.

113. The method of claim 104, wherein said second nucleic acid molecule is a vector.

114. The method of claim 104, wherein said third nucleic acid molecule is a vector.

115. The method of claim 104, wherein said first nucleic acid molecule is a linear nucleic acid molecule.

116. The method of claim 115, wherein said first nucleic acid molecule is a blunt-end nucleic acid molecule.

117. The method of claim 104, wherein said first nucleic acid molecule is a PCR product.

118. The method of claim 104, further comprising inserting said first product polynucleotide construct into a host cell.

119. The method of claim 104, further comprising inserting said second product polynucleotide construct into a host cell.
120. The method of claim 104, wherein said second and/or said third nucleic acid molecules comprises at least one additional nucleic acid sequence selected from the group consisting of a selectable marker, a cloning site, a restriction site, a promoter, an operator, an operon, a nucleotide sequence encoding a gene product which allows for negative selection, an origin of replication, a nucleotide sequence which encodes a repressor of at least one promoter, and a gene or partial gene.
121. The method of claim 104, wherein said first, second, third and fourth recombination sites are selected from the group consisting of: (a) *attB* sites, (b) *attP* sites, (c) *attL* sites, (d) *attR* sites, (e) *lox* sites, (f) *psi* sites, (g) *dif* sites, (h) *cer* sites, (i) *frt* sites, and mutants, variants, and derivatives of the recombination sites of (a), (b), (c), (d), (e), (f), (g), (h), or (i) which retain the ability to undergo recombination.
122. The method of claim 104, wherein said topoisomerase is a type I topoisomerase.
123. The method of claim 122, wherein said type I topoisomerase is a type IB topoisomerase.
124. The method of claim 123, wherein said type IB topoisomerase is selected from the group consisting of eukaryotic nuclear type I topoisomerase and a poxvirus topoisomerase.
125. The method of claim 124, wherein said poxvirus topoisomerase is produced by or isolated from a virus selected from the group consisting of vaccinia virus, Shope fibroma virus, ORF virus, fowlpox virus, molluscum contagiosum virus and *Amsacta moorei* entomopoxvirus.

126. The method of claim 104, wherein said first product polynucleotide construct and said third nucleic acid molecule are combined in the presence of at least one recombination protein.
127. The method of claim 126, wherein said recombination protein is selected from the group consisting of: (a) Cre, (b) Int, (c) IHF, (d) Xis, (e) Fis, (f) Hin, (g) Gin, (h) Cin, (i) Tn3 resolvase, (j) TndX, (k) XerC, and (l) XerD.
128. The method of claim 126, wherein said recombination protein is Cre.
129. A vector selected from the group consisting of pET104-DEST, pET104/GW/*lacZ*, pET104/D-TOPO, pET104/D/*lacZ*, pcDNA6/Biotag<sup>TM</sup>-DEST, pcDNA6/Biotag<sup>TM</sup>-GW/*lacZ*, pcDNA6/Biotag<sup>TM</sup>/D-TOPO, pcDNA6/Biotag<sup>TM</sup>/*lacZ*, pMT/Biotag<sup>TM</sup>-DEST, and pMT/Biotag<sup>TM</sup>/GW-*lacZ*.
130. A kit comprising the isolated nucleic acid molecule of claim 1.
131. The kit of claim 130, further comprising one or more components selected from the group consisting of one or more topoisomerases, one or more recombination proteins, one or more vectors, one or more polypeptides having polymerase activity, one or more host cells, and one or more support matrices complexed with avidin or an avidin analog.
132. A kit comprising the isolated nucleic acid molecule of claim 39.
133. The kit of claim 132, further comprising one or more components selected from the group consisting of one or more topoisomerases, one or more recombination proteins, one or more vectors, one or more

-100-

polypeptides having polymerase activity, one or more host cells, and one or more support matrices complexed with avidin or an avidin analog.

134. A kit comprising the isolated nucleic acid molecule of claim 80.
135. The kit of claim 134, further comprising one or more components selected from the group consisting of one or more topoisomerases, one or more recombination proteins, one or more vectors, one or more polypeptides having polymerase activity, one or more host cells, and one or more support matrices complexed with avidin or an avidin analog.
136. A host cell comprising a polynucleotide construct that encodes a fusion protein capable of being post-translationally modified, said polynucleotide construct produced according to the method of claim 19.
137. A host cell comprising a polynucleotide construct that encodes a fusion protein capable of being post-translationally modified, said polynucleotide construct produced according to the method of claim 60.
138. A host cell comprising a polynucleotide construct that encodes a fusion protein capable of being post-translationally modified, said polynucleotide construct produced according to the method of claim 104.
139. A method of producing a fusion protein that comprises an amino acid sequence tag, said method comprising:
- (a) obtaining the host cell of claim 136; and
  - (b) culturing said host cell under conditions wherein said fusion protein is produced by said host cell.
140. The method of claim 139, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.

141. The method of claim 140, further comprising culturing said host cell under conditions wherein said fusion protein is post-translationally modified in said host cell.
142. The method of claim 140, further comprising culturing said host cell under conditions wherein said fusion protein is biotinylated in said host cell.
143. The method of claim 139, further comprising:
- (a) treating said host cell such that said fusion protein is released from said host cell; and
  - (b) contacting said fusion protein with a detecting composition comprising a molecule that is capable of interacting with said amino acid sequence tag or with a molecular entity that is attached to said amino acid sequence tag.
144. The method of claim 143, wherein said fusion protein is a biotinylated fusion protein, and said detecting composition comprises avidin or an avidin analogue.
145. A method of producing a fusion protein that comprises an amino acid sequence tag, said method comprising:
- (a) obtaining the host cell of claim 137; and
  - (b) culturing said host cell under conditions wherein said fusion protein is produced by said host cell.
146. The method of claim 145, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.
147. The method of claim 146, further comprising culturing said host cell under conditions wherein said fusion protein is post-translationally modified in said host cell.

148. The method of claim 146, further comprising culturing said host cell under conditions wherein said fusion protein is biotinylated in said host cell.
149. The method of claim 145, further comprising:
- (a) treating said host cell such that said fusion protein is released from said host cell; and
  - (b) contacting said fusion protein with a detecting composition comprising a molecule that is capable of interacting with said amino acid sequence tag or with a molecular entity that is attached to said amino acid sequence tag.
150. The method of claim 149, wherein said fusion protein is a biotinylated fusion protein, and said detecting composition comprises avidin or an avidin analogue.
151. A method of producing a fusion protein that comprises an amino acid sequence tag, said method comprising:
- (a) obtaining the host cell of claim 138; and
  - (b) culturing said host cell under conditions wherein said fusion protein is produced by said host cell.
152. The method of claim 151, wherein said amino acid sequence tag is an amino acid sequence that is capable of being post-translationally modified.
153. The method of claim 152, further comprising culturing said host cell under conditions wherein said fusion protein is post-translationally modified in said host cell.




-103-

154. The method of claim 152, further comprising culturing said host cell under conditions wherein said fusion protein is biotinylated in said host cell.

155. The method of claim 151, further comprising:

- (a) treating said host cell such that said fusion protein is released from said host cell; and
- (b) contacting said fusion protein with a detecting composition comprising a molecule that is capable of interacting with said amino acid sequence tag or with a molecular entity that is attached to said amino acid sequence tag.

156. The method of claim 155, wherein said post-translationally modified fusion protein is a biotinylated fusion protein, and said detecting composition comprises avidin or an avidin analogue.



CAAGGAGATGGCGCCCAACAGTCCCCCGGCCACGGGGCTGCCACCATACCCACGCCGAAACAAGCGCTC  
ATGAGCCCGAAGTGGCGAGCCCGATCTTCCCCATCGGTGATGTCGGCGATATAGGCGCCAGCAACCGCAC  
CTGTGGCGCCGGTGATGCCGGCCACGATGCGTCCGGCGTAGAGGATCGAGATCTCGATCCCGCGAAATTA  
ATACGACTCACTATAGGGGAATTGTGAGCGGATAACAATTCCCCCTCTAGAAATAATTTTGTTTAACTTTA  
AGAAGGAGATATACATATGGGCGCCGGCACCCCGGTGACCGCCCCGCTGGCGGGCACTATCTGGAAGGTG  
CTGGCCAGCGAAGGCCAGACGGTGCCCGCAGGCGAGGTGCTGCTGATTCTGGAAGCCATGAAGATGGAAA  
CCGAAATCCGCGCCGCGCAGGCCGGGACCGTGC CGGTATCGCGGTGAAAGCCGGCGACGCGGTGGCGGT  
CGGCGACACCCTGATGACCCTGGCGGGCTCTGGATCCGATCTGTACGACGATGACGATAAGGGAATTATC  
ACAAGTTTGTACAAAAAGCTGAACGAGAAAACGTAAAATGATATAAAATATCAATATATTAAATTAGATTT  
TGCATAAAAAACAGACTACATAATACTGTAAAAACACAACATATCCAGTCACTATGGCGGCCGATTAGGC  
ACCCAGGCTTTACACTTTATGCTTCCGGCTCGTATAATGTGTGGATTTTGAGTTAGGATCCGGCGAGAT  
TTTCAGGAGCTAAGGAAGCTAAAATGGAGAAAAAAATCACTGGATATACCACCGTTGATATATCCCAATG  
GCATCGTAAAGAACATTTTGAGGCATTTTCAGTCAGTTGCTCAATGTACCTATAACCAGACCGTTTCAGCTG  
GATATTACGGCCTTTTTAAAGACCGTAAAGAAAAATAAGCACAAGTTTTATCCGGCCTTTATTACATTC  
TTGCCCGCCTGATGAATGCTCATCCGGAATTCGTATGGCAATGAAAGACGGTGAGCTGGTGATATGGGA  
TAGTGTTACCCCTTGTTACACCGTTTTCCATGAGCAAACGTAAACGTTTTTCATCGCTCTGGAGTGAATAC  
CACGACGATTTCCGGCAGTTTCTACACATATATTCGCAAGATGTGGCGTGTTACGGTGAAAACCTGGCCT  
ATTTCCCTAAAGGGTTTATTGAGAATATGTTTTTCGTCTCAGCCAATCCCTGGGTGAGTTTACCAGTTT  
TGATTTAAACGTGGCCAATATGGACAACCTTCTTCGCCCCCGTTTTTACCATGGGCAATATTATACGAA  
GGCGACAAGGTGCTGATGCCGCTGGCGATTTCAGGTTTCATCATGCCGTCTGTGATGGCTTCCATGTCGGCA  
GAATGCTTAATGAATTACAACAGTACTGCGATGAGTGGCAGGGCGGGGCGTAAACCGGTGGATCCGGCTT  
ACTAAAAGCCAGATAACAGTATGCGTATTTGCGCGCACCGGTGCTAGCGTATACCCGAAGTATGTCAAAA  
AGAGGTGTGCTATGAAGCAGCGTATTACAGTGACAGTTGACAGCGACAGCTATCAGTTGCTCAAGGCATA  
TATGATGTCAATATCTCCGGTCTGGTAAGCACAACCATGCAGAATGAAGCCCGTCTGTGCGTGCCGAAC  
GCTGGAAAGCGGAAAATCAGGAAGGGATGGCTGAGGTGCGCCCGTTTTATTGAAATGAACGGCTCTTTTGC  
TGACGAGAACAGGGACTGGTGAAATGCAGTTTAAAGTTTACACCTATAAAAGAGAGAGCCGTTATCGTCT  
GTTTGTGGATGTACAGAGTGATATTATTGACACGCCCCGGGCGACGGATGGTGATCCCCCTGGCTAGGCA  
CGTCTGCTGTCAGATAAAGTCTCCCGTGAACCTTACC CGGTGGTGATATCGGGGATGAAAGCTGGCGCA  
TGATGACCACCGATATGGCCAGTGTGCCGGTCTCCGTTATCGGGGAAGAAGTGGCTGATCTCAGCCACCG  
CGAAAATGACATCAAAAACGCCATTAACCTGATGTTCTGGGGAATATAAATGTCAGGCTCCGTTATACAC  
AGCCAGTCTGCAGGTCGACCATAGTGAAGTGGATATGTTGTGTTTTACAGTATTATGTAGTCTGTTTTTTA  
TGCAAAATCTAATTTAATATATTGATATTTATATCATTTTACGTTTCTCGTTCAGCTTTCTTGTAACAAG  
TGGTGATAATTAATTAAGATAGCTCAGATCCGGCTGCTAACAAAGCCCCGAAAGGAAGCTGAGTTGGCTGC  
TGCCACCGCTGAGCAATAACTAGCATAACCCCTTGGGGCCTCTAAACGGGTCTTGAGGGGTTTTTTTGCTG  
AAAGGAGGAAGTATATCCGGATATCCCGCAAGAGGCCCGGCAGTACCGGCATAACCAAGCCTATGCCTAC  
AGCATCCAGGGTGACGGTGCCGAGGATGACGATGAGCGCATTTGTTAGATTTTCATACAGGTGCCTGACTG  
CGTTAGCAATTTAACTGTGATAAACTACCGCATTAAGCTAGCTTATCGATGATAAGCTGTCAAACATGA  
GAATTAATTTCTTGAAGACGAAAGGGCCTCGTGATACGCCATTTTTTATAGGTTAATGTCATGATAATAAT  
GGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGTTTATTTTTCTAA  
ATACATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATTTGAAAAAGGA  
AGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCTTTTTTGCGGCATTTTGCCCTCCTGTTTT  
TGCTCAGCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATC  
GAAGTGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTTCGCCCCGAAGAACGTTTTTCCAATGATGAGCA  
CTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTGTGACGCCGGGCAAGAGCAACTCGGTGCGCG  
CATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATG  
ACAGTAAGAGAATTATGCAGTGTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGACAA  
CGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCAACAACATGGGGGATCATGTAAGTTCGCCCTTGATCG  
TTGGGAACCGGAGCTGAATGAAGCCATACCAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCA  
ACAACGTTGCGCAAACTATTAAGTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAAGACTGGA  
TGGAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAA


Fig. 2A


ATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGT  
ATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAG  
GTGCCTCACTGATTAAGCATTGGTAAGTGTGACACCAAGTTTACTCATATATACTTTAGATTGATTTAA  
ACTTCATTTTTTAATTTAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAA  
CGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTT  
TTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAACCACCGCTACCAGCGGTGGTTTGTGTGCCGGATCA  
AGAGCTACCAACTCTTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCGCAGATACCAAATACTGTCTTCTA  
GTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCC  
TGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTCTGTCTTACCGGGTTGGACTCAAGACGATAGTTACC  
GGATAAGGCGCAGCGGTGGGGTGAACGGGGGGTTTCGTGCACACAGCCAGCTTGGAGCGAACGACCTAC  
ACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGGAGAAAGCGGACA  
GGTATCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCTTGGTA  
TCTTTATAGTCCGTGTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGG  
CGGAGCCTATGAAAAACGCCAGCAACGCGGCCTTTTTACGGTTCTTGGCCTTTTGTGCTGGCCTTTTGTCTC  
ACATGTTCTTCTGCGTTATCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATAC  
CGCTCGCCGACGCCAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCCTGATGCGG  
TATTTCTCCTTACGCATCTGTGCGGTATTTACACCGCATATATGGTGCCTCTCAGTACAATCTGCTC  
TGATGCCGCATAGTTAAGCCAGTATACACTCCGCTACGTGACTGGGTCTAGGCTGCGCCCCGAT  
ACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGCTGCTCCCGGCATCCGCTTACAGCAAGCTGT  
GACCGTCTCCGGGAGCTGCATGTGTGAGAGTTTTACCGTCTATCACGAAACGCGCAGGACAGCTGCGG  
TAAAGCTCATCAGCGTGGTCTGAAGCGATTACAGATGTCTGCCTGTTTCATCCGCGTCCAGCTCGTTGA  
GTTTCTCCAGAAGCGTTAATGTCTGGCTTCTGATAAAGCGGGCCATGTTAAGGGCGGTTTTTCTGTTT  
GGTCACTGATGCCCTCCGTGTAAGGGGGATTTCTGTTTCATGGGGGTAATGATACCGATGAAACGAGAGAGG  
ATGCTCACGATACGGGTACTGATGATGAACATGCCCGGTTACTGGAACGTTGTGAGGGTAAACAAGTGG  
CGGTATGGATGCGGCGGGACCAGAGAAAAATCACTCAGGGTCAATGCCAGCGCTTCGTTAATACAGATGT  
AGGTGTTCCACAGGGTAGCCAGCAGCATCCTGCGATGCAGATCCGGAACATAATGGTGCAGGGCGCTGAC  
TTCCGCGTTTCCAGACTTTACGAAACACGGAACCGAAGACCATTTCATGTTGTTGCTCAGGTGCGAGACG  
TTTTGCAGCAGCAGTCCGTTTACGTTCTCGCTCGCGTATCGGTGATTTCATTCTGCTAACCCAGTAAGGCAACC  
CCGCCAGCCTAGCCGGGTCTCAACGACAGGAGCAGCATATGCGCACCCGTTGGCCAGGACCCAACGCTG  
CCCGAGATGCGCCGCGTGGCGCTGCTGGAGATGGCGGACGCGATGGATATGTTCTGCCAAGGGTTGGTTT  
GCGCATTCACAGTTCTCCGCAAGAATTGATTGGCTCCAATTCTTGGAGTGGTGAATCCGTTAGCGAGGTG  
CCGCCGGCTTCCATTAGGTGAGGTGGCCCGGCTCCATGCACCGCGACGCAACGCGGGGAGGCAGACAA  
GGTATAGGGCGCGCGCTACAATCCATGCCAACCCGTTCCATGTGCTCGCCGAGGCGGCATAAATCGCCGT  
GACGATCAGCGGTCCAGTGATCGAAGTTAGGCTGGTAAGAGCCGCGAGCGATCCTTGAAGCTGTCCCTGA  
TGGTCTCATCTACCTGCCCTGGACAGCATGGCTGCAACGCGGGCATCCCGATGCCGCCGGAAGCGAGAA  
GAATCATAATGGGGAAGGCCATCCAGCCTCGCGTCCGGAACGCCAGCAAGACGTAGCCAGCGCGTCCGGC  
CGCCATGCCGGCGATAATGGCCTGCTTCTCGCCGAAACGTTTGGTGGCGGGACAGTGACGAAGGCTTGA  
GCGAGGGCGTGCAAGATTCCGAATACCGCAAGCGACAGGCCGATCATCGTCCGCTCCAGCGAAAGCGGT  
CCTCGCCGAAAATGACCCAGAGCGTCCCGGCACCTGTCTACGAGTTGCATGATAAAGAAGACAGTCAT  
AAGTGCGGCGACGATAGTCATGCCCCGCGCCACCGGAAGGAGCTGACTGGGTGGAAGGCTCTCAAGGGC  
ATCGGTGCGAGATCCCGGTGCCATAATGAGTGAGCTAATTACATTAATTGCGTTGCGCTCACTGCCCGCTT  
TCCAGTCCGGGAAACCTGTGCTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGC  
TATTGGGCGCCAGGGTGGTTTTCTTTTACCAGTGAGACGGGCAACAGCTGATTGCCCTTACCAGCCTG  
GCCCTGAGAGAGTTGCAGCAAGCGGTCCACGCTGGTTTGGCCCGAGCAGGCGAAAATCCTGTTTGATGGTG  
GTTAACGGCGGGATATAACATGAGCTGTCTTCGGTATCGTCGTATCCCACTACCGAGATATCCGCACCAA  
CGCGCAGCCCGGACTCGGTAATGGCGCGCATTGCGCCAGCGCCATCTGATCGTTGGCAACCAGCATCGC  
AGTGGGAACGATGCCCTCATTCAGCATTTGCATGGTTTGTGAAAACCGGACATGGCACTCCAGTCCGCT  
TCCCGTTCCGCTATCGGCTGAATTTGATTGCGAGTGAGATATTTATGCCAGCCAGCCAGACGCGAGCGG  
CCGAGACAGAACTAATGGGCCCCGTAACAGCGCGATTGCTGGTGACCCAATGCCAGCAGATGCTCCAC  
GCCAGTCCGCTACCGTCTTCATGGGAGAAAAATAACTGTTGATGGGTGTCTGGTCAGAGACATCAAGA

Fig. 2B

AATAACGCCGGAACATTAGTGCAGGCAGCTTCCACAGCAATGGCATCCTGGTCATCCAGCGGATAGTTAA  
TGATCAGCCCACTGACGCGTTGCGCGAGAAGATTGTGCACCGCCGCTTTACAGGCTTCGACGCCGCTTCG  
TTCTACCATCGACACCACCACGCTGGCAGCCAGTTGATCGGCGCGAGATTTAATCGCCGCGACAATTTGC  
GACGGCGCGTGCAGGGCCAGACTGGAGGTGGCAACGCCAATCAGCAACGACTGTTTGCCCGCCAGTTGTT  
GTGCCACGCGGTTGGGAATGTAATTCAGCTCCGCCATCGCCGCTTCCACTTTTTCCCGCGTTTTTCGCAGA  
AACGTGGCTGGCCTGGTTTACCACGCGGGAAACGGTCTGATAAGAGACACCGGCATACTCTGCGACATCG  
TATAACGTTACTGGTTTTACATTACCAACCCTGAATTGACTCTCTTCCGGGCGCTATCATGCCATACCGC  
GAAAGGTTTTGCGCCATTCGATGGTGTCCGGGATCTCGACGCTCTCCCTTATGCGACTCCTGCATTAGGA  
AGCAGCCCAGTAGTAGGTTGAGGCCGTTGAGCACCGCCGCCGCAAGGAATGGTGCATG

Fig. 2C





CAAGGAGATGGCGCCCAACAGTCCCCCGGCCACGGGGCCTGCCACCATACCCACGCCGAAACAAGCGCTC  
ATGAGCCCGAAGTGGCGAGCCCGATCTTCCCCATCGGTGATGTCGGCGATATAGGCGCCAGCAACCGCAC  
CTGTGGCGCCGGTATGCCGGCCACGATGCGTCCGGCGTAGAGGATCGAGATCTCGATCCCGCGAAATTA  
ATACGACTCACTATAGGGGAATTGTGAGCGGATAACAATTCCCCCTCTAGAAATAATTTTGTTTAACTTTA  
AGAAGGAGATATACATATGGGCGCCGGCACCCCGGTGACCGCCCCGCTGGCGGGCACTATCTGGAAGGTG  
CTGGCCAGCGAAGGCCAGACGGTGGCCGCAGGCGAGGTGCTGCTGATTCTGGAAGCCATGAAGATGGA  
CCGAAATCCGCGCCGCGCAGGCCGGGACCGTGCGCGGTATCGCGGTGAAAGCCGGCGACGCGGTGGCGGT  
CGGCGACACCCTGATGACCCTGGCGGGCTCTGGATCCGATCTGTACGACGATGACGATAAGGGAATTGAT  
CCCTTCACCAAGGGCGAGCTCAGATCCGGCTGCTAACAAAGCCCGAAAGGAAGCTGAGTTGGCTGCTGCC  
ACCGCTGAGCAATAACTAGCATAACCCCTTGGGGCCTCTAAACGGGTCTTGAGGGGTTTTTTGCTGAAAG  
GAGGAACATATATCCGGATATCCCGCAAGAGGCCCGGCAGTACCGGCATAACCAAGCCTATGCCTACAGCA  
TCCAGGGTGACGGTGCCGAGGATGACGATGAGCGCATTGTTAGATTTCATACACGGTGCCTGACTGCGTT  
AGCAATTTAACTGTGATAAACTACCGCATTAAAGCTAGCTTATCGATGATAAGCTGTCAAACATGAGAAT  
TAATTCCTGAAGACGAAAGGGCCTCGTGATACGCCTATTTTTATAGGTTAATGTCTATGATAATAATGGTT  
TCTTAGACGTCAGGTGGCCTTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGCTTTATTTCTAAATAC  
ATTCAAATATGTATCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAG  
TATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCTTTTTTTCGCGCATTTTGCCTTCCTGTTTTTGTCT  
CACCCAGAAACGCTGGTGAAAGTAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTTACATCGAAC  
TGGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTT  
TAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTGTGACGCCGGGCAAGAGCAACTCGGTGCGCGCATA  
CACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAG  
TAAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAAACTGCGGCCAACTTACTTCTGACAACGAT  
CGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAACCTGCCTTGATCGTTGG  
GAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAA  
CGTTGCGCAAACTATTAACCTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGATGGA  
GGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGCTGATAAATCT  
GGAGCCGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCG  
TAGTTATCTACACGACGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGC  
CTCACTGATTAAGCATTGGTAAGTGTGACACCAAGTTTACTCATATATACTTTAGATTGATTTAAACTT  
CATTTTTAATTTAAAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCCTTAACGCTG  
AGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCT  
GCGCGTAATCTGCTGCTTGCAAACAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCAAGAG  
CTACCAACTCTTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCGCAGATACCAAATACTGTCCTTCTAGTGT  
AGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCCTACATACCTCGCTCTGCTAATCCTGTT  
ACCAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGTTGGACTCAAGACGATAGTTACCGGAT  
AAGGCGCAGCGGTGCGGCTGAACGGGGGGTTCGTGCACACAGCCAGCTTGGAGCGAACGACCTACACCG  
AACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTA  
TCCGGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCTTGGTATCTT  
TATAGTCCTGTGGGTTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGA  
GCCTATGGAACAAACGCCAGCAACGCGGCTTTTTACGGTCTCTGGCCTTTTGCTGGCCTTTTGCTCACAT  
GTTCTTTTCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCGCT  
CGCCGAGCCGAACGACGCGCAGCGAGTCAGTGAGCGAGGAAGCGGAAGAGCGCCTGATGCGGTATT  
TTCTCCTTACGCATCTGTGCGGTATTTACACCCGCATATATGGTGCACTCTCAGTACAATCTGCTCTGAT  
GCCGCATAGTTAAGCCAGTATACACTCCGCTATCGCTACGTGACTGGGTGCTGCTGCGCCCCGACACCC  
GCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCCGGCATCCGCTTACAGACAAGCTGTGACC  
GTCTCCGGGAGCTGCATGTGTGAGAGTTTTACCGTCATCACCGAAACGCGGAGGACGCTGCGGTAAA  
GCTCATCAGCGTGGTGTGAAGCGATTACAGATGTCTGCCTGTTTATCCGCTCCAGCTCGTTGAGTTT  
CTCCAGAAGCGTTAATGTCTGGCTTCTGATAAAGCGGGCCATGTTAAGGGCGGTTTTTCTGTTTGGTC  
ACTGATGCCTCCGTGTAAGGGGGATTCTGTTTATGGGGTAATGATACCGATGAAACGAGAGAGGATGC  
TCACGATACGGGTTACTGATGATGAACATGCCCGGTTACTGGAACGTTGTGAGGGTAAACAACCTGGCGGT

Fig. 5A



ATGGATGCGGCGGGACCAGAGAAAAATCACTCAGGGTCAATGCCAGCGCTTCGTTAATACAGATGTAGGT  
GTTCCACAGGGTAGCCAGCAGCATCCTGCGATGCAGATCCGGAACATAATGGTGCAGGGCGCTGACTTCC  
GCGTTTCCAGACTTTACGAAACACGGAAACCGAAGACCATTTCATGTTGTTGCTCAGGTTCGAGACGTTTT  
GCAGCAGCAGTCGCTTACGTTGCTCGCTCGCGTATCGGTGATTTCATTCTGCTAACAGTAAGGCAACCCCGC  
CAGCCTAGCCGGGTCTCAACGACAGGAGCAGCATCATGCCGACCCGTGGCCAGGACCCAACGCTGCCCG  
AGATGCGCCGCGTGGGGCTGCTGGAGATGGCGGACGCGATGGATATGTTCTGCCAAGGGTTGGTTTGCGC  
ATTCACAGTTCTCCGCAAGAATTGATTGGCTCCAATTCTTGGAGTGGTGAATCCGTTAGCGAGGTGCCGC  
CGGCTTCCATTAGGTTCGAGGTGGCCCGGCTCCATGCACCGCGACGCAACGCGGGGAGGCAGACAAGGTA  
TAGGGCGGCGCCTACAATCCATGCCAACCCTTCCATGTGCTCGCCGAGGCGGCATAAATCGCCGTGACG  
ATCAGCGGTCCAGTGATCGAAGTTAGGCTGGTAAGAGCCGCGAGCGATCCTTGAAGCTGTCCCTGATGGT  
CGTCATCTACCTGCCTGGACAGCATGGCCTGCAACGCGGGCATCCCGATGCCGCCGGAAGCGAGAAGAAT  
CATAATGGGGAAGGCCATCCAGCCTCGCGTCGCGAACGCCAGCAAGACGTAGCCAGCGCGTCCGGCCGCC  
ATGCCGGCGATAATGGCCTGCTTCTCGCCGAAACGTTTGGTGGCGGGACCAGTGACGAAGGCTTGAGCGA  
GGGCGTGCAAGATTCCGAATACCGCAAGCGACAGGCCGATCATCGTCGCGCTCCAGCGAAAGCGGTCTCTC  
GCCGAAATGACCCAGAGCGCTGCCGGCACCTGTCTACGAGTTGTCATGATAAAGAAGACAGTCATAAGT  
GCGGCGACGATAGTCATGCCCCGCGCCACCGGAAGGAGCTGACTGGGTGAAGGCTCTCAAGGGCATCG  
GTCGAGATCCCGGTGCTTAATGAGTGAGCTAACTTACATTAATTGCGTTGCGCTCACTGCCCGCTTTCCA  
GTCGGGAAACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCCTATT  
GGGCGCCAGGGTGGTTTTTCTTTTACCAGTGAGACGGGCAACAGCTGATTGCCCTTACCAGCCTGGCCC  
TGAGAGAGTTGCAGCAAGCGGTCCACGCTGGTTTGGCCCCAGCAGGCGAAAATCCTGTTTGATGGTGGTTA  
ACGGCGGGATATAACATGAGCTGTCTTCGGTATCGTCGTATCCCACTACCGAGATATCCGCACCAACGCG  
CAGCCCGGACTCGGTAATGGCGCGCATTGCGCCAGCGCCATCTGATCGTTGGCAACCAGCATCGCAGTG  
GGAACGATGCCCTCATTAGCATTTGCATGGTTTGTGAAAACCGGACATGGCACTCCAGTCGCCTTCCC  
GTTCCGCTATCGGCTGAATTTGATTGCGAGTGAGATATTTATGCCAGCCAGCCAGACGCAGACGCGCCGA  
GACAGAACTTAATGGGCCCCTAACAGCGCGATTTGCTGGTGACCAATGCGACCAGATGCTCCACGCCC  
AGTCGCGTACCGTCTTCATGGGAGAAAATAATACTGTTGATGGGTGTCTGGTCAGAGACATCAAGAAATA  
ACGCCGGAACATTAGTGAGGAGCTTCCACAGCAATGGCATCCTGGTCATCCAGCGGATAGTTAATGAT  
CAGCCCACTGACGCGTTGCGCGAGAAGATTGTGCACCGCCGCTTTACAGGCTTCGACGCGCGTTCGTTCT  
ACCATCGACACCACCGCTGGCACCCAGTTGATCGGCGCGAGATTTAATCGCCGCGACAATTTGCGACG  
GCGCGTGCCAGGGCCAGACTGGAGGTGGCAACGCCAATCAGCAACGACTGTTTGCCCGCCAGTTGTTGTGC  
CACGCGTTGGGAATGTAATTCAGCTCCGCCATCGCCGCTTCCACTTTTCCCGCGTTTTCGCAGAAACG  
TGGCTGGCCTGGTTTACCACGCGGGAACGGTCTGATAAGAGACACCGGCATACTCTGCGACATCGTATA  
ACGTTACTGGTTTACATTACCCACCCTGAATTGACTCTCTTCCGGGCGCTATCATGCCATACCGCGAAA  
GGTTTTGCGCCATTGATGGTGTCCGGGATCTCGACGCTCTCCCTTATGCGACTCCTGCATTAGGAAGCA  
GCCAGTAGTAGGTTGAGGCCGTTGAGCACCGCCGCGCGCAAGGAATGGTGCATG

Fig. 5B




Fig. 6




Fig.7

GACGGATCGGGAGATCTCCCGATCCCCATGCGTCTCAGTACAATCTGCTCTGATGCCGCATAGTT  
AAGCCAGTATCTGCTCCCTGCTTGTGTGTTGGAGGTCGCTGAGTAGTGCGCGAGCAAAATTTAAGCTACA  
ACAAGGCAAGGCTTGACCGACAATTGCATGAAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCG  
ATGTACGGGCCAGATATACGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTC  
ATTAGTTCATAGCCCATATATGGAGTTCGCGTTACATAACTTACGGTAAATGGCCCCGCTGGCTGACCG  
CCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCC  
ATTGACGTCAATGGGTGGACTATTTACGGTAAACTGCCACTTGGCAGTACATCAAGTGATCATATGCC  
AAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCCGCTGGCATTATGCCCAGTACATGACCTTA  
TGGGACTTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGC  
AGTACATCAATGGGCGTGGATAGCGGTTTGACTCACGGGGATTTCGAAGTCTCCACCCCATTGACGTCAA  
TGGGAGTTTTGTTTTGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACCTCGCCCCATTGACG  
CAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCTCTGGCTAAGTACGAAACCCA  
CTGCTTACTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCCAAGCTGGCTAGCGTTTTAACTT  
AAGCTTACCATGGGCGCGGCCACCCCGGTGACCGCCCCGCTGGCGGGCACTATCTGGAAGGTGCTGGCCA  
GCGAAGGCCAGACGGTGGCCGCGAGGCGAGGTGCTGCTGATTCTGGAAGCCATGAAGATGGAAACCGAAAT  
CCGCGCCGCGCAGGCCGGGACCGTGGCGGCTATCGCGGTGAAAGCCGCGACGCGGTGGCGGTGGCGGAC  
ACCCTGATGACCCTGGCGGGCTCTGGATCCGATCTGTACGACGATGACGATAAGGTACATCAAACAAGTT  
TGTACAAAAAGCTGAACGAGAAACGTAAAATGATATAAATATCAATATATTAAATTAGATTTTGCATAA  
AAAACAGACTACATAATACTGTAAACACAACATATCCAGTCACTATGGCGGCCGATTAGGCACCCAG  
GCTTTACACTTTATGCTTCCGGCTCGTATAATGTGTGGATTTTGTAGTTAGGATCCGGCGAGATTTTCAGG  
AGCTAAGGAAGCTAAAATGGAGAAAAAATCACTGGATATACCACCGTTGATATATCCCAATGGCATCGT  
AAAGAACATTTTGAGGCATTTTCACTCAGTTGCTCAATGTACCTATAACCAGACCGTTTCACTGGATATTA  
CGGCCTTTTTAAAGACCGTAAAGAAAAATAAGCACAAGTTTTATCCGGCCTTTATTACATTCTTGCCCG  
CCTGATGAATGCTCATCCGGAATTCCGTATGGCAATGAAAGACGGTGAGCTGGTGATATGGGATAGTGTT  
CACCTTGTACACCGTTTTCCATGAGCAAACTGAAACGTTTTTCATCGCTCTGGAGTGAATACCACGACG  
ATTTCCGGCAGTTTTCTACACATATATTGCAAGATGTGGCGTGTACGGTGAAAACCTGGCCTATTTCCC  
TAAAGGGTTTTATTGAGAATATGTTTTTCTGCTCAGCCAATCCCTGGGTGAGTTTACCAGTTTTGATTTA  
AACGTGGCCAATATGGACAACCTTCTCGCCCCCGTTTTTACCATGGGCAATATTATACGGAAGGCGACA  
AGGTGCTGATGCCGCTGGCGATTACAGTTTATCATGCGCTCTGTGATGGCTTCCATGTGCGCAGAATGCT  
TAATGAATTACAACAGTACTGCGATGAGTGGCAGGGCGGGCGTAAACGCGTGGATCCGGCTTACTAAAA  
GCCAGATAACAGTATGCGTATTTGCGCGCTCGCGAACCGGTGTATACCCGAAGTATGTCAAAAAGAGGTG  
TGCTATGAAGCAGCGTATTACAGTGACAGTTGACAGCGACAGCTATCAGTTGCTCAAGGCATATATGATG  
TCAATATCTCCGCTCTGGTAAGCACAACCATGCAGAATGAAGCCCGTCTGCTGCGTGCCGAACGCTGGAA  
AGCGGAAAATCAGGAAGGGATGGCTGAGGTGCGCCGTTTTATTGAAATGAACGGCTCTTTTGCTGACGAG  
AACAGGGACTGGTGAAATGCAGTTTAAAGTTTACACCTATAAAGAGAGAGCCGTTATCGTCTGTTTGTG  
GATGTACAGAGTGATATTATTGACACGCCCCGGCGACGGATGGTGATCCCCCTGGCCAGTGCACGTCTGC  
TGTCAGATAAAGTCTCCCGTGAACTTTACCCGGTGGTGATATCGGGGATGAAAGCTGGCGCATGATGAC  
CACCGATATGGCCAGTGTGCCGCTTCCGTTATCGGGGAAGAAGTGGCTGATCTCAGCCACCGCGAAAAT  
GACATCAAAAACGCCATTAACTGATGTTCTGGGGAATATAAATGTCAGGCTCCGTTATACACAGCCAGT  
CTGCAGGTGACCATAGTGACTGGATATGTTGTGTTTACAGTATTATGTAGTCTGTTTTTATGCAAAA  
TCTAATTTAATATATTGATATTTATATCATTTTTACGTTTCTCGTTTCAAGCTTTCTGTACAAAGTGGTGAT  
AATTAATTAAGATCTAGAGGGCCCGTTAAACCCGCTGATCAGCCTCGACTGTGCCCTTCTAGTTGCCAGC  
CATCTGTTGTTTGGCCCTCCCCGCTGCCTTCCCTGACCCCTGGAAGGTGCCACTCCCACTGTCTTTCCTA  
ATAAATGAGGAAATTGCATCGCATTTGTCTGAGTAGGTGTCATTCTATTCTGGGGGGTGGGGTGGGGCAG  
GACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGGATGCGGTGGGCTCTATGGCTTCTG  
AGGCGGAAAGAACCAGCTGGGGCTCTAGGGGGTATCCCCACGCGCCCTGTAGCGGCGCATTAAGCGCGGC  
GGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCGCTCCTTTTCGCTTTC  
TTCCCTTCTTTCTCGCCACGTTTCGCGGCTTTCCCGCTCAAGCTCTAAATCGGGGCATCCCTTTAGGGT  
TCCGATTTAGTGCTTTACGGCACCTCGACCCAAAAAATCTGATTAGGGTGATGGTTCACGTAGTGGGCC  
ATCGCCCTGATAGACGGTTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCT

Fig. 8A

CAAACCTGGAACAACACTCAACCCTATCTCGGTCTATTCTTTTGATTTATAAGGGATTTTGGGGATTTCCG  
CCTATTGGTTAAAAAATGAGCTGATTTAAACAAAAATTTAACCGGAATTAATTCTGTGGAATGTGTGTCAG  
TTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGGCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCA  
GCAACCAGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGT  
CAGCAACCATAGTCCCGCCCCTAACTCCGCCCCTAACTCCGCCCAGTTCGCGCCATTCTCC  
GCCCCATGGCTGACTAATTTTTTTTATTTATGAGAGGCGGAGGCCGCTCTGCCTCTGAGCTATTCCAG  
AAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTCCCGGGAGCTTGTATATCCATTT  
TCGGATCTGATCAGCACGTGTTGACAATTAATCATCGGCATAGTATATCGGCATAGTATAATACGACAAG  
GTGAGGAACATAACCATGGCCAAGCCTTTGTCTCAAGAAGAATCCACCCTCATTGAAAGAGCAACGGCTA  
CAATCAACAGCATCCCCATCTCTGAAGACTACAGCGTCGCCAGCGCAGCTCTCTCTAGCGACGGCCGCAT  
CTTCACTGGTGTCAATGTATATCATTTTTACTGGGGACCTTGTGCAGAACTCGTGGTGTGGGCACTGCT  
GCTGCTGCGGCAGCTGGCAACCTGACTTGTATCGTCGCGATCGGAAATGAGAACAGGGGCATCTTGAGCC  
CCTGCGGACGGTGCCGACAGGTGCTTCTCGATCTGCATCCTGGGATCAAAGCCATAGTGAAGGACAGTGA  
TGGACAGCCGACGGCAGTTGGGATTCGTGAATTGCTGCCCTCTGGTTATGTGTGGGAGGGCTAAGCACTT  
CGTGGCCGAGGAGCAGGACTGACACGTGCTACGAGATTTGATTCACCGCCGCTTCTATGAAAGGTTG  
GGCTTCGGAATCGTTTTCCGGGACGCCGGCTGGATGATCCTCCAGCGCGGGGATCTCATGTGGAGTTCT  
TCGCCCCACCCAACTTGTATTGTCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTTTAC  
AAATAAAGCATTTTTTCACTGCATTCTAGTTGTGGTTTGTCCAACTCATCAATGTATCTTATCATGTC  
TGTATACCGTTCGACCTCTAGCTAGAGCTTGGCGTAATCATGGTCATAGCTGTTTCTGTGTGAATTGTT  
ATCCGCTCACAAATCCACACAACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAATGAGT  
GAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCGCTTCCAGTCGGGAAACCTGTCGTGCCAGCTG  
CATTAAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCCTATTGGGCGCTCTTCCGCTTCTCGCTCA  
CTGACTCGCTGCGCTCGGTGCTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTT  
ATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGT  
AAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCT  
CAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAGATACAGGCGTTTCCCCCTGGAAGCTCCCTCGT  
GCGCTCTCCTGTTCCGACCTGCCGCTTACCGGATACCTGTCCGCTTCTCCCTTCGGGAAGCGTGCGG  
CTTTCTCAATGCTCAGCTGTAGGTATCTCAGTTCGGTGTAGGTGCTTCGCTCCAAGCTGGGCTGTGTGC  
ACGAACCCCCGTTAGCCCCGACCGCTGCGCCTTATCCGGTAACATATCGTCTTGAGTCCAACCCGGTAAG  
ACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCT  
ACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGC  
TGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGG  
TGTTTTTTTTTTGTTTGAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTT  
TCTACGGGGTCTGACGCTCAGTGGAAACGAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAA  
GGATCTTACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAAC  
TTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTTCATCC  
ATAGTTGCCTGACTCCCCGTGCTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTG  
CAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGC  
CGAGCGCAGAAGTGGTCTGCAACTTTATCCGCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGA  
GTAAGTAGTTCCGCAAGTTAATAGTTTGGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCT  
CGTCGTTTGGTATGGCTTCATTTCAGCTCCGCTTCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTT  
GTGCAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGCAAGTAAGTTGGCCGAGTGTATCA  
CTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCTATGCCATCCGTAAGATGCTTTTCTGTGACTG  
GTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGGCCGGCGTCAAT  
ACGGGATAATACCGCGCCACATAGCAGAAGTTTAAAGTGCTCATCATTGGAAAACGTTCTTCCGGGGCGA  
AACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCAACCACTGATCTT  
CAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGG  
AATAAGGGCGACACGGAATGTTGAATACTCATACTCTTCTTTTTCAATATTATTGAAGCATTTATCAG  
GGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCA  
CATTTCCCCGAAAAGTGCCACCTGACGTC

Fig. 8B




Fig. 9




Fig. 10

GACGGATCGGGAGATCTCCCGATCCCCTATGGTCGACTCTCAGTACAATCTGCTCTGATGCCGCATAGTT  
AAGCCAGTATCTGCTCCCTGCTTGTGTGTTGGAGGTCGCTGAGTAGTGCGCGAGCAAAATTTAAGCTACA  
ACAAGGCAAGGCTTGACCGACAATTGCATGAAGAATCTGCTTAGGGTTAGGCGTTTTGCGCTGCTTCGCG  
ATGTACGGGCCAGATATACGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTC  
ATTAGTTCATAGCCCATATATGGAGTTCGCGCTTACATAACTTACGGTAAATGGCCCGCTGGCTGACCG  
CCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCC  
ATTGACGTCAATGGGTGGACTATTTACGGTAACTGCCCCTTGGCAGTACATCAAGTGTATCATATGCC  
AAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTA  
TGGGACTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGC  
AGTACATCAATGGGCGTGGATAGCGGTTTGAATCACGGGGATTTCGAAGTCTCCACCCCATTTGACGTCAA  
TGGGAGTTTGTTTTGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTCCGCCCCATTGACG  
CAAATGGGCGGTAGGCGGTGACGGTGGGAGGTCTATATAAGCAGAGCTCTCTGGCTAACTAGAGAACCCA  
CTGCTTACTGGCTTATCGAAAATTAATACGACTCACTATAGGGAGACCCAAGCTGGCTAGCGTTTTAACTT  
AAGCTTACCATGGGCGCGGCACCCCGGTGACCGCCCCGCTGGCGGGCCTATCTGGAAGGTGCTGGCCA  
GCGAAGGCCAGACGGTGGCCGCGAGGCGAGGTGCTGCTGATTCTGGAAGCCATGAAGATGGAAACCGAAAT  
CCGCGCCGCGCAGGCCGGGACCGTGCAGCGGTATCGCGGTGAAAGCCGGCGACGCGGTGGCGGTGGCGGAC  
ACCTGATGACCCCTGGCGGGCTCTGGATCCGATCTGTACGACGATGACGATAAGGTACCTAGGATCCAGT  
GTGGTGAATTGATCCCTTACCAAGGGCGTCGAGTCTAGAGGGCCCGTTTAAACCCGCTGATCAGCCTC  
GACTGTGCCTTCTAGTTGCCAGCCATCTGTTGTTTGGCCCTCCCCCGTGCTTCTTGACCCCTGGAAGGT  
GCCACTCCCACTGTCTTTTCTTAATAAAATGAGGAAATTCATCGCATTTGTCTGAGTAGGTGTCTTCTA  
TTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGGA  
TGCGGTGGGCTCTATGGCTTCTGAGGCGGAAAGAACAGCTGGGGCTCTAGGGGGTATCCCCACGCGCCC  
TGTAGCGGCGCATTAAGCGCGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCCC  
TAGCGCCCGCTCCTTTTCGCTTTCTTCCCTTCTTCTCGCCACGTTTCGCGGCTTTCCCCGTCAAGCTCT  
AAATCGGGGCATCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAAACTTGATTAG  
GGTGATGGTTACGATAGTGGGCCATCGCCCTGATAGACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGT  
TCTTTAATAGTGGACTCTTGTTCAAACTGGAACAACACTCAACCTATCTCGGTCTATTCTTTTGATTT  
ATAAGGGATTTTGGGGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATTTAACAATAATTAACGCGAAT  
TAATCTGTGGAATGTGTGTGAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGGCAGGCAGAAGTATGC  
AAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTCCCCAGGCTCCCCAGGCAGGCAGAAGTAT  
GCAAGCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCCATCCCGCCCCCTAACT  
CCGCCCCAGTTCCGCCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCG  
CCTCTGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCT  
CCCGGGAGCTTGTATATCCATTTTCGGATCTGATCAGCAGCTGTTGACAATTAATCATCGGCATAGTATA  
TCGGCATAGTATAATACGACAAGGTGAGGAATAAACCATGGCCAAGCCTTTGTCTCAAGAAGAATCCAC  
CCTCATTGAAAGAGCAACGGCTACAATCAACAGCATCCCCATCTCTGAAGACTACAGCGTCGCCAGCGCA  
GCTCTCTCTAGCGACGGCCGCATCTTCACTGGTGTCAATGTATATCATTTTACTGGGGGACCTTGTGCAG  
AACTCGTGGTGCTGGGCACTGCTGCTGCTGCGGCAGCTGGCAACCTGACTTGTATCGTCGCGATCGGAAA  
TGAGAACAGGGGCATCTTGAGCCCCTGCGGACGGTGCCGACAGGTGCTTCTCGATCTGCATCCTGGGATC  
AAAGCCATAGTGAAGGACAGTGATGGACAGCCGACGGCAGTTGGGATTCTGTAATTGCTGCCCTCTGGTT  
ATGTGTGGGAGGGCTAAGCACTTCGTGGCCGAGGAGCAGGACTGACACGTGCTACGAGATTTGATTCCA  
CCGCGCCTTCTATGAAAGGTTGGGCTTCGGAATCGTTTTCCGGGACGCCGGCTGGATGATCTCCAGCG  
CGGGGATCTCATGCTGGAGTTCTTCGCCCACCCCACTTGTTTTATTGCAGCTTATAATGGTTACAAATAA  
AGCAATAGCATCACAATTTTACAATAAAGCATTTTTTTCACTGCATTCTAGTTGTGGTTTTGTCCAAC  
TCATCAATGTATCTTATCATGTCTGTATACCGTCGACCTCTAGCTAGAGCTTGGCGTAATCATGGTCATA  
GCTGTTTTCTGTGTGAAATTGTTATCCGCTCACAATTCACAACAACATACGAGCCGGAAGCATAAAGTGT  
AAAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCGCTTTCCAGT  
CGGGAACCTGTCTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTTGCGTATTGG  
GCGCTCTTCCGCTTCTCGCTCACTGACTCGCTGCGCTCGGTGCTTCGGCTGCGGCGAGCGGTATCAGCT  
CACTCAAAGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAG

Fig. 11A



GCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGA  
CGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCG  
TTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCT  
TTCTCCCTTCGGGAAGCGTGGCGCTTCTCAATGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGT  
TCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCGACCGCTGCGCCTTATCCGGTAAGTAT  
CGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCA  
GAGCGAGGTATGTAGGCGGTGCTACAGAGTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGAC  
AGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGC  
AAACAAACCACCGCTGGTAGCGGTGGTTTTTTTGTGTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGAT  
CTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAACGAAAACCTCACGTTAAGGGAT  
TTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCA  
ATCTAAAGTATATATGAGTAACTTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAG  
CGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGG  
CTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCAGCTCCAGATTTATCAGCA  
ATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAACTTTATCCGCCTCCATCCAGTCTA  
TTAATTGTTGCCGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGGCGAACGTTGTTGCCATTGC  
TACAGGCATCGTGGTGTACGCTCGTCGTTTGGTATGGCTTCATTCAGCTCCGGTTCQCAACGATCAAGG  
CGAGTTACATGATCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCCCTCCGATCGTTGTCAGAA  
GTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTATGCCATC  
CGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCG  
AGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCA  
TTGGAAAACGTTCTTCGGGGCGAAAACCTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACC  
CACTCGTGCACCCAACCTGATCTTCAGCATCTTTTACTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGA  
AGGCAAAATGCCGCAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTT  
AATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAA  
TAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACGTC

Fig. 11 B




Fig. 12




Fig. 13

TCGCGCGTTTCGGTGATGACGGTGAAAACCTCTGACACATGCAGCTCCCGGAGACGGTCACAGCTTGTCT  
GTAAGCGGATGCCGGGAGCAGACAAGCCCGTCAGGGCGCGTCAGCGGGTGTGGCGGGTGTGCGGGCTGG  
CTTAACATATGCGGCATCAGAGCAGATTGTACTGAGAGTGCACCATATGCGGTGTGAAATACCGCACAGAT  
GCGTAAGGAGAAAATACCGCATCAGGCGCCATTGCGCCATTGAGGCTGCGCAACTGTTGGGAAGGGCGATC  
GGTGCGGGCCTCTTCGCTATTACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGTTGGGTA  
ACGCCAGGGTTTTCCAGTCACGACGTTGTAAAACGACGGCCAGTGCCAGTGAATTAATTCGTTGCAGGA  
CAGGATGTGGTGCCCGATGTGACTAGCTCTTTGCTGCAGGCGCTCCTATCCTCTGGTTCCGATAAGAGAC  
CCAGAACTCCGGCCCCCACC GCCCACC GCCCACC CCATACATATGTGGTACGCAAGTAAGAGTGCCTGC  
GCATGCCCCATGTGCCCCACCAAGAGTTTTGCATCCCATACAAGTCCCCAAAGTGAGAAACCGAACCAAT  
TCTTCGCGGGCAGAACAAAAGCTTCTGCACACGTCTCCACTCGAATTTGGAGCCGGCCGGCGTGTGCAAA  
AGAGGTGAATCGAACGAAAGACCCGTGTGTAAAGCCGCGTTTTCCAAAATGTATAAAACCGAGAGCATCTG  
GCCAATGTGCATCAGTTGTGGTCAGCAGCAAAATCAAGTGAATCATCTCAGTGCAACTAAAGGGGGGATC  
TAGCGTTTTAAACTTAAGCTTACCATGGGCGCCGGCACCCCGGTGACCGCCCCGCTGGCGGGCACTATCTG  
GAAGGTGCTGGCCAGCGAAGGCCAGACGGTGGCCGAGGCGAGGTGCTGCTGATTCTGGAAGCCATGAAG  
ATGGAACCGAAATCCGCGCCGCGCAGGCGGGGACCCTGCGCGGTATCGCGGTGAAAGCCGGCGACGCGG  
TGGCGGTGCGCGACACCCTGATGACCCTGGCGGGCTCTGGATCCGATCTGTACGACGATGACGATAAGGT  
ACATCAAACAAGTTTGTACAAAAAGCTGAACGAGAAACGTAATAATGATATAAATATCAATATATTAAAT  
TAGATTTTGCATAAAAAACAGACTACATAACTGTAAACACAACATATCCAGTCACATATGGCGGCCGC  
ATTAGGCACCCAGGCTTTACACTTTATGCTTCCGGCTCGTATAATGTGTGGATTTTGGATTAGGATCCG  
GCGAGATTTTCAGGAGCTAAGGAAGCTAAATGGAGAAAAAATCACTGGATATACCACCGTTGATATAT  
CCCAATGGCATCGTAAAGAACATTTTGAGGCATTTTCAGTCAGTTGCTCAATGTACCTATAACCAGACCGT  
TCAGCTGGATATTACGGCCTTTTTAAAGACCGTAAAGAAAAAATAGCAAGTTTTATCCGGCCTTTATT  
CACATTCTTGCCCGCTGATGAATGCTCATCCGAATTCGCTATGGCAATGAAAGACGGTGAGCTGGTGA  
TATGGGATAGTGTTACCCCTTGTTACACCGTTTTCCATGAGCAAACTGAAACGTTTTATCGCTCTGGAG  
TGAATACCACGACGATTTCCGGCAGTTTCTACACATATATTCGCAAGATGTGGCGTGTACCGTGAAAC  
CTGGCCTATTTCCCTAAAGGGTTTTATTGAGAATATGTTTTTCGTCTCAGCCAATCCCTGGGTGAGTTCA  
CCAGTTTTGATTTAAACGTGGCCAATATGGACAACCTCTTCGCCCCGTTTTACCATGGGCAAAATATTA  
TACGCAAGGCGACAAGGTGCTGATGCCGCTGGCGATTGAGGTTTCATCATGCCGCTCTGTGATGGCTTCCAT  
GTCGGCAGAATGCTTAATGAATTACAACAGTACTGCGATGAGTGGCAGGGCGGGGCGTAAACGCGTGGAT  
CCGGCTTACTAAAAGCCAGATAACAGTATGCGTATTTGCGCGCTCGCGAACC GGTTGATACCCGAAGTAT  
GTCAAAAAGAGGTGTGCTATGAAGCAGCGTATTACAGTGACAGTTGACAGCGACAGCTATCAGTTGCTCA  
AGGCATATATGATGTCAATATCTCCGGTCTGGTAAGCACAACCATGCAGAATGAAGCCCGTCTGCTGCGT  
GCCGAACGCTGGAAAGCGGAAAATCAGGAAGGGATGGCTGAGGTGCGCCGGTTTTATTGAAATGAACGGCT  
CTTTTGCTGACGAGAACAGGGACTGGTGAATGCAGTTAAGGTTTACACCTATAAAAGAGAGAGCCGTT  
ATCGTCTGTTTGTGGATGTACAGAGTGATATTATTGACACGCCCGGGCGACGGATGGTGATCCCCCTGGC  
CAGTGCACGCTGCTGTCAGATAAAGTCTCCCGTGAACCTTTACCCGGTGGTGATATCGGGGATGAAAGC  
TGGCGCATGATGACCACCGATATGGCCAGTGTGCCGGTCTCCGTTATCGGGGAAGAAGTGGGTGATCTCA  
GCCACCGCGAAAATGACATCAAAAACGCCATTAACTGATGTTCTGGGGAATATAAATGTCAGGCTCCGT  
TATACACAGCCAGTCTGCAGGTCGACCATAGTGACTGGATATGTTGTGTTTTACAGTATTATGTAGTCTG  
TTTTTTATGCAAAATCTAATTTAATATATTGATATTTATATCATTTTACGTTTCTCGTTTCAGCTTTCTTG  
TACAAAGTGGTGATAATTAATTAAGATCTAGAGGGCCCGTTAAACCCGCTGATCAGCCTCGACTGTGCC  
TTCTAAGATCCAGACATGATAAGATACATTGATGAGTTTGGACAAACCACAACCTAGAATGCAGTGAAAAA  
AATGCTTTATTTGTGAAATTTGTGATGCTATTGCTTTATTTGTAACCATTATAAGCTGCAATAAACAAGT  
TAACAACAACAATTGCATTCATTTTATGTTTCAGGTTTCAGGGGGAGGTGTGGGAGGTTTTTTAAAGCAAG  
TAAACCTCTACAAATGTGGTATGGCTGATTATGATCAGTCGACCTGCAGGCATGCAAGCTTGGCGTAAT  
CATGGTCATAGCTGTTTCCGTGTGTGAAATTTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAG  
CATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCC  
GCTTTCCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGGCGGTT  
TGGTATTTGGGCGCTCTTCCGCTTCTCGCTCACTGACTCGCTGCGCTCGGTCTCGGCTGCGGCGAGC  
GGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATG

Fig. 14A

TGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCC  
GCCCCCTGACGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAG  
ATACCAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATAC  
CTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATCTCAGTTCGG  
TGAGGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCGACCGCTGCGCCTTATC  
CGGTAACATATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAAC  
AGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACA  
CTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTC  
TTGATCCGGCAAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGA  
AAAAAGGATCTCAAGAAGATCCTTTGATCTTTCTACGGGGTCTGACGCTCAGTGGAAACGAAAACCTCAC  
GTAAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAAAATGAAG  
TTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCA  
CCTATCTCAGCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGA  
TACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGA  
TTTATCAGCAATAAACAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAACTTTATCCGCCTCC  
ATCCAGTCTATTAATTGTTGCCGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGCGCAACGTTG  
TTGCCATTGCTACAGGCATCGTGGTGTACGCTCGTCTGTTTGGTATGGCTTCATTACGCTCCGGTTCCCA  
ACGATCAAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCCCTCCGATC  
GTTGTCAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTG  
TCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTAT  
GCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAA  
GTGCTCATCATTTGGAACGTTCTTCGGGGCGAAAACCTCTCAAGGATCTTACCGCTGTTGAGATCCAGTT  
CGATGTAACCCACTCGTGCACCCAACCTGATCTTCAGCATCTTTTACTTTACCAGCGTTTCTGGGTGAGC  
AAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTC  
TTCCTTTTCAATATTATTGAAGCATTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTA  
TTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGACGTCTAAGAAAC  
CATTATTATCATGACATTAACCTATAAAAAATAGGCGTATCACGAGGCCCTTTCGT

Fig. 14B




Fig. 15

- Shaded regions correspond to those DNA sequences transferred from the entry clone into the pET104-DEST™ vector by recombination. Non-shaded regions are derived from the pET104-DEST™ vector.
- Bases 568 and 2230 of the pET104-DEST™ sequence are marked.
- The biotin binding site is labeled with a \*.

2121 ATAGGCGGCCA GCAACCGCAC CTGTGGCCGC EGTGATGCGC GCCACGATGC GTCCGGCGTA GAGGATCGAG AATCTGATCC

T7 promoter/primer site  
T7 promoter lac operator

201 CGCGAAATTATA ATACGACTCA CTATAGGGGA ATTGTGAGCG GATAACMAAT CCCCTCTAGA AATAATTTTG TTTAACTTTA

Biotag™

RBS Met Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp Lys Val  
281 AGAAGGAGAT ATACAT ATG GGC GCC GGC ACC CCG GTG ACC GCC CCG CTG GCG GGC ACT ATC TGG AAG GTG

Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu Leu Ile Leu Gln Ala Met Lys Met  
351 CTG GCC AGC GAA GGC CAG ACC GTG GCC GCA GGC GAG GTG CTG CTG ATT CTG GAA GCC ATG AAG ATG

Biotin binding site

417 Glu Thr Glu Ile Arg Ala Ala Gln Ala Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala  
GAA ACC GAA ATC CGC GCC GCG CAG GCC GGG ACC GTG CCG GGT ATC CCG GTG AAA GCC GGC GAC CCG

Biotag™ forward priming site EK recognition site

483 Val Ala Val Gly Asp Thr Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr Asp Asp Asp  
GTG GCG GTC GGC GAC ACC CTG ATG ACC CTG CCG GGC TCT GGA TCC GAT CTG TAC GAC GAT GAC GAT

688 2219

Lys Gly Ile Ile Thr Ser Leu Tyr Lys Lys Ala Gly ... \*\*  
549 AAG GGA ATT ATC ACA AGT TT [REDACTED] GENE [REDACTED] TTGTACAAG  
CCT TAA TAG TGT TCA AAC ATG TTT [REDACTED] TTTC

EK cleavage site attB1 attB2 T7 reverse priming site

2241 TGGTGATAAT TAATTAAGAT AGCTCAGATC CGGCTGCTAA CAAGGCCGA AAGGAAGCTG AGTTGGCTGC TGCCACCGET  
ACCACTATTA

2321 GAGCAATAAC TAGCATAC

Fig. 16




Fig.17



**Flow Chart**

The flow chart below describes the general steps required to clone and express your blunt-end PCR product.




Fig. 18




Fig. 19

- Shaded regions correspond to those DNA sequences transferred from the entry clone into the pcDNA6/Biotag-DEST™ vector by recombination. Non-shaded regions are derived from the pcDNA6/Biotag-DEST™ vector.
- Bases 1191 and 2853 of the pcDNA6/Biotag-DEST™ sequence are marked.
- The biotin binding site is labeled with a \*.
- Potential stop codons are underlined.




Fig. 20

**Flow Chart**

The flow chart below outlines the experimental steps necessary to clone and express your blunt-end PCR product.




Fig. 21




Fig. 22

- Shaded regions correspond to those DNA sequences transferred from the entry clone into the pMT/Biotag™-DEST vector by recombination. Non-shaded regions are derived from the pMT/Biotag™-DEST vector.
- Bases 1135 and 2797 of the pMT/Biotag™-DEST sequence are marked.
- The biotin binding site is labeled with a \*.
- Potential stop codons are underlined.

5' end of metallothionein promoter

411 CGTTGCAGGA CAGGATGTGG TGCCCGATGT GACTAGCTCT TTGCTGCAGG CCGTCCTATC CTCTGGTTCC GATAAGAGAC CCAGAATCC

Metal regulatory region

501 GGCCCCCACC CCCCCACCG CCCCCATA CATATGTGGT ACGCAAGTAA GAGTGCTGCG GCATGCCCCA TGTGCCCCAC CAAGATTTT

Metal regulatory

region

591 GCATCCCAT AAGTCCCCA AAGTGAGAA CCGAACCAAT TCTTCGCGGG CAGAACAAA GCTTCTGCAC AGGTCTCCAC TCGAATTTGG

Metal regulatory regions

681 AGCCGECGG CGTGTGCAA AGAGGTGAAT CGACCAAG ACCCGTGTGT AAAGCCGGT TTCCAAATG TATAAACCG AGAGCATCTG

Metal regulatory region

Metal regulatory regions

TATA

Start of transcription

771 GCCAATGTGC ATCAGTTGTG GTCAGCAGCA AAATCAAGTG ATCATCTCA GTGCAACTAA AGGGGGGATC TAGCGTTTAA ACTTAAAGCTT

Biolog™

861 Met Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp Lys Val Leu Ala Ser Glu  
ACC ATG GGC GCC GGC ACC CCG GTG ACC GCC CCG CTG GCG GGC ACT ATC TGG AAG GTG CTG GCC AGC GAA

Gly Gln Thr Val Ala Ala Gly Glu Val Leu Leu Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg  
930 GGC CAG ACG GTG GCC GCA GGC GAG GTC CTG CTG ATT CTG GAA GCC ATG AAG ATG GAA ACC GAA ATC CGC

Blotin binding site

999 Ala Ala Gln Ala Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala Val Gly Asp Thr  
GCC GCG CAG GCC GGG ACC GTG CCG GGT ATC GCG GTG AAA GCC GGC GAC GCG GTG GCG GTC GGC GAC ACC

EK recognition site

EK cleavage site

1135

1068 Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr Asp Asp Asp Asp Lys Val His Gln Thr Ser Leu  
CTG ATG ACC CTG GCG GGC TCT GGA TCC GAT CTG TAC GAC GAT GAC GAT AAG GTA CAT CAA ACA AGT TGT  
CAT GTA GTT TGT TCA AAC

1787

Tyr Lys Lys Ala Gly ... \*\*\*

1137

ATG TTT ... GENE ... TTTCACC ACTATAATT

al(B1)

al(B2)

Fig. 23




Fig. 24




Fig. 25



MAIL STOP PCT

Applicant: INVITROGEN CORPORATION

Docket: 0942.551PC03

PCT Int'l. Appl. No: PCT/US03/21339

Atty: RWE/FRC

PCT Int'l. Filing Date: 08 July 2003

For: **METHODS AND COMPOSITIONS FOR THE PRODUCTION,  
IDENTIFICATION AND PURIFICATION OF FUSION PROTEINS**

When receipt stamp is placed hereon, the USPTO acknowledges receipt of the following documents:

Submission of Formal Drawings Under PCT Rule 11.13 (32 sheets, depicting figures 1 through 25) for the above application.

RWE/FRC/CAG/lmw

IN THE PCT/US RECEIVING OFFICE  
UNDER THE PATENT COOPERATION TREATY

In re International Application of: INVITROGEN CORPORATION

International Application No.: PCT/US03/21339

International Filing Date: 08 July 2003

Title: **METHODS AND COMPOSITIONS FOR THE PRODUCTION, IDENTIFICATION AND  
PURIFICATION OF FUSION PROTEINS**


Commissioner for Patents  
Mail Stop PCT  
Alexandria, VA 22313-1450

**SUBMISSION OF FORMAL DRAWINGS  
UNDER PCT RULE 11.13**

Sir:

In accordance with PCT Rule 11.13, the undersigned hereby submits 32 sheets of formal drawings (comprising figures 1 through 25) in connection with the above international application. The undersigned requests that the formal figures be transmitted to the International Bureau as soon as possible for technical preparations of the international publication document.

Respectfully submitted,



Robert W. Esmond  
Attorney for Applicant  
Reg. No. 32,893

Date: 13 November 2003

RWE/FRC/CAG/lmw  
Enclosures:  
Formal Drawings - 32 Sheets

Applicant: INVITROGEN CORPORATION.

MAIL STOP PCT

PCT Int'l. Appl. No: PCT/US03/21339

Docket: 0942.551PC03

Int'l. Filing Date: 8 July 2003

Atty: RWE/FRC

For: METHODS AND COMPOSITIONS FOR THE PRODUCTION,  
IDENTIFICATION AND PURIFICATION OF FUSION PROTEINS

When receipt stamp is placed hereon, the USPTO acknowledges receipt of the following documents:

SUBMISSION OF NUCLEOTIDE AND/OR AMINO ACID SEQUENCE LISTING COMPLYING WITH STANDARD,  
SUBMITTING A DISKETTE CONTAINING SEQUENCE LISTING AND PAPER COPY.

RWE/FRC/CAG/lmw

IN THE INTERNATIONAL SEARCHING AUTHORITY (ISA/US)

UNDER THE PATENT COOPERATION TREATY

In re International Application of: INVITROGEN CORPORATION

International Application No.: PCT/US03/21339

International Filing Date: 08 July 2003

Title: **METHODS AND COMPOSITIONS FOR THE PRODUCTION,  
IDENTIFICATION AND PURIFICATION OF FUSION PROTEINS**

Commissioner for Patents  
Mail Stop PCT  
Alexandria, VA 22313-1450


ATTENTION: Lissie Marquis

SUBMISSION OF NUCLEOTIDE  
AND/OR AMINO ACID SEQUENCE LISTING COMPLYING WITH STANDARD  
(under PCT Rule 13<sup>ter</sup>.1(a) and (c) and Administrative Instructions,  
Section 208 and Annex C)

Ms. Marquis:

The undersigned hereby submits a diskette containing the sequence listing and a paper copy thereof. The undersigned confirms that the diskette containing the sequence listing does not include matter which goes beyond the disclosure of the international application as filed. Moreover, the paper copy of the sequence listing and the computer readable form are the same.

Respectfully submitted,

  
Robert W. Esmond  
Attorney for Applicant  
Reg. No. 32,893

Date: 13 November 2003

RWE/FRC/CAG/lmw

Enclosures:  
Paper Copy of Sequence Listing  
Diskette

SEQUENCE LISTING

<110> Invitrogen Corporation

<120> Methods and Compositions for the Production, Identification and Purification of Fusion Proteins

<130> 0942.551PC03

<140> PCT/US03/21339

<141> 2003-07-08

<150> 60/393,756

<151> 2002-07-08

<150> 60/396,627

<151> 2002-07-19

<150> 60/417,172

<151> 2002-10-10

<160> 34

<170> PatentIn version 3.2

<210> 1

<211> 7618

<212> DNA

<213> Artificial

<220>

<223> pET104-DEST

<400> 1

caaggagatg gcgccaaca gtcccccggc cacggggcct gccaccatac ccacgccgaa	60
acaagcgctc atgagccga agtggcgagc ccatcttcc ccatcggtga tgctggcgat	120
ataggcgcca gcaaccgcac ctgtggcgcc ggtgatgccg gccacgatgc gtccggcgta	180
gaggatcgag atctcgatcc cgcgaaatta atacgactca ctatagggga attgtgagcg	240
gataacaatt cccctctaga aataattttg tttaacttta agaaggagat atacatatgg	300
gcgccggcac cccggtgacc gccccgctgg cgggcactat ctggaaggtg ctggccagcg	360
aaggccagac ggtggccgca ggcgaggtgc tgctgattct ggaagccatg aagatggaaa	420
ccgaaatccg cgccgcgcag gccgggaccg tgcgcggtat cgcggtgaaa gccggcgacg	480
cgggtggcggc cggcgacacc ctgatgaccc tggcgggctc tggatccgat ctgtacgacg	540
atgacgataa ggaattatc acaagtttgt acaaaaaagc tgaacgagaa acgtaaaatg	600
atataaatat caatatatta aattagattt tgcataaaaa acagactaca taatactgta	660
aaacacaaca tatccagtca ctatggcggc cgcattaggc accccaggct ttacacttta	720
tgcttccggc tcgtataatg tgtggatttt gagttaggat ccggcgagat tttcaggagc	780
taagggaagct aaaatggaga aaaaaatcac tggatatacc accgttgata tatcccaatg	840

gcatcgtaaa	gaacattttg	aggcattttca	gtcagttgct	caatgtacct	ataaccagac	900
cgttcagctg	gatattacgg	ccttttttaa	gaccgtaaag	aaaaataagc	acaagtttta	960
tccggccttt	attcacattc	ttgcccgcct	gatgaatgct	catccggaat	tccgtatggc	1020
aatgaaagac	ggtgagctgg	tgatatggga	tagtgttcac	ccttgttaca	ccgtttttcca	1080
tgagcaaact	gaaacgtttt	catcgctctg	gagtgaatac	cacgacgatt	tccggcagtt	1140
tctacacata	tattcgcaag	atgtggcgctg	ttacggtgaa	aacctggcct	atttccttaa	1200
agggtttatt	gagaatatgt	ttttcgtctc	agccaatccc	tgggtgagtt	tcaccagttt	1260
tgatttaaac	gtggccaata	tggacaactt	cttcgcccc	gttttcacca	tgggcaaata	1320
ttatacgcaa	ggcgacaagg	tgctgatgcc	gctggcgatt	caggttcctc	atgccgtctg	1380
tgatggcttc	catgtcggca	gaatgcttaa	tgaattacaa	cagtactgcg	atgagtggca	1440
gggcggggcg	taaacgcgtg	gatccggctt	actaaaagcc	agataacagt	atgcgtattt	1500
gcgcgcaccg	gtgctagcgt	atacccgaag	tatgtcaaaa	agaggtgtgc	tatgaagcag	1560
cgtattacag	tgacagttga	cagcgacagc	tatcagttgc	tcaaggcata	tatgatgtca	1620
atatctccgg	tctggtaagc	acaaccatgc	agaatgaagc	ccgtcgtctg	cgtgccgaac	1680
gctggaaagc	ggaaaatcag	gaagggatgg	ctgaggtcgc	ccggtttatt	gaaatgaacg	1740
gctcttttgc	tgacgagaac	agggactggt	gaaatgcagt	ttaagggtta	cacctataaa	1800
agagagagcc	gttatcgtct	gtttgtggat	gtacagagtg	atattattga	cacgcccggg	1860
cgacggatgg	tgatccccct	ggccagtgca	cgtctgctgt	cagataaagt	ctcccgtgaa	1920
ctttaccg	tggtgcatat	cggggatgaa	agctggcgca	tgatgaccac	cgatatggcc	1980
agtgtgccgg	tctccgttat	cggggaagaa	gtggctgac	tcagccaccg	cgaaaatgac	2040
atcaaaaacg	ccattaacct	gatgttctgg	ggaatataaa	tgtcaggctc	cgttatacac	2100
agccagtctg	caggctcgacc	atagtactg	gatatgttgt	gttttacagt	attatgtagt	2160
ctgtttttta	tgcaaaatct	aatttaatat	attgatattt	atatcatttt	acgtttctcg	2220
ttcagctttc	ttgtacaaag	tggtgataat	taattaagat	agctcagatc	cggctgctaa	2280
caaagcccga	aaggaagctg	agttggctgc	tgccaccgct	gagcaataac	tagcataacc	2340
ccttggggcc	tctaaacggg	tcttgagggg	ttttttgctg	aaaggaggaa	ctatatccgg	2400
atatcccgcg	agaggcccgg	cagtaccggc	ataaccaagc	ctatgcctac	agcatccagg	2460
gtgacggtgc	cgaggatgac	gatgagcgca	ttgttagatt	tcatacacgg	tgccctgactg	2520
cgtttagcaat	ttaactgtga	taaactaccg	cattaaagct	agcttatcga	tgataagctg	2580
tcaaacatga	gaattaattc	ttgaagacga	aagggcctcg	tgatacgctt	atttttatag	2640
gttaatgtca	tgataataat	ggtttcttag	acgtcaggtg	gcacttttctg	gggaaatgtg	2700

cgcggaaccc ctatttgttt atttttctaa atacattcaa atatgtatcc gctcatgaga 2760  
caataaccct gataaatgct tcaataatat tgaaaaagga agagtatgag tattcaacat 2820  
ttccgtgtcg cccttattcc cttttttgcg gcattttgcc ttctgtttt tgctcaccca 2880  
gaaacgctgg tgaaagtaaa agatgctgaa gatcagttgg gtgcacgagt gggttacatc 2940  
gaactggatc tcaacagcgg taagatcctt gagagttttc gccccgaaga acgttttcca 3000  
atgatgagca ctttttaaagt tctgctatgt ggcgcggtat tatcccgtgt tgacgccggg 3060  
caagagcaac tcggtcgccg catacactat tctcagaatg acttggttga gtactcacca 3120  
gtcacagaaa agcatcttac ggatggcatg acagtaagag aattatgcag tgctgccata 3180  
accatgagtg ataacactgc ggccaactta cttctgacaa cgatcggagg accgaaggag 3240  
ctaaccgctt ttttgacaaa catgggggat catgtaactc gccttgatcg ttgggaaccg 3300  
gagctgaatg aagccatacc aaacgacgag cgtgacacca cgatgcctgc agcaatggca 3360  
acaacgttgc gcaaactatt aactggcgaa ctacttactc tagcttcccg gcaacaatta 3420  
atagactgga tggaggcgga taaagttgca ggaccacttc tgcgctcggc ccttccggct 3480  
ggctggttta ttgctgataa atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca 3540  
gcaactggggc cagatggtaa gccctcccgt atcgtagtta tctacacgac ggggagtcag 3600  
gcaactatgg atgaacgaaa tagacagatc gctgagatag gtgcctcact gattaagcat 3660  
tggttaactgt cagaccaagt ttactcatat atactttaga ttgatttaaa acttcatttt 3720  
taatttaaaa ggatctaggt gaagatcctt tttgataatc tcatgaccaa aatcccttaa 3780  
cgtgagtttt cgttccactg agcgtcagac cccgtagaaa agatcaaagg atcttcttga 3840  
gatccttttt ttctgcgct aatctgctgc ttgcaaacia aaaaaccacc gctaccagcg 3900  
gtggttttgt tgccggatca agagctacca actctttttc cgaaggtaac tggcttcagc 3960  
agagcgcaga taccaaatac tgtccttcta gtgtagccgt agttaggcca ccacttcaag 4020  
aactctgtag caccgcctac atacctcgt ctgctaacc tgttaccagt ggctgctgcc 4080  
agtggcgata agtcgtgtct taccgggttg gactcaagac gatagttacc ggataaggcg 4140  
cagcggtcgg gctgaacggg gggttcgtgc acacagccca gcttggagcg aacgacctac 4200  
accgaactga gatacctaca gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga 4260  
aaggcggaca ggtatccggt aagcggcagg gtcggaacag gagagcgcac gagggagctt 4320  
ccagggggaa acgcctggta tctttatagt cctgtcgggt ttgccacct ctgacttgag 4380  
cgtcgatttt tgtgatgctc gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg 4440  
gcctttttac ggttccctggc cttttgctgg ccttttgctc acatgttctt tctgegtta 4500  
tcccctgatt ctgtggataa ccgtattacc gcctttgagt gagctgatac cgctcgcgcg 4560

agccgaacga ccgagcgcag cgagtcagtg agcgaggaag cggaagagcg cctgatgcgg	4620
tattttctcc ttacgcattct gtgcggtatt tcacaccgca tatatggtgc actctcagta	4680
caatctgctc tgatgccgca tagttaagcc agtatacact ccgctatcgc tacgtgactg	4740
ggtcattggct gcgccccgac acccgccaac acccgctgac gcgcccctgac gggcttgtct	4800
gctccccgca tccgcttaca gacaagctgt gaccgtctcc gggagctgca tgtgtcagag	4860
gttttcaccg tcatcaccga aacgcgcgag gcagctgcgg taaagctcat cagcgtggtc	4920
gtgaagcgat tcacagatgt ctgcctgttc atccgcgtcc agctcgttga gtttctccag	4980
aagcgttaat gtctggcttc tgataaagcg ggccatgtta agggcggttt tttcctgttt	5040
ggtcactgat gcctccgtgt aagggggatt tctgttcattg ggggtaatga taccgatgaa	5100
acgagagagg atgctcacga tacgggttac tgatgatgaa catgcccggg tactggaacg	5160
ttgtgagggt aaacaactgg cggatatgat gcggcgggac cagagaaaaa tctctcaggg	5220
tcaatgccag cgcttcgtta atacagatgt aggtgttcca cagggtagcc agcagcatcc	5280
tgcatgacag atccggaaca taatggtgca gggcgctgac ttccgcgttt ccagacttta	5340
cgaaacacgg aaaccgaaga ccattcatgt tggtgctcag gtcgcagacg ttttgacgca	5400
gcagtcgctt cacgttcgct cgcgtatcgg tgattcattc tgctaaccag taaggcaacc	5460
ccgccagcct agccgggtcc tcaacgacag gagcacgac atgcgcaccc gtggccagga	5520
cccaacgctg cccgagatgc gccgcgtgcg gctgctggag atggcggacg cgatggatat	5580
gttctgcca ggggttggtt gcgcattcac agttctccgc aagaattgat tggctccaat	5640
tcttgagtg gtgaatccgt tagcgagggt ccgcccgtt ccattcaggt cgaggtggcc	5700
cggctccatg caccgcgacg caacgcgggg aggcagacaa ggtatagggc ggcgcctaca	5760
atccatgcca acccgttcca tgtgctcgcc gaggcggcat aaatcgccgt gacgatcagc	5820
ggtccagtga tcgaagttag gctggtaaga gccgcgagcg atccttgaag ctgtccctga	5880
tggtcgtcat ctacctgcct ggacagcatg gcctgcaacg cgggcatccc gatgccgccc	5940
gaagcgagaa gaatcataat ggggaaggcc atccagcctc gcgtcgcgaa cgccagcaag	6000
acgtagccca gcgcgtcggc cgccatgccg gcgataatgg cctgcttctc gccgaaacgt	6060
ttggtggcgg gaccagtgc gaaggcttga gcgagggcgt gcaagattcc gaataccgca	6120
agcgacaggc cgatcatcgt cgcgtccag cgaaagcggc cctcgccgaa aatgaccag	6180
agcgtgccg gcacctgtcc tacgagttgc atgataaaga agacagtcatt aagtgcggcg	6240
acgatagtca tgccccgcgc ccaccggaag gagctgactg ggttgaaggc tctcaagggc	6300
atcggtcgag atcccgggtgc ctaatgagtg agctaactta cattaattgc gttgcgctca	6360
ctgcccgtt tccagtcggg aaacctgtcg tgccagctgc attaatgaat cggccaacgc	6420



gcgggggagag gcgggtttgcg tattggggcgc caggggtgggt tttcttttca ccagtgcagac 6480  
gggcaacagc tgattgccct tcaccgcctg gccctgagag agttgcagca agcgggtccac 6540  
gctgggtttgc cccagcaggc gaaaatcctg tttgatgggtg gttaacggcg ggatataaca 6600  
tgagctgtct tcggtatcgt cgtatccac taccgagata tccgcaccaa cgcgcagccc 6660  
ggactcggta atggcgcgca ttgcgcccag cgccatctga tcgttggcaa ccagcatcgc 6720  
agtgggaacg atgccctcat tcagcatttg catggtttgt tgaaaaccgg acatggcact 6780  
ccagtcgcct tcccgttccg ctatcggctg aatttgattg cgagtgcgat atttatgcc 6840  
gccagccaga cgcagacgcg ccgagacaga acttaatggg cccgctaaca gcgcgatttg 6900  
ctggtgacct aatgcgacca gatgctccac gccagtcgc gtaccgtctt catgggagaa 6960  
aataatactg ttgatgggtg tctggtcaga gacatcaaga aataacgccg gaacattagt 7020  
gcaggcagct tccacagcaa tggcatcctg gtcattccagc ggatagttaa tgatcagccc 7080  
actgacgcgt tgcgcgagaa gattgtgcac cgccgcttta caggcttcga cgccgcttcg 7140  
ttctaccatc gacaccacca cgctggcacc cagttgatcg gcgcgagatt taatcgccgc 7200  
gacaatttgc gacggcgcggt gcagggccag actggagggtg gcaacgcaa tcagcaacga 7260  
ctgtttgccc gccagttgtt gtgccacgcg gttgggaatg taattcagct ccgccatcgc 7320  
cgcttccact ttttcccgcg ttttcgcaga aacgtggctg gcctggttca ccacgcggga 7380  
aacggtctga taagagacac cggcactc tgcgacatcg tataacgtta ctggtttcac 7440  
attcaccacc ctgaattgac tctcttcgg gcgctatcat gccataccgc gaaagggttt 7500  
gcgccattcg atggtgtccg ggatctcgac gctctccctt atgcgactcc tgcattagga 7560  
agcagcccag tagtaggttg aggccgttga gcaccgccgc cgcaaggaat ggtgcatg 7618

<210> 2  
<211> 5934  
<212> DNA  
<213> Artificial

<220>  
<223> pET104/D-TOPO

<400> 2  
caaggagatg gcgcccaca gtccccggc cacggggcct gccaccatac ccacgccgaa 60  
acaagcgctc atgagcccga agtggcgagc ccgatcttcc ccatcgggtga tgtcggcgat 120  
ataggcgcca gcaaccgcac ctgtggcgcc ggtgatgccg gccacgatgc gtccggcgta 180  
gaggatcgag atctcgatcc cgcgaaatta atacgactca ctatagggga attgtgagcg 240  
gataacaatt cccctctaga aataattttg ttttaacttta agaaggagat atacatatgg 300  
gcgccggcac cccggtgacc gccccgctgg cgggcactat ctggaagggtg ctggccagcg 360

aaggccagac ggtggccgca ggcgaggtgc tgctgattct ggaagccatg aagatggaaa	420
ccgaaatccg cgccgcgcag gccgggaccg tgcgcggtat cgcggtgaaa gccggcgacg	480
cggtggcggt cggcgacacc ctgatgaccc tggcgggctc tggatccgat ctgtacgacg	540
atgacgataa ggggaattgat cccttcacca agggcgagct cagatccggc tgctaacaaa	600
gcccgaagg aagctgagtt ggctgctgcc accgctgagc aataactagc ataaccctt	660
ggggcctcta aacgggtctt gaggggtttt ttgctgaaag gaggaactat atccggatat	720
cccgaagag gcccggcagt accggcataa ccaagcctat gcctacagca tccaggggtga	780
cggtgccgag gatgacgatg agcgcattgt tagatttcat acacggtgcc tgactgcgtt	840
agcaatttaa ctgtgataaa ctaccgcatt aaagctagct tatcgatgat aagctgtcaa	900
acatgagaat taattcttga agacgaaagg gcctcgtgat acgcctatct ttataggtta	960
atgtcatgat aataatgggt tcttagacgt cagggtggcac ttttcgggga aatgtgcgcg	1020
gaacccttat ttgtttatct ttctaaatac attcaaatac gtatccgctc atgagacaat	1080
aaccctgata aatgcttcaa taatattgaa aaaggaagag tatgagtatt caacatttcc	1140
gtgtgcctt tattcccttt tttgcggcat tttgccttcc tgtttttgct caccagaaa	1200
cgctggtgaa agtaaaagat gctgaagatc agttgggtgc acgagtgggt tacatcgaac	1260
tggatctcaa cagcggtgaa atccttgaga gttttcgccc cgaagaacgt tttccaatga	1320
tgagcacttt taaagttctg ctatgtggcg cggattatc ccggtgtgac gccgggcaag	1380
agcaactcgg tcgccgcata cactattctc agaatgactt ggttgagtac tcaccagtca	1440
cagaaaagca tcttacggat ggcatgacag taagagaatt atgcagtgct gccataacca	1500
tgagtgataa cactgcggcc aacttacttc tgacaacgat cggaggaccg aaggagctaa	1560
ccgctttttt gcacaacatg ggggatcatg taactcgcct tgatcggttg gaaccggagc	1620
tgaatgaagc cataccaaac gacgagcgtg acaccacgat gcctgcagca atggcaacaa	1680
cgttgcgcaa actattaact ggcgaaactac ttactctagc tccccggcaa caattaatag	1740
actggatgga ggcggataaa gttgcaggac cacttctgcg ctcgccctt ccggtcggt	1800
ggtttattgc tgataaatct ggagccggtg agcgtgggtc tcgcggtatc attgcagcac	1860
tggggccaga tggtaagccc tcccgtatcg tagttatcta caccgacggg agtcaggcaa	1920
ctatggatga acgaaataga cagatcgctg agataggtgc ctactgatt aagcattgg	1980
aactgtcaga ccaagtttac tcatatatac tttagattga tttaaaactt cattttta	2040
ttaaaaggat ctaggtgaag atcctttttg ataattctcat gacccaaatc ccttaacgtg	2100
agttttcggt cactgagcg tcagaccccg tagaaaagat caaaggatct tcttgagatc	2160
ctttttttct gcgcgtaatc tgctgcttgc aaacaaaaaa accaccgcta ccagcggtg	2220

tttgtttgcc	ggatcaagag	ctaccaactc	tttttccgaa	ggtaactggc	ttcagcagag	2280
cgcagatacc	aaatactgtc	cttctagtgt	agccgtagtt	aggccaccac	ttcaagaact	2340
ctgtagcacc	gcctacatac	ctcgtctctg	taatcctgtt	accagtggct	gctgccagtg	2400
gcgataagtc	gtgtcttacc	gggttggact	caagacgata	gttaccggat	aaggcgcagc	2460
ggtcgggctg	aacggggggt	tcgtgcacac	agcccagctt	ggagcgaacg	acctacaccg	2520
aactgagata	cctacagcgt	gagctatgag	aaagcgccac	gcttcccga	gggagaaaagg	2580
cggacaggta	tccggtaaag	ggcagggtcg	gaacaggaga	gcgcacgagg	gagcttccag	2640
ggggaaacgc	ctggtatctt	tatagtcctg	tcgggtttcg	ccacctctga	cttgagcgtc	2700
gatttttgtg	atgctcgtca	ggggggcgga	gcctatggaa	aaacgccagc	aacgcggcct	2760
ttttacgggt	cctggccttt	tgctggcctt	ttgctcacat	gttctttcct	gcgttatccc	2820
ctgattctgt	ggataaccgt	attaccgcct	ttgagtgagc	tgataccgct	cgccgcagcc	2880
gaacgaccga	gcgcagcgag	tcagtgcgag	aggaagcgga	agagcgccct	atgcgggtatt	2940
ttctccttac	gcctctgtgc	ggtatttcac	accgcatata	tggtgcactc	tcagtacaat	3000
ctgctctgat	gccgcatagt	taagccagta	tacactccgc	tatcgctacg	tgactgggtc	3060
atggctgcgc	cccgcacccc	gccaacaccc	gctgacgcgc	cctgacgggc	ttgtctgctc	3120
ccggcatccg	cttacagaca	agctgtgacc	gtctccggga	gctgcatgtg	tcagaggttt	3180
tcaccgtcat	caccgaaacg	cgcgaggcag	ctgcggtaaa	gctcatcagc	gtggtcgtga	3240
agcgattcac	agatgtctgc	ctgttcatcc	gcgtccagct	cgttgagttt	ctccagaagc	3300
gttaatgtct	ggcttctgat	aaagcgggcc	atgttaaggg	cggttttttc	ctgtttggtc	3360
actgatgcct	ccgtgtaagg	gggatttctg	ttcatggggg	taatgatacc	gatgaaacga	3420
gagaggatgc	tcacgatacg	ggttactgat	gatgaacatg	cccggttact	ggaacgttgt	3480
gagggtaaac	aactggcggg	atggatgcgg	cgggaccaga	gaaaaatcac	tcagggtcaa	3540
tgccagcgct	tcgttaatac	agatgtaggt	gttccacagg	gtagccagca	gcctcctgcg	3600
atgcagatcc	ggaacataat	ggtgcagggc	gctgacttcc	gcgtttccag	actttacgaa	3660
acacggaaac	cgaagaccat	tcattgttgt	gctcaggctg	cagacgtttt	gcagcagcag	3720
tcgcttcacg	ttcgctcgcg	tatcggtgat	tcattctgct	aaccagtaag	gcaaccccg	3780
cagcctagcc	gggtcctcaa	cgacaggagc	acgatcatgc	gcacccgtgg	ccaggacca	3840
acgctgcccc	agatgcgccg	cgtgcggctg	ctggagatgg	cggacgcgat	ggatatgttc	3900
tgccaagggg	tggtttgcgc	attcacagtt	ctccgcaaga	attgattggc	tccaattctt	3960
ggagtgggtga	atccgttagc	gaggtgccgc	cggcttccat	tcaggctcag	gtggcccggc	4020
tccatgcacc	gcgacgcaac	gcggggaggc	agacaaggta	tagggcggcg	cctacaatcc	4080

atgccaaacc	gttccatgtg	ctcgccgagg	cggcataaat	cgccgtgacg	atcagcggtc	4140
cagtgatcga	agttaggctg	gtaagagccg	cgagcgatcc	ttgaagctgt	cctgatgggt	4200
cgtcatctac	ctgcctggac	agcatggcct	gcaacgcggg	catcccgatg	ccgccggaag	4260
cgagaagaat	cataatgggg	aaggccatcc	agcctcgcg	cgcgaaacgc	agcaagacgt	4320
agcccagcgc	gtcggccgcc	atgccggcga	taatggcctg	cttctcgccg	aaacgtttgg	4380
tggcgggacc	agtgcgaag	gcttgagcga	gggcgtgcaa	gattccgaat	accgcaagcg	4440
acaggccgat	catcgtcg	ctccagcgaa	agcggtcctc	gccgaaaatg	accagagcgc	4500
ctgccggcac	ctgtcctacg	agttgcatga	taaagaagac	agtcataagt	gcggcgacga	4560
tagtcatgcc	ccgcgcccac	cggaaggagc	tgactgggtt	gaaggctctc	aagggcacgc	4620
gtcgagatcc	cggtgcctaa	tgagtgaagt	aacttacatt	aattgcgttg	cgctcactgc	4680
ccgctttcca	gtcgggaaac	ctgtcgtgcc	agctgcatta	atgaatcggc	caacgcgcgc	4740
ggagaggcgc	tttgcgtatt	gggcgccagg	gtgggttttc	ttttcaccag	tgagacgggc	4800
aacagctgat	tgcccttcac	cgccggccc	tgagagagtt	gcagcaagcg	gtccacgctg	4860
gtttgcccc	gcaggcgaaa	atcctgtttg	atggtggtta	acggcgggat	ataacatgag	4920
ctgtcttcgc	tatcgtcgta	tcccactacc	gagatatccg	caccaacgcg	cagcccggac	4980
tcggtaattg	cgcgcatcgc	gcccagcgcc	atctgatcgt	tggcaaccag	catcgcatg	5040
ggaacgatgc	cctcattcag	catttgcatg	gtttgttgaa	aaccggacat	ggcactccag	5100
tcgccttccc	gttcgcgtat	cggtgaatt	tgattgcgag	tgagatattt	atgccagcca	5160
gccagacgca	gacgcgccga	gacagaactt	aatgggccc	ctaacagcgc	gatttgctgg	5220
tgacccaatg	cgaccagatg	ctccacgccc	agtcgcgtac	cgtcttcacg	ggagaaaata	5280
atactgttga	tgggtgtctg	gtcagagaca	tcaagaaata	acgcgggaac	attagtgcag	5340
gcagcttcca	cagcaatggc	atcctggtea	tccagcggat	agttaatgat	cagccactg	5400
acgcgttgcg	cgagaagatt	gtgcaccgcc	gctttacagg	cttcgacgcc	gcttcgttct	5460
accatcgaca	ccaccacgct	ggcaccaggt	tgatcggcgc	gagatttaat	cgccgcgaca	5520
atttgcgacg	gcgcgtgcag	ggccagactg	gaggtggcaa	cgccaatcag	caacgactgt	5580
ttgcccgcca	gttgttgtgc	cacgcggttg	ggaatgtaat	tcagctccgc	catcgccgct	5640
tccacttttt	cccgcgtttt	cgcagaaaac	tggtggcct	ggttcaccac	gcgggaaacg	5700
gtctgataag	agacaccggc	atactctgcg	acatcgtata	acgttactgg	tttcacattc	5760
accaccctga	attgactctc	ttccgggcgc	tatcatgcca	taccgcgaaa	ggttttgcgc	5820
cattcgatgg	tgtcgggat	ctcgacgctc	tcccttatgc	gactcctgca	ttaggaagca	5880
gccagtagt	aggttgaggc	cgttgagcac	cgccgccgca	aggaatgggtg	catg	5934

<210> 3  
<211> 6959  
<212> DNA  
<213> Artificial

<220>  
<223> pcDNA/Biotag-DEST -

<400> 3  
gacggatcgg gagatctccc gatccccctat ggtcgactct cagtacaatc tgctctgatg 60  
ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggtcgct gagtagtgcg 120  
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc 180  
ttagggttag gcgttttgcg ctgcttcgcg atgtacgggc cagatatacg cgttgacatt 240  
gattattgac tagttattaa tagtaatcaa ttacggggtc attagttcat agcccatata 300  
tggagttccg cgttacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc 360  
cccgccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc 420  
attgacgtca atgggtggac tatttacggg aaactgccca cttggcagta catcaagtgt 480  
atcatatgcc aagtacgcc cctattgacg tcaatgacgg taaatggccc gcctggcatt 540  
atgcccagta catgacctta tgggactttc ctacttggca gtacatctac gtattagtca 600  
tcgctattac catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggtttg 660  
actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc 720  
aaaatcaacg ggactttcca aaatgtcgta acaactccgc ccattgacg caaatgggcg 780  
gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact agagaacca 840  
ctgcttactg gcttatcgaa attaatacga ctactatag ggagacccaa gctggctagc 900  
gtttaaactt aagcttacca tgggcgcg cgccccgtg accgccccgc tggcgggcac 960  
tatctggaag gtgctggcca gcgaaggcca gacggtggcc gcaggcgagg tgctgctgat 1020  
tctggaagcc atgaagatgg aaaccgaaat ccgcgcgcgc caggccggga ccgtgcgcgc 1080  
tatcgcggtg aaagccggcg acgcggtggc ggtcggcgac accctgatga ccctggcggg 1140  
ctctggatcc gatctgtacg acgatgacga taaggtacat caaacaagtt tgtacaaaaa 1200  
agctgaacga gaaacgtaaa atgatataaa tatcaatata ttaaattaga ttttgcataa 1260  
aaaacagact acataatact gtaaaacaca acatatccag tcactatggc ggccgcatta 1320  
ggcaccaccg gctttacact ttatgcttcc ggctcgtata atgtgtggat tttgagttag 1380  
gatccggcga gattttcagg agctaaggaa gctaaaatgg agaaaaaat cactggatat 1440  
accaccgttg atatatccca atggcatcgt aaagaacatt ttgaggcatt tcagtcagtt 1500  
gctcaatgta cctataacca gaccgttcag ctggatatta cggccttttt aaagaccgta 1560  
aagaaaaata agcacaagtt ttatccggcc ttatttcaca ttcttgccc cctgatgaat 1620

gctcatccgg aattccgtat ggcaatgaaa gacggtgagc tggatgatg ggatagtgtt	1680
cacccttggt acaccgtttt ccatgagcaa actgaaacgt tttcatcgct ctggagtga	1740
taccacgacg atttccggca gtttctacac atatattcgc aagatgtggc gtgttacggt	1800
gaaaacctgg cctattttccc taaagggttt attgagaata tgtttttcgt ctcagccaat	1860
ccctgggtga gtttcaccag ttttgattta aacgtggcca atatggacaa cttcttcgcc	1920
cccgttttca ccatgggcaa atattatacg caaggcgaca aggtgctgat gccgctggcg	1980
attcaggttc atcatgccgt ctgtgatggc ttccatgtcg gcagaatgct taatgaatta	2040
caacagtact gcgatgagtg gcagggcggg gcgtaaacgc gtggatccgg cttactaaaa	2100
gccagataac agtatgcgta tttgcgcgct cgcgaaccgg tgtatacccg aagtatgtca	2160
aaaagagggt tgctatgaag cagcgtatta cagtacagt tgacagcgac agctatcagt	2220
tgctcaaggc atatatgatg tcaatatctc cggtctggta agcacaacca tgcagaatga	2280
agcccgctgt ctgcgtgccg aacgctggaa agcggaaaat caggaaggga tggctgaggt	2340
cgcccggttt attgaaatga acggctcttt tgctgacgag aacagggact ggtgaaatgc	2400
agtttaaggt ttacacctat aaaagagaga gccgttatcg tctgtttgtg gatgtacaga	2460
gtgatattat tgacacgccc gggcgacgga tggatgatccc cctggccagt gcacgtctgc	2520
tgtcagataa agtctcccgt gaactttacc cggtgggtgca tatcggggat gaaagctggc	2580
gcatgatgac caccgatatg gccagtgtgc cggtctccgt tatcggggaa gaagtggctg	2640
atctcagcca ccgcgaaaat gacatcaaaa acgccattaa cctgatgttc tggggaatat	2700
aaatgtcagg ctccgttata cacagccagt ctgcaggctc accatagtga ctggatatgt	2760
tgtgttttac agtattatgt agtctgtttt ttatgcaaaa tctaatttaa tatattgata	2820
tttatatcat ttacgtttc tcgttcagct ttcttgtaaa aagtgggtgat aattaattaa	2880
gatctagagg gcccgtttaa acccgctgat cagcctcgac tgtgccttct agttgccagc	2940
catctgttgt ttgcccctcc cccgtgcctt ccttgaccct ggaaggtgcc actcccactg	3000
tcctttccta ataaaatgag gaaattgcat cgcattgtct gagtaggtgt cattctattc	3060
tgggggggtgg ggtggggcag gacagcaagg gggaggattg ggaagacaat agcaggcatg	3120
ctgggggatgc ggtgggctct atggcttctg aggcggaaaag aaccagctgg ggctctaggg	3180
ggtatcccca cgcgccctgt agcggcgcat taagcgcggc ggggtgtggtg gttacgcgca	3240
gcgtgaccgc tacacttgcc agcgccctag cgcccgctcc tttcgcttcc ttccttcct	3300
ttctcgccac gttcgccggc tttccccgtc aagctctaaa tcggggcatc cctttagggt	3360
tccgatttag tgctttacgg cacctcgacc ccaaaaaact tgattagggt gatggttcac	3420
gtagtgggccc atcgccctga tagacggttt ttcgcccttt gacgttgag tccacgttct	3480

ttaatagtgg	actcttggtc	caaactggaa	caacactcaa	ccctatctcg	gtctattctt	3540
ttgatttata	agggattttg	gggatttcgg	cctattgggt	aaaaaatgag	ctgatttaac	3600
aaaaatttaa	cgcgaaattaa	ttctgtggaa	tgtgtgtcag	ttaggggtgtg	gaaagtcctc	3660
aggctcccca	ggcaggcaga	agtatgcaaa	gcatgcatct	caattagtca	gcaaccaggt	3720
gtggaaagtc	cccaggctcc	ccagcaggca	gaagtatgca	aagcatgcat	ctcaattagt	3780
cagcaaccat	agtcctcgcc	ctaactccgc	ccatcccgcc	cctaactccg	ccagttccg	3840
cccattctcc	gccccatggc	tgactaattt	tttttattta	tgcagaggcc	gaggccgcct	3900
ctgcctctga	gctattccag	aagtagtgag	gaggcttttt	tggaggccta	ggcttttgca	3960
aaaagctccc	gggagcttgt	atatccattt	tggatctga	tcagcacgtg	ttgacaatta	4020
atcatcggca	tagtatatcg	gcatagtata	atacgacaag	gtgaggaact	aaaccatggc	4080
caagcctttg	tctcaagaag	aatccaccct	cattgaaaga	gcaacggcta	caatcaacag	4140
catccccatc	tctgaagact	acagcgctgc	cagcgagct	ctctctagcg	acggccgcat	4200
cttcaactgg	gtcaatgtat	atcattttac	tgggggacct	tgtgcagaac	tcgtgggtgt	4260
gggcactgct	gctgctgcgg	cagctggcaa	cctgacttgt	atcgctcgga	tcggaaatga	4320
gaacaggggc	atcttgagcc	cctgcggacg	gtgccgacag	gtgcttctcg	atctgcatcc	4380
tgggatcaaa	gccatagtga	aggacagtga	tggacagccg	acggcagttg	ggattcgtga	4440
attgctgccc	tctggttatg	tgtgggaggg	ctaagcactt	cgtggccgag	gagcaggact	4500
gacacgtgct	acgagatttc	gattccaccg	ccgccttcta	tgaaagggtg	ggcttcggaa	4560
tcgttttccg	ggacgcgggc	tggatgatcc	tccagcgcg	ggatctcatg	ctggagttct	4620
tcgcccaccc	caacttgttt	attgcagctt	ataatggtta	caaataaagc	aatagcatca	4680
caaatttcac	aaataaagca	tttttttcac	tgcattctag	ttgtggtttg	tccaaactca	4740
tcaatgtatc	ttatcatgtc	tgtataccgt	cgacctctag	ctagagcttg	gcgtaatcat	4800
ggtcatagct	gtttcctgtg	tgaaattggt	atccgctcac	aattccacac	aacatacgag	4860
ccggaagcat	aaagtgtaaa	gcctgggggtg	cctaatagag	gagctaactc	acattaattg	4920
cgttgcgctc	actgcccgt	ttccagtcgg	gaaacctgtc	gtgccagctg	cattaatgaa	4980
tcggccaacg	cgcggggaga	ggcggtttgc	gtattgggcg	ctcttcgct	tcctcgctca	5040
ctgactcgct	gcgctcggtc	gttcggctgc	ggcgagcggt	atcagctcac	tcaaaggcgg	5100
taatacggtt	atccacagaa	tcaggggata	acgcaggaaa	gaacatgtga	gcaaaaggcc	5160
agcaaaaggc	caggaaccgt	aaaaaggccg	cgttgctggc	gtttttccat	aggctccgcc	5220
cccctgacga	gcatcacaaa	aatcgacgct	caagtcagag	gtggcgaaac	ccgacaggac	5280
tataaagata	ccaggcggtt	ccccctggaa	gtccctcgt	gcgctctcct	gttccgaccc	5340

```

tgcgcgttac cggatacctg tccgcctttc tcccttcggg aagcgtggcg ctttctcaat 5400
gtcacgctg taggtatctc agttcgggtg aggtcggtcg ctccaagctg ggctgtgtgc 5460
acgaaccccc cggttcagccc gaccgctgcg ccttatccgg taactatcgt cttgagtcca 5520
acccggtaag acacgactta tcgccactgg cagcagccac tggtaacagg attagcagag 5580
cgaggatatgt aggcgggtgct acagagttct tgaagtgggt gcctaactac ggctacacta 5640
gaaggacagt atttgggtatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg 5700
gtagctcttg atccggcaaa caaaccaccg ctggtagcgg tggttttttt gtttgcaagc 5760
agcagattac gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt tctacggggt 5820
ctgacgctca gtggaacgaa aactcacgtt aagggatttt ggtcatgaga ttatcaaaaa 5880
ggatcttcac ctagatcctt ttaaattaaa aatgaagttt taaatcaatc taaagtatat 5940
atgagtaaac ttgggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga 6000
tctgtctatt tcgttcatcc atagttgcct gactccccgt cgtgtagata actacgatac 6060
gggaggggctt accatctggc ccagtgctg caatgatacc gcgagacca cgctcaccgg 6120
ctccagattt atcagcaata aaccagccag ccggaagggc cgagcgaga agtggctctg 6180
caactttatc cgctccatc cagtctatta attggtgccg ggaagctaga gtaagtagtt 6240
cgccagttaa tagtttgcc aacgttggtt ccattgtac aggcacgtg gtgtcacgt 6300
cgtcgtttgg tatggcttca ttcagctccg gtcccaacg atcaaggcga gttacatgat 6360
cccccatggt gtgcaaaaaa gcggttagct ccttcgggtc tccgatcgtt gtcagaagta 6420
agttggccgc agtgttatca ctcatgggta tggcagcact gcataattct cttactgtca 6480
tgccatccgt aagatgcttt tctgtgactg gtgagtactc aaccaagtca ttctgagaat 6540
agtgtatgcg gcgaccgagt tgctcttgcc cggcgtaaat acgggataat accgcgccac 6600
atagcagaac tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga aaactctcaa 6660
ggatcttacc gctgttgaga tccagttcga tgtaaccac tcgtgcaccc aactgatctt 6720
cagcatcttt tactttcacc agcgtttctg ggtgagcaaa aacaggaagg caaatgccg 6780
caaaaaaggg aataagggcg acacggaaat gttgaatact catactcttc ctttttcaat 6840
attattgaag catttatcag ggttattgtc tcatgagcgg atacatattt gaatgtattt 6900
agaaaaataa acaaataggg gttccgcgca catttccccg aaaagtgcc cctgacgtc 6959

```

<210> 4

<211> 5302

<212> DNA

<213> Artificial

<220>

<223> pcDNA6/Biotag/D-TOPO



<400> 4  
gacggatcgg gagatctccc gatcccctat ggtcgaactct cagtacaatc tgctctgatg 60  
ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggtcgct gagtagtgcg 120  
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc 180  
ttaggggttag gcgttttgcg ctgcttcgcg atgtacgggc cagatatacg cgttgacatt 240  
gattattgac tagttattaa tagtaatcaa ttacggggtc attagttcat agcccatata 300  
tggagttccg cgttacataa cttacggtaa atggcccgcg tggctgaccg cccaacgacc 360  
ccgcccatt gacgtcaata atgacgtatg tcccatagt aacgccaata gggactttcc 420  
attgacgtca atgggtggac tatttacggg aaactgccca cttggcagta catcaagtgt 480  
atcatatgcc aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt 540  
atgccagta catgacctta tgggactttc ctacttgga gtacatctac gtattagtca 600  
tcgtattac catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggtttg 660  
actcacgggg atttccaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc 720  
aaaatcaacg ggactttcca aaatgtcgta acaactcgc cccattgacg caaatgggcg 780  
gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact agagaacca 840  
ctgcttactg gcttatcgaa attaatacga ctactatag ggagacccaa gctggctagc 900  
gtttaaactt aagcttacca tgggcgcggg caccgccgtg accgcccgc tggcgggcac 960  
tatctggaag gtgctggcca gcgaaggcca gacggtggcc gcaggcgagg tgctgctgat 1020  
tctggaagcc atgaagatgg aaaccgaaat ccgcgcgcg caggccggga ccgtgcgcg 1080  
tatcgcggtg aaagccggcg acgcggtggc ggtcggcgac accctgatga ccctggcggg 1140  
ctctggatcc gatctgtacg acgatgacga taaggtaact aggatccagt gtggtggaat 1200  
tgatcccttc accaagggcg tcgagtctag agggcccgtt taaaccgct gatcagcctc 1260  
gactgtgcct tctagttgcc agccatctgt tgtttgcccc tccccgtgc ctcccttgac 1320  
cctggaaggt gccactccca ctgtcctttc ctaataaaat gaggaattg catcgcattg 1380  
tctgagtagg tgtcattcta ttctgggggg tggggtgggg caggacagca agggggagga 1440  
ttgggaagac aatagcaggc atgctgggga tgcggtgggc tctatggctt ctgaggcgga 1500  
aagaaccagc tggggctcta gggggtatcc ccacgcgcc tgtagcggcg cattaagcgc 1560  
ggcgggtgtg gtggttacgc gcagcgtgac cgtacactt gccagcgcgc tagcgcgcgc 1620  
tcctttcgct ttcttccctt cctttctcgc caggttcgcc ggctttcccc gtcaagctct 1680  
aaatcggggc atcccttttag ggttccgatt tagtgcttta cggcacctcg accccaaaaa 1740  
acttgattag ggtgatggtt cacgtagtgg gccatcgccc tgatagacgg tttttcgccc 1800  
tttgacgttg gagtccacgt tctttaatag tggactcttg ttccaaactg gaacaacact 1860

caaccctatc tcggtctatt cttttgattt ataagggatt ttggggattt cggcctattg	1920
gttaaaaaat gagctgattt aacaaaaatt taacgcgaat taattctgtg gaatgtgtgt	1980
cagttagggt gtggaaagtc cccaggctcc ccaggcaggc agaagtatgc aaagcatgca	2040
tctcaattag tcagcaacca ggtgtggaaa gtccccaggc tccccagcag gcagaagtat	2100
gcaaagcatg catctcaatt agtcagcaac catagtcccc cccctaactc cgcccatccc	2160
gcccctaact ccgcccagtt ccgcccattc tccgccccat ggctgactaa ttttttttat	2220
ttatgcagag gccgaggccg cctctgcctc tgagctattc cagaagtagt gaggaggctt	2280
ttttggaggc ctaggctttt gcaaaaagct cccgggagct tgtatatcca ttttcggatc	2340
tgatcagcac gtgttgacaa ttaatcatcg gcatagtata tcggcatagt ataatacgac	2400
aagggtgagga actaaaccat ggccaagcct ttgtctcaag aagaatccac cctcattgaa	2460
agagcaacgg ctacaatcaa cagcatcccc atctctgaag actacagcgt cgccagcgca	2520
gctctctcta gcgacggccg catcttcact ggtgtcaatg tatatcattt tactggggga	2580
ccttgtgcag aactcgtggt gctgggcact gctgctgctg cggcagctgg caacctgact	2640
tgtatcgtcg cgatcggaaa tgagaacagg ggcattctga gcccctgcgg acggtgccga	2700
cagggtgctt tcgatctgca tcctgggatc aaagccatag tgaaggacag tgatggacag	2760
ccgacggcag ttgggattcg tgaattgctg ccctctgggt atgtgtggga gggctaagca	2820
cttcgtggcc gaggagcagg actgacacgt gctacgagat ttcgattcca ccgccgcctt	2880
ctatgaaagg ttgggcttcg gaatcgtttt ccgggacgcc ggctggatga tcctccagcg	2940
cggggatctc atgctggagt tcttcgcca cccaacttg tttattgcag cttataatgg	3000
ttacaaataa agcaatagca tcacaaattt cacaaataaa gcattttttt cactgcattc	3060
tagttgtggt ttgtccaaac tcatcaatgt atcttatcat gtctgtatac cgtcgacctc	3120
tagctagagc ttggcgtaat catggtcata gctgtttcct gtgtgaaatt gttatccgct	3180
cacaattcca cacaacatac gagccggaag cataaagtgt aaagcctggg gtgcctaattg	3240
agtgcgctaa ctcacattaa ttgcgttgcg ctactgccc gcttttcagt cgggaaacct	3300
gtcgtgccag ctgcattaat gaatcggcc aacgcggggg agaggcggtt tgcgtattgg	3360
gcgctcttcc gcttcctcgc tcaactgact gctgcgctcg gtcgttcggc tgcggcgagc	3420
ggtatcagct cactcaaagg cggtaatacg gttatccaca gaatcagggg ataacgcagg	3480
aaagaacatg tgagcaaaaag gccagcaaaa ggccaggaac cgtaaaaagg ccgcgttgct	3540
ggcgtttttc cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca	3600
gagggtggcg aacccgacag gactataaag ataccaggcg tttccccctg gaagctccct	3660
cgtgcgctct cctgttccga ccctgccgct taccggatac ctgtccgcct ttctcccttc	3720

gggaagcgtg gcgctttctc aatgctcacg ctgtaggtat ctcagttcgg tgtaggtcgt 3780  
tcgctccaag ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct gcgccttctc 3840  
cggtaactat cgtcttgagt ccaaccgggt aagacacgac ttatcgccac tggcagcagc 3900  
cactggtaac aggattagca gagcgaggta tgtaggcggt gctacagagt tcttgaagtg 3960  
gtggcctaac tacggctaca ctagaaggac agtatttggt atctgcgctc tgctgaagcc 4020  
agttaccttc ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag 4080  
cgggtggtttt tttgtttgca agcagcagat tacgcgcaga aaaaaaggat ctcaagaaga 4140  
tcctttgctc ttttctacgg ggtctgacgc tcagtggaaac gaaaactcac gttaagggat 4200  
tttggctcatg agattatcaa aaaggatctt cacctagatc cttttaaatt aaaaatgaag 4260  
ttttaaatca atctaaagta tatatgagta aacttggtct gacagttacc aatgcttaat 4320  
cagtgagga cctatctcag cgatctgtct atttcgttca tccatagttg cctgactccc 4380  
cgctcgtgtg ataactacga tacgggaggg cttaccatct ggccccagtg ctgcaatgat 4440  
accgcgagac ccacgctcac cggtccaga tttatcagca ataaaccagc cagccggaag 4500  
ggccgagcgc agaagtggtc ctgcaacttt atccgcctcc atccagtcta ttaattgttg 4560  
ccgggaagct agagtaagta gttcgccagt taatagtttg cgcaacgttg ttgccattgc 4620  
tacaggcctc gtggtgtcac gctcgtcgtt tggtaggct tcattcagct ccggttccca 4680  
acgatcaagg cgagttacat gatcccccat gttgtgcaaa aaagcgggta gtccttcgg 4740  
tcctccgctc gttgtcagaa gtaagttggc cgcagtggtta tcaactcatgg ttatggcagc 4800  
actgcataat tctcttactg tcatgccatc cgtaagatgc ttttctgtga ctggtgagta 4860  
ctcaaccaag tcattctgag aatagtgtat gcggcgaccg agttgctctt gcccgcgctc 4920  
aatacgggat aataccgcgc cacatagcag aactttaaaa gtgctcatca ttggaaaacg 4980  
ttcttcgggg cgaaaactct caaggatctt accgctgttg agatccagtt cgatgtaacc 5040  
cactcgtgca cccaactgat cttcagcctc ttttactttc accagcggtt ctgggtgagc 5100  
aaaaacagga aggcaaaatg ccgcaaaaaa gggaataagg gcgacacgga aatgttgaat 5160  
actcactctc ttcctttttc aatattattg aagcatttat cagggttatt gtctcatgag 5220  
cggatacata tttgaatgta tttagaaaaa taaacaaata ggggttccgc gcacatttcc 5280  
ccgaaaagtg ccacctgacg tc 5302

<210> 5  
<211> 5375  
<212> DNA  
<213> Artificial  
  
<220>  
<223> PMT/Biotag-DEST

<400> 5  
tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccc gagacggtca 60  
cagcttgtct gtaagcggat gccgggagca gacaagcccc tcagggcgcg tcagcgggtg 120  
ttggcgggtg tcggggctgg cttaactatg cggtatcaga gcagattgta ctgagagtgc 180  
accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc 240  
attcgccatt caggctgcgc aactgttggg aagggcgatc ggtgcgggcc tcttcgctat 300  
tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagttgggta acgccagggt 360  
tttcccagtc acgacgttgt aaaacgacgg ccagtgccag tgaattaatt cgttgcagga 420  
caggatgtgg tgcccgatgt gactagctct ttgctgcagg ccgtcctatc ctctggttcc 480  
gataagagac ccagaactcc ggccccccac cgcccaccgc ccccccata catatgtggt 540  
acgcaagtaa gattgcctgc gcatgcccc tgtgccccac caagagtttt gcatcccata 600  
caagtcccca aagtggagaa ccgaaccaat tcttcgcggg cagaacaaaa gcttctgcac 660  
acgtctccac tcgaatttgg agccggccgg cggtgtgcaa agaggtgaat cgaacgaaag 720  
acccgtgtgt aaagccgcgt ttccaaaatg tataaaaccg agagcatctg gccaatgtgc 780  
atcagttgtg gtcagcagca aaatcaagt aatcatctca gtgcaactaa aggggggatc 840  
tagcgtttaa acttaagctt accatgggcg ccggcacccc ggtgaccgcc ccgctggcgg 900  
gcactatctg gaagggtgct gccagcgaag gccagacggg ggccgcaggc gaggtgctgc 960  
tgattctgga agccatgaag atggaaaccg aaatccgcgc cgcgcaggcc gggaccgtgc 1020  
gcggtatcgc ggtgaaagcc ggcgacgcgg tggcggtcgg cgacaccctg atgaccctgg 1080  
cgggctctgg atccgatctg tacgacgatg acgataaggc acatcaaaca agtttgtaca 1140  
aaaaagctga acgagaaacg taaaatgata taaatatcaa tatattaaat tagattttgc 1200  
ataaaaaaca gactacataa tactgtaaaa cacaacatat ccagtcacta tggcgccgcg 1260  
attaggcacc ccaggcttta cactttatgc ttccggctcg tataatgtgt ggattttgag 1320  
ttaggatccg gcgagatttt caggagctaa ggaagctaaa atggagaaaa aaatcactgg 1380  
atataccacc gttgatatat cccaatggca tcgtaaagaa cattttgagg catttcagtc 1440  
agttgctcaa tgtacctata accagaccgt tcagctggat attacggcct ttttaaagac 1500  
cgtaaagaaa aataagcaca agttttatcc ggcctttatt cacattcttg cccgcctgat 1560  
gaatgctcat ccggaattcc gtatggcaat gaaagacggg gagctggtga tatgggatag 1620  
tgttcaccct tgttacaccg ttttccatga gcaaactgaa acgttttcat cgctctggag 1680  
tgaataccac gacgatttcc ggagtttct acacatatat tcgcaagatg tggcgtgtta 1740  
cggtgaaaac ctggcctatt tccctaaagg gtttattgag aatatgtttt tcgtctcagc 1800  
caatccctgg gtgagtttca ccagttttga tttaaacgtg gccaatatgg acaacttctt 1860

cgcccccggtt ttcacccatgg gcaaataatta tacgcaaggc gacaagggtgc tgatgccgct 1920  
ggcgattcag gttcatcatg ccgctctgtga tggcttccat gtcggcagaa tgcttaata 1980  
attacaacag tactgcgatg agtggcaggg cggggcgtaa acgcgtggat ccggcttact 2040  
aaaagccaga taacagtatg cgtatctgcg cgctcgcgaa ccggtgtata cccgaagtat 2100  
gtcaaaaaga ggtgtgctat gaagcagcgt attacagtga cagttgacag cgacagctat 2160  
cagttgctca aggcatatat gatgtcaata tctccggtct ggtaagcaca accatgcaga 2220  
atgaagcccg tcgtctgcgt gccgaacgct ggaaagcggg aaatcaggaa gggatggctg 2280  
aggctgccccg gtttattgaa atgaacggct cttttgctga cgagaacagg gactggtgaa 2340  
atgcagttta aggtttacac ctataaaaaga gagagccggt atcgtctgtt tgtggatgta 2400  
cagagtata ttattgacac gcccgggcga cggatgggtga tccccctggc cagtgcacgt 2460  
ctgctgtcag ataaagtctc ccggtgaactt taccgggtgg tgcataatcg ggatgaaagc 2520  
tggcgcata tgaccaccga tatggccagt gtgccggtct ccgttatcgg ggaagaagtg 2580  
gctgatctca gccaccgcga aaatgacatc aaaaacgcca ttaacctgat gttctgggga 2640  
atataaatgt caggctccgt tatacacagc cagtctgcag gtcgaccata gtgactggat 2700  
atgttgtgtt ttacagtatt atgtagtctg ttttttatgc aaaatctaatt ttaatatatt 2760  
gatatttata tcattttacg tttctcgttc agctttcttg tacaaagtgg tgataattaa 2820  
ttaagatcta gagggccggt ttaaaccgcg tgatcagcct cgactgtgcc ttctaagatc 2880  
cagacatgat aagatacatt gatgagtttg gacaaaccac aactagaatg cagtgaaaaa 2940  
aatgctttat ttgtgaaatt tgtgatgcta ttgctttatt tgtaaccatt ataagctgca 3000  
ataaacaagt taacaacaac aattgcattc attttatgtt tcagggttcag ggggaggtgt 3060  
gggaggtttt ttaaagcaag taaaacctct acaaagtggg tatggctgat tatgatcagt 3120  
cgacctgcag gcatgcaagc ttggcgtaat catggtcata gctgtttcct gtgtgaaatt 3180  
gttatccgct cacaattcca cacaacatac gagccggaag cataaagtgt aaagcctggg 3240  
gtgcctaata agtgagctaa ctcacattaa ttgcgttgcg ctactgccc gctttccagt 3300  
cgggaaacct gtcgtgccag ctgcattaat gaatcgcca acgcgcgggg agaggcggtt 3360  
tgcgatttgg gcgctcttcc gcttctctgc tctactgactc gctgcgctcg gtcgttcggc 3420  
tgcggcgagc ggtatcagct cactcaaagg cggtaatacg gttatccaca gaatcagggg 3480  
ataacgcagg aaagaacatg tgagcaaaaag gccagcaaaa ggccaggaaac cgtaaaaagg 3540  
ccgcgttgct ggcgtttttc cataggctcc gccccctga cgagcatcac aaaaatcgac 3600  
gctcaagtca gaggtggcga aaccgcagag gactataaag ataccaggcg tttccccctg 3660  
gaagctccct cgtgcgctct cctgttccga cctgcccgt taccggatac ctgtccgcct 3720

ttctcccttc	gggaagcgtg	gcgctttctc	atagctcacg	ctgtaggtat	ctcagttcgg	3780
tgtaggtcgt	tcgctccaag	ctgggctgtg	tgcacgaacc	ccccgttcag	cccgaccgct	3840
gcgccttata	cggtaactat	cgtcttgagt	ccaacccggt	aagacacgac	ttatcgccac	3900
tggcagcagc	cactggtaac	aggattagca	gagcgaggta	tgtaggcggt	gctacagagt	3960
tcttgaagtg	gtggcctaac	tacggctaca	ctagaaggac	agtatttggt	atctgcgctc	4020
tgctgaagcc	agttaccttc	ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	4080
ccgctggtag	cggtggtttt	tttgtttgca	agcagcagat	tacgcgcaga	aaaaaaggat	4140
ctcaagaaga	tcctttgata	ttttctacgg	ggtctgacgc	tcagtggaac	gaaaactcac	4200
gttaagggat	tttggtcattg	agattatcaa	aaaggatctt	cacctagata	cttttaaatt	4260
aaaaatgaag	ttttaaatca	atctaaagta	tatatgagta	aacttggtct	gacagttacc	4320
aatgcttaat	cagtgaggca	cctatctcag	cgatctgtct	atttcgttca	tccatagttg	4380
cctgactccc	cgctcgttag	ataactacga	tacgggaggg	cttaccatct	ggccccagtg	4440
ctgcaatgat	accgcgagac	ccacgctcac	cggtccaga	tttatcagca	ataaaccagc	4500
cagccggaag	ggccgagcgc	agaagtggtc	ctgcaacttt	atccgcctcc	atccagtcta	4560
ttaattgttg	ccgggaagct	agagtaagta	gttcgccagt	taatagtttg	cgcaacgttg	4620
ttgccattgc	tacaggcata	gtggtgtcac	gctcgtcgtt	tggtagggct	tcattcagct	4680
ccggttccca	acgatcaagg	cgagttacat	gatccccat	gttgtgcaaa	aaagcgggta	4740
gctccttcgg	tcctccgata	gttgtcagaa	gtaagttggc	cgcagtgtta	tcactcatgg	4800
ttatggcagc	actgcataat	tctcttactg	tcatgccata	cgtaagatgc	ttttctgtga	4860
ctggtgagta	ctcaaccaag	tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	4920
gccccgcgtc	aatacgggat	aataccgcgc	cacatagcag	aactttaaaa	gtgctcatca	4980
ttggaaaacg	ttcttcgggg	cgaaaactct	caaggatctt	accgctgttg	agatccagtt	5040
cgatgtaacc	cactcgtgca	cccaactgat	cttcagcata	ttttactttc	accagcgttt	5100
ctgggtgagc	aaaaacagga	aggcaaatg	ccgcaaaaaa	gggaataagg	gcgacacgga	5160
aatgttgaat	actcatactc	ttcctttttc	aatattattg	aagcatttat	caggggttatt	5220
gtctcatgag	cggatacata	tttgaatgta	tttagaaaaa	taaacaaata	ggggttccgc	5280
gcacatttcc	ccgaaaagtg	ccacctgacg	tctaagaaac	cattattata	atgacattaa	5340
cctataaaaa	taggcgtata	acgaggcctt	ttcgt			5375

<210> 6

<211> 72

<212> PRT

<213> Klebsiella pneumoniae

<400> 6

Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp Lys  
1 5 10 15

Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu Leu  
20 25 30

Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln Ala  
35 40 45

Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala Val  
50 55 60

Gly Asp Thr Leu Met Thr Leu Ala  
65 70

<210> 7

<211> 115

<212> PRT

<213> Mus musculus

<400> 7

Lys Ala Leu Ala Val Ser Asp Leu Asn Arg Ala Gly Gln Arg Gln Val  
1 5 10 15

Phe Phe Glu Leu Asn Gly Gln Leu Arg Ser Ile Leu Val Lys Asp Thr  
20 25 30

Gln Ala Met Lys Glu Met His Phe His Pro Lys Ala Leu Lys Asp Val  
35 40 45

Lys Gly Gln Ile Gly Ala Pro Met Pro Gly Lys Val Ile Asp Ile Lys  
50 55 60

Val Ala Ala Gly Asp Lys Val Ala Lys Gly Gln Pro Leu Cys Val Leu  
65 70 75 80

Ser Ala Met Lys Met Glu Thr Val Val Thr Ser Pro Met Glu Gly Thr  
85 90 95

Ile Arg Lys Val His Val Thr Lys Asp Met Thr Leu Glu Gly Asp Asp  
100 105 110

Leu Ile Leu  
115

<210> 8

<211> 123  
 <212> PRT  
 <213> Propionibacterium shermanii

<400> 8

Met Lys Leu Lys Val Thr Val Asn Gly Thr Ala Tyr Asp Val Asp Val  
 1 5 10 15

Asp Val Asp Lys Ser His Glu Asn Pro Met Gly Thr Ile Leu Phe Gly  
 20 25 30

Gly Gly Thr Gly Gly Ala Pro Ala Pro Arg Ala Ala Gly Gly Ala Gly  
 35 40 45

Ala Gly Lys Ala Gly Glu Gly Glu Ile Pro Ala Pro Leu Ala Gly Thr  
 50 55 60

Val Ser Lys Ile Leu Val Lys Glu Gly Asp Thr Val Lys Ala Gly Gln  
 65 70 75 80

Thr Val Leu Val Leu Glu Ala Met Lys Met Glu Thr Glu Ile Asn Ala  
 85 90 95

Pro Thr Asp Gly Lys Val Glu Lys Val Leu Val Lys Glu Arg Asp Ala  
 100 105 110

Val Gln Gly Gly Gln Gly Leu Ile Lys Ile Gly  
 115 120

<210> 9  
 <211> 122  
 <212> PRT  
 <213> Homo sapiens

<400> 9

Gly Ser Cys Val Glu Val Asp Val His Arg Leu Ser Asp Gly Gly Leu  
 1 5 10 15

Leu Leu Ser Tyr Asp Gly Ser Ser Tyr Thr Thr Tyr Met Lys Glu Glu  
 20 25 30

Val Asp Arg Tyr Arg Ile Thr Ile Gly Asn Lys Thr Cys Val Phe Glu  
 35 40 45

Lys Glu Asn Asp Pro Ser Val Met Arg Ser Pro Ser Ala Gly Lys Leu  
 50 55 60

Ile Gln Tyr Ile Val Glu Asp Gly Gly His Val Phe Ala Gly Gln Cys



65		70		75		80									
Tyr	Ala	Glu	Ile	Glu	Val	Met	Lys	Met	Val	Met	Thr	Leu	Thr	Ala	Val
				85					90					95	
Glu	Ser	Gly	Cys	Ile	His	Tyr	Val	Lys	Arg	Pro	Gly	Ala	Ala	Leu	Asp
			100					105					110		
Pro	Gly	Cys	Val	Leu	Ala	Lys	Met	Gln	Leu						
		115					120								
<210>	10														
<211>	156														
<212>	PRT														
<213>	Escherichia coli														
<400>	10														
Met	Asp	Ile	Arg	Lys	Ile	Lys	Lys	Leu	Ile	Glu	Leu	Val	Glu	Glu	Ser
1				5					10					15	
Gly	Ile	Ser	Glu	Leu	Glu	Ile	Ser	Glu	Gly	Glu	Glu	Ser	Val	Arg	Ile
			20					25					30		
Ser	Arg	Ala	Ala	Pro	Ala	Ala	Ser	Phe	Pro	Val	Met	Gln	Gln	Ala	Tyr
		35					40					45			
Ala	Ala	Pro	Met	Met	Gln	Gln	Pro	Ala	Gln	Ser	Asn	Ala	Ala	Ala	Pro
	50					55					60				
Ala	Thr	Val	Pro	Ser	Met	Glu	Ala	Pro	Ala	Ala	Ala	Glu	Ile	Ser	Gly
65					70				75					80	
His	Ile	Val	Arg	Ser	Pro	Met	Val	Gly	Thr	Phe	Tyr	Arg	Thr	Pro	Ser
			85						90					95	
Pro	Asp	Ala	Lys	Ala	Phe	Ile	Glu	Val	Gly	Gln	Lys	Val	Asn	Val	Gly
			100					105					110		
Asp	Thr	Leu	Cys	Ile	Val	Glu	Ala	Met	Lys	Met	Met	Asn	Gln	Ile	Glu
		115					120					125			
Ala	Asp	Lys	Ser	Gly	Thr	Val	Lys	Ala	Ile	Leu	Val	Glu	Ser	Gly	Gln
	130					135					140				
Pro	Val	Glu	Phe	Asp	Glu	Pro	Leu	Val	Val	Ile	Glu				
145					150					155					

<210> 11  
<211> 216  
<212> DNA  
<213> *Klebsiella pneumoniae*

<400> 11  
ggcgccggca ccccggtgac cgccccgctg gcgggcacta tctggaaggt gctggccagc 60  
gaaggccaga cgggtggccgc aggcgaggtg ctgctgattc tggaagccat gaagatggaa 120  
accgaaatcc gcgccgcgca ggccgggacc gtgcgcggta tcgcggtgaa agccggcgac 180  
gcggtggcgg tcggcgacac cctgatgacc ctggcg 216

<210> 12  
<211> 345  
<212> DNA  
<213> *Mus musculus*

<400> 12  
aaagccctgg ctgtaagcga cctgaaccgt gctggccaga ggcaggtgtt ctttgaactc 60  
aatgggcagc ttcgatccat tctggttaaa gacaccagg ccatgaagga gatgcacttc 120  
catcccaagg ctttgaagga tgtgaagggc caaattgggg ccccgatgcc tgggaaggtc 180  
atagacatca aggtggcagc aggggacaag gtggctaagg gccagcccct ctgtgtgctc 240  
agcgccatga agatggagac tgtggtgact tcgccatgg agggcactat ccgaaagggt 300  
catgttacca aggacatgac tctggaaggc gacgacctca tccta 345

<210> 13  
<211> 369  
<212> DNA  
<213> *Propionibacterium shermanii*

<400> 13  
atgaaactga aggtaacagt caacggcact gcgtatgacg ttgacgttga cgtcgacaag 60  
tcacacgaaa acccgatggg caccatcctg ttcggcggcg gcaccggcgg cgcgccggca 120  
ccgcgcgcag caggtggcgc aggcgccggt aaggccggag agggcgagat tcccgtccg 180  
ctggccggca ccgtctccaa gatcctcgtg aaggaggggt acacgggtcaa ggctggtcag 240  
accgtgctcg ttctcgaggc catgaagatg gagaccgaga tcaacgctcc caccgacggc 300  
aaggtcgaga aggtccttgt caaggagcgt gacgccgtgc agggcgggtca gggctctcatc 360  
aagatcggc 369

<210> 14  
<211> 366  
<212> DNA  
<213> *Homo sapiens*

<400> 14  
ggctcatgtg tagaagtaga tgtacatcgg ctgagtgacg gtggactgct cttgtcctat 60

gatggcagca gttacaccac gtatatgaag gaggaagtag acagatatcg catcacaatt	120
ggcaataaaaa cctgtgtgtt tgagaaggaa aatgacccat cggatgatgcg ctcaccttct	180
gctgggaagt taatccagta cattgtagaa gatggaggtc atgtgtttgc cggccagtgc	240
tatgcagaga ttgaggtaat gaagatggta atgactttga cagctgtgga gtctggctgt	300
atccattacg tcaagcgtcc tggagcagct cttgaccctg gctgtgtact cgccaaaatg	360
caactg	366

<210> 15  
 <211> 468  
 <212> DNA  
 <213> Escherichia coli

<400> 15	
atggatattc gtaagattaa aaaactgac gagctggttg aagaatcagg catctccgaa	60
ctggaaattt ctgaaggcga agagtcagta cgcattagcc gtgcagctcc tgccgcaagt	120
ttccctgtga tgcaacaagc ttacgctgca ccaatgatgc agcagccagc tcaatctaac	180
gcagccgctc cggcgaccgt tccttccatg gaagcgccag cagcagcgga aatcagtggg	240
cacatcgtag gttccccgat ggttggtagt ttctaccgca cccaagccc ggacgcaaaa	300
gcgttcatcg aagtgggtca gaaagtcaac gtgggcgata ccctgtgcat cgttgaagcc	360
atgaaaatga tgaaccagat cgaagcggac aaatccggtg ccgtgaaagc aattctggtc	420
gaaagtggac aaccggtaga atttgacgag ccgctggctg tcatcgag	468

<210> 16  
 <211> 8  
 <212> PRT  
 <213> Artificial

<220>  
 <223> FLAG epitope

<400> 16

Asp Tyr Lys Asp Asp Asp Asp Lys  
 1 5

<210> 17  
 <211> 8  
 <212> PRT  
 <213> Artificial

<220>  
 <223> FLAG epitope

<400> 17

Asp Tyr Lys Asp Glu Asp Asp Lys  
 1 5

<210> 18  
<211> 9  
<212> PRT  
<213> Artificial

<220>  
<223> Strep epitope

<400> 18

Ala Trp Arg His Pro Gln Phe Gly Gly  
1 5

<210> 19  
<211> 11  
<212> PRT  
<213> Artificial

<220>  
<223> VSV-G epitope

<400> 19

Tyr Thr Asp Ile Glu Met Asn Arg Leu Gly Lys  
1 5 10

<210> 20  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> poly-His epitope

<400> 20

His His His His His His  
1 5

<210> 21  
<211> 13  
<212> PRT  
<213> Artificial

<220>  
<223> Influenza epitope

<400> 21

Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ile Glu Gly Arg  
1 5 10

<210> 22  
<211> 11  
<212> PRT  
<213> Artificial

<220>  
<223> Human c-myc epitope  
<400> 22

Glu Gln Lys Leu Leu Ser Glu Glu Asp Leu Asn  
1 5 10

<210> 23  
<211> 3  
<212> PRT  
<213> Artificial

<220>  
<223> tripeptide epitope  
<400> 23

Glu Glu Phe  
1

<210> 24  
<211> 5  
<212> PRT  
<213> Artificial

<220>  
<223> enterokinase (EK) recognition site  
<400> 24

Asp Asp Asp Asp Lys  
1 5

<210> 25  
<211> 467  
<212> DNA  
<213> Artificial

<220>  
<223> pET104-DEST vector

<220>  
<221> CDS  
<222> (177)..(464)

<220>  
<221> misc\_feature  
<222> (466)..(467)  
<223> n is a, c, g, or t

<400> 25  
ataggcgcca gcaaccgcac ctgtggcgcc ggtgatgccg gccacgatgc gtccggcgta 60  
gaggatcgag atctcgatcc cgcgaaatta atacgactca ctatagggga attgtgagcg 120  
gataacaatt cccctctaga aataattttg tttaacttta agaaggagat atacat atg 179  
Met

1

ggc gcc ggc acc ccg gtg acc gcc ccg ctg gcg ggc act atc tgg aag 227  
Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp Lys  
5 10 15

gtg ctg gcc agc gaa ggc cag acg gtg gcc gca ggc gag gtg ctg ctg 275  
Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu Leu  
20 25 30

att ctg gaa gcc atg aag atg gaa acc gaa atc cgc gcc gcg cag gcc 323  
Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln Ala  
35 40 45

ggg acc gtg cgc ggt atc gcg gtg aaa gcc ggc gac gcg gtg gcg gtc 371  
Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala Val  
50 55 60 65

ggc gac acc ctg atg acc ctg gcg ggc tct gga tcc gat ctg tac gac 419  
Gly Asp Thr Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr Asp  
70 75 80

gat gac gat aag gga att atc aca agt ttg tac aaa aaa gca ggc tnn 467  
Asp Asp Asp Lys Gly Ile Ile Thr Ser Leu Tyr Lys Lys Ala Gly  
85 90 95

<210> 26

<211> 96

<212> PRT

<213> Artificial

<220>

<223> pET104-DEST vector .

<400> 26

Met Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp  
1 5 10 15

Lys Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu  
20 25 30

Leu Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln  
35 40 45

Ala Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala  
50 55 60

Val Gly Asp Thr Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr  
65 70 75 80

Asp Asp Asp Asp Lys Gly Ile Ile Thr Ser Leu Tyr Lys Lys Ala Gly  
85 90 95

<210> 27  
 <211> 449  
 <212> DNA  
 <213> Artificial

<220>  
 <223> pET104/D-TOPO vector

<220>  
 <221> CDS  
 <222> (177) .. (449)

<400> 27  
 ataggcgcca gcaaccgcac ctgtggcgcc ggtgatgccg gccacgatgc gtccggcgta 60  
 gaggatcgag atctcgatcc cgcgaaatta atacgactca ctatagggga attgtgagcg 120  
 gataacaatt cccctctaga aataattttg tttaacttta agaaggagat atacat atg 179  
 Met  
 1  
 ggc gcc ggc acc ccg gtg acc gcc ccg ctg gcg ggc act atc tgg aag 227  
 Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp Lys  
 5 10 15  
 gtg ctg gcc agc gaa ggc cag acg gtg gcc gca ggc gag gtg ctg ctg 275  
 Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu Leu  
 20 25 30  
 att ctg gaa gcc atg aag atg gaa acc gaa atc cgc gcc gcg cag gcc 323  
 Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln Ala  
 35 40 45  
 ggg acc gtg cgc ggt atc gcg gtg aaa gcc ggc gac gcg gtg gcg gtc 371  
 Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala Val  
 50 55 60 65  
 ggc gac acc ctg atg acc ctg gcg ggc tct gga tcc gat ctg tac gac 419  
 Gly Asp Thr Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr Asp  
 70 75 80  
 gat gac gat aag gga att gat ccc ttc acc 449  
 Asp Asp Asp Lys Gly Ile Asp Pro Phe Thr  
 85 90

<210> 28  
 <211> 91  
 <212> PRT  
 <213> Artificial

<220>  
 <223> pET104/D-TOPO vector

<400> 28

Met Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp  
 1 5 10 15

Lys Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu

<210>	29
<211>	450
<212>	DNA
<213>	Artificial

```
<220>
<221> CDS
<222> (160) .. (447)
```

[illegible]



55 60 65

atg acc ctg gcg ggc tct gga tcc gat ctg tac gac gat gac gat aag 414  
Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr Asp Asp Asp Asp Lys  
70 75 80 85

gta cat caa aca agt ttg tac aaa aaa gca ggc tnn 450  
Val His Gln Thr Ser Leu Tyr Lys Lys Ala Gly  
90 95

<210> 30  
<211> 96  
<212> PRT  
<213> Artificial

<220>  
<223> pcDNA/Biotag-DEST vector

<400> 30

Met Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp  
1 5 10 15

Lys Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu  
20 25 30

Leu Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln  
35 40 45

Ala Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala  
50 55 60

Val Gly Asp Thr Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr  
65 70 75 80

Asp Asp Asp Asp Lys Val His Gln Thr Ser Leu Tyr Lys Lys Ala Gly  
85 90 95

<210> 31  
<211> 453  
<212> DNA  
<213> Artificial

<220>  
<223> pcDNA6/Biotag/D-TOPO

<220>  
<221> CDS  
<222> (160)..(453)

<400> 31

cccattgacg caaatgggcg gtaggcgtgt acggtgggag gtctatataa gcagagctct 60

ctggctaact agagaacca ctgcttactg gcttatcgaa attaatacga ctactatag 120

ggagacccaa gctggctagc gtttaaactt aagcttacc atg ggc gcc ggc acc 174  
Met Gly Ala Gly Thr  
1 5

ccg gtg acc gcc ccg ctg gcg ggc act atc tgg aag gtg ctg gcc agc 222  
Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp Lys Val Leu Ala Ser  
10 15 20

gaa ggc cag acg gtg gcc gca ggc gag gtg ctg ctg att ctg gaa gcc 270  
Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu Leu Ile Leu Glu Ala  
25 30 35

atg aag atg gaa acc gaa atc cgc gcc gcg cag gcc ggg acc gtg cgc 318  
Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln Ala Gly Thr Val Arg  
40 45 50

ggc atc gcg gtg aaa gcc ggc gac gcg gtg gcg gtc ggc gac acc ctg 366  
Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala Val Gly Asp Thr Leu  
55 60 65

atg acc ctg gcg ggc tct gga tcc gat ctg tac gac gat gac gat aag 414  
Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr Asp Asp Asp Asp Lys  
70 75 80 85

gta cct agg atc cag tgt ggt gga att gat ccc ttc acc 453  
Val Pro Arg Ile Gln Cys Gly Gly Ile Asp Pro Phe Thr  
90 95

<210> 32

<211> 98

<212> PRT

<213> Artificial

<220>

<223> pcDNA6/Biotag/D-TOPO

<400> 32

Met Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp  
1 5 10 15

Lys Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu  
20 25 30

Leu Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln  
35 40 45

Ala Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala  
50 55 60

Val Gly Asp Thr Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr  
65 70 75 80

Asp Asp Asp Asp Lys Val Pro Arg Ile Gln Cys Gly Gly Ile Asp Pro  
85 90 95

Phe Thr

<210> 33  
<211> 744  
<212> DNA  
<213> Artificial

<220>  
<223> pMT/Biotag-DEST vector

<220>  
<221> CDS  
<222> (454)..(741)

<220>  
<221> misc\_feature  
<222> (743)..(744)  
<223> n is a, c, g, or t

<400> 33  
cgttgcagga caggatgtgg tgcccgatgt gactagctct ttgctgcagg ccgtcctatc 60  
ctctggttcc gataagagac ccagaactcc ggccccccac cgcccaccgc cccccccata 120  
catatgtggt acgcaagtaa gtagtcctgc gcatgccccca tgtgccccac caagagtttt 180  
gcatcccata caagtccccca aagtggagaa ccgaaccaat tcttcgcggg cagaacaaaa 240  
gcttctgcac acgtctccac tcgaatttgg agccggccgg cgtgtgcaaa agaggtgaat 300  
cgaacgaaaag acccgtgtgt aaagccgcgt ttccaaaatg tataaaaccg agagcatctg 360  
gccaatgtgc atcagttgtg gtcagcagca aaatcaagtg aatcatctca gtgcaactaa 420  
aggggggatc tagcgtttaa acitaaagctt acc atg ggc gcc ggc acc ccg gtg 474  
Met Gly Ala Gly Thr Pro Val  
1 5  
acc gcc ccg ctg gcg ggc act atc tgg aag gtg ctg gcc agc gaa ggc 522  
Thr Ala Pro Leu Ala Gly Thr Ile Trp Lys Val Leu Ala Ser Glu Gly  
10 15 20  
cag acg gtg gcc gca ggc gag gtg ctg ctg att ctg gaa gcc atg aag 570  
Gln Thr Val Ala Ala Gly Glu Val Leu Leu Ile Leu Glu Ala Met Lys  
25 30 35  
atg gaa acc gaa atc cgc gcc gcg cag gcc ggg acc gtg cgc ggt atc 618  
Met Glu Thr Glu Ile Arg Ala Ala Gln Ala Gly Thr Val Arg Gly Ile  
40 45 50 55  
gcg gtg aaa gcc ggc gac gcg gtg gcg gtc ggc gac acc ctg atg acc 666  
Ala Val Lys Ala Gly Asp Ala Val Ala Val Gly Asp Thr Leu Met Thr  
60 65 70  
ctg gcg ggc tct gga tcc gat ctg tac gac gat gac gat aag gta cat 714  
Leu Ala Gly Ser Gly Ser Asp Leu Tyr Asp Asp Asp Asp Lys Val His  
75 80 85  
caa aca agt ttg tac aaa aaa gca ggc tnn 744

Gln Thr Ser Leu Tyr Lys Lys Ala Gly  
90 95

<210> 34

<211> 96

<212> PRT

<213> Artificial

<220>

<223> pMT/Biotag-DEST vector

<400> 34

Met Gly Ala Gly Thr Pro Val Thr Ala Pro Leu Ala Gly Thr Ile Trp  
1 5 10 15

Lys Val Leu Ala Ser Glu Gly Gln Thr Val Ala Ala Gly Glu Val Leu  
20 25 30

Leu Ile Leu Glu Ala Met Lys Met Glu Thr Glu Ile Arg Ala Ala Gln  
35 40 45

Ala Gly Thr Val Arg Gly Ile Ala Val Lys Ala Gly Asp Ala Val Ala  
50 55 60

Val Gly Asp Thr Leu Met Thr Leu Ala Gly Ser Gly Ser Asp Leu Tyr  
65 70 75 80

Asp Asp Asp Asp Lys Val His Gln Thr Ser Leu Tyr Lys Lys Ala Gly  
85 90 95